

# Integrating biodiversity assessments into local conservation planning: the importance of assessing suitable data sources

Thibaut Ferraille\*<sup>1,2</sup>, Christian Kerbiriou<sup>3,4</sup>, Charlotte Bigard<sup>5</sup>, Fabien Claireau<sup>3,4</sup>, John D. Thompson<sup>1</sup>

<sup>1</sup> CEFE, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France

<sup>2</sup> Naturalia environnement, Avignon, France

<sup>3</sup> Centre d'Ecologie et des Sciences de la Conservation (CESCO), Muséum national d'Histoire naturelle, Centre National de la Recherche Scientifique, Sorbonne Université, Paris, France

<sup>4</sup> Centre d'Ecologie et des Sciences de la Conservation (CESCO), Muséum national d'Histoire naturelle, Station de Biologie Marine, Concarneau, France

<sup>5</sup> Agence Bretonne de la Biodiversité, Brest, France

\*Corresponding author

Correspondence: [thibaut.ferraille@gmail.com](mailto:thibaut.ferraille@gmail.com)

## ABSTRACT

Strategic Environmental Assessment (SEA) of land-use planning is a fundamental tool to minimize environmental impacts of artificialization. In this context, Systematic Conservation Planning (SCP) tools based on Species Distribution Models (SDM) are frequently used for the elaboration of spatially exhaustive biodiversity diagnostics. Despite the paradigm of “garbage in – garbage out” that emphasises the importance of testing the suitability of data for SDM and priority conservation areas, the assessment of database sources remains relatively rare. In addition, the lack of practical recommendations for the use of open-access databases by SEA stakeholders remains a problem. The aim of this study is to explore the quality of data sources that can be used ~~by stakeholders in SEA~~ to assess priority conservation areas in SEA. The study ~~was done using~~ used data for nine taxonomic groups (commonly used in inventories for environmental impact assessment) and three databases available to SEA stakeholders. Three local administrative entities in very different socio-ecological contexts were used to examine three main issues : (i) the suitability of local versus ~~national regional or country~~ databases for assessing conservation priorities, (ii) differences among taxonomic groups or territories in terms of the suitability of databases, (iii) the importance of the quality of databases for the application of SDM to assess priority conservation areas. Our study provides several clear messages for potential users of open-access databases. First, the need for prudence in the interpretation of biodiversity maps. Second, ~~for SDM,~~ the collection of individual databases at the ~~national country~~ scale is necessary to complete local data and ensure the suitability of ~~models~~SDM in a local context. Third, a data driven approach can lead to the use of notably different species communities to identify priority conservation areas when compared to the community in the original database. Finally, we propose a workflow to guide SEA stakeholders through the process of data rationalization and use in conservation planning.

**Keywords:** Data-driven approach, Species Distribution Models (SDM), Strategic Environmental Assessment (SEA), Systematic Conservation Planning (SCP)

Land-use change, in particular urban land expansion, leads to artificialization of habitats and soils and is one of the major causes of the loss of biodiversity (Maxwell et al., 2016; IPBES, 2019). The reduction and fragmentation of natural habitats leads to population declines and species extinction (Fahrig, 1997; Horváth et al., 2019; Lino et al., 2019), as well as biotic homogenization ~~that, i.e. mostly the extinction of specialist species and the introduction of exotic species, which~~ involves an increase in genetic, taxonomic and functional similarity (Olden and Rooney, 2006; Zambrano et al., 2019) (Olden and Rooney, 2006; Zambrano et al., 2019).

A major tool to limit artificialization is the mitigation hierarchy used in environmental assessments studies. This approach consists of three sequential steps: “avoid” impacts, “reduce or minimize” impacts not avoided and “offset” residual impacts (Bull et al., 2016; Maron et al., 2016). However, in the current application of the mitigation hierarchy several weaknesses prevent it from achieving the goal of “No Net Loss” of biodiversity (Quétier et al., 2014; Bezombes et al., 2019). Avoidance is poorly implemented despite the fact that it is the first and most efficient step of the hierarchy (Bigard et al., 2017; Phalan et al., 2018). What is more, the mitigation hierarchy is mostly applied in a project-by-project approach without scaling up (Pope et al., 2013; Bigard et al., 2017), which limits proper consideration of fragmentation issues (Gontier et al., 2006) and cumulative impacts (Whitehead et al., 2017), including those of multiple small projects (Bigard et al., 2017).

To anticipate avoidance measures, Strategic Environmental Assessment (SEA) of land-use planning is a global and fundamental tool to minimize environmental impacts (Baker et al., 2005). SEA provides for the integration of avoidance measures early in the land-use planning process through environmental assessment of policies, plans and programs (Fundingsland Tetlow and Hanusch, 2012; Bigard et al., 2020). However, the implementation of SEA is often based on biodiversity diagnostic maps that are rarely complete and exhaustive. Indeed, biodiversity diagnostics are rarely exclusively based on empirical observations from field surveys (Phalan et al., 2018) and usually use areas and documents already known (e.g., protected areas and green infrastructures).

Spatial modelling provides a tool for the elaboration of spatially exhaustive diagnostics of biodiversity maps for land-use and conservation planning (Almenar et al., 2019; Tarabon et al., 2019; Bigard et al., 2020; Tulloch et al., 2019; Baker et al., 2021; Boileau et al., 2022). Among these methods of biodiversity modelling, Species Distribution Models (SDM) are widely used to predict suitable habitat for species based empirical observations (Guisan et al., 2017; Zurell et al., 2020) and are increasingly used in conservation planning (Guisan et al., 2017; Domisch et al., 2019; Baker et al., 2021). Systematic Conservation Planning (SCP) tools are also particular pertinent to identify priority biodiversity stakes and avoid the adoption of an *ad hoc* approach (Margules and Pressey, 2000; Pressey and Bottrill, 2008) in order to inform SEA (Tulloch et al., 2019).

The management of databases and their use for conservation planning is a critical issue for the application of such methods to practical conservation planning. The databases available for SEA stakeholders (i.e. decision makers, environmental consultants and conservation managers) are often limited because of data sensitivity or ownership issues, although more and more ~~programmes~~ programs contain data that are publicly available and use of them can be made without any particular attention to their quality (Costello and Wieczorek, 2014; Tittensor et al., 2014) ~~and they are generally unfamiliar to SEA stakeholders~~. Surprisingly however, despite the prevailing recognition of the “garbage in – garbage out” that emphasises the critical importance of the quality of data (Sanders and Saxe, 2017; Canbek, 2022), an examination of data suitability is relatively rare in local conservation planning (Rondinini et al., 2006; Hermoso et al., 2015a). In this context, some authors argue the necessity of examining the sensitivity of model results to the nature of the datasets that are used (Sanders and Saxe, 2017; Clare et al., 2019; Velazco et al., 2020). SDM studies generally use data that has not been designed specifically for this type of analysis, and is often comprised of presence-only data, hence the need for a rigorous assessment of sampling biases (Beck et al., 2014; Botella et al., 2018; Guisan et al., 2017). Another particularly important point that can influence distribution modelling is the spatial extent of the data, and in particular the question of whether to use only local data or those collected on a larger scale (Baker et al., 2021; Meyer, 2007). The choice and possible combination of data sources is part of this problem due to the fact that they often vary considerably in their design, the gradients covered, and potential sampling biases (Fletcher et al., 2019; Boyd et al., 2023). Basically, the use of available databases requires a rigorous test of their quality and pertinence (Zuckerberg et al., 2011), especially when used for analyses such as SDM (Tulloch et al., 2016; Domisch et al., 2019). Confidence in the models must be assessed through the use of metrics adapted to the data (Guisan et al., 2017; Leroy et al., 2018). As recognised by Clare et al., (2019), the lack of practical

98 recommendations for the use of databases that differ in terms of their quality and pertinence by public  
99 authorities or other institutions remains a serious problem.


100 The overall goal of this study is to test the influence of different database sources that can be used by SEA  
101 stakeholders to map priority conservation areas in SEAs based on SCP. To do so, we studied three local  
102 administrative territories that occur in different socio-ecological contexts in France. The study has three main  
103 objectives. First, ~~in terms of the quantity of data available we assess the content of three open-access~~  
104 ~~databases~~ for nine taxonomic groups commonly used in naturalist inventories in environmental ~~impact~~  
105 ~~assessment studies, we assess the variation in the . We evaluate their suitability of the available databases in~~  
106 ~~terms of data quantity for SDM application, at three scales (local, regional and national) for application of SDM.~~  
107 ~~Second, by focusing on country). SDM and SCP analyses were performed for two taxonomic groups with~~  
108 ~~different dispersion capacities, home range sizes and environmental(Aves and Papilionidae) to test the~~  
109 ~~hypothesis that sampling bias and differences in ecological response scales, of species may influence the~~  
110 ~~identification priority conservation areas. Second, we explore the influence of databases for on the application~~  
111 ~~of SDM to assess priority conservation areas. Third, we analyse the influence of this data-driven approach on~~  
112 ~~the composition of species communities that are ultimately considered in priority conservation areas relative~~  
113 ~~to the actual communities in the original databases.~~ Third, we analyse the influence of this data-driven  
114 approach on the composition of species communities that are ultimately considered used in the  
115 analysis identification of priority conservation areas relative to the actual communities in the original  
116 databases.

## 117 2 - Methods

### 118 2.1 - Study sites

119 To assess the availability and suitability of pertinent data sources, we selected three French local  
120 administrative entities in charge of land-use planning: Lodévois-Larzac (T1), Brocéliande (T2), La Rochelle (T3)  
121 ~~(Figure 1, Table 1). We selected these territories). We selected these study sites~~ on the basis their contrasting  
122 social, ecological and geographical contexts in order to examine patterns of variation of data suitability among  
123 sites- ~~(Figure 1, Table 1). For example, each of the three territories is composed of study sites have~~ different  
124 ecosystems and bioclimates and the sites vary in terms of urbanization pressures ~~(from 3% to 28% artificial~~  
125 ~~land-use cover for sites T1 and T2 respectively) and . This is due to the presence of protected areas a major~~  
126 ~~city (La Rochelle) in their territories (from 2% to 70% of the protected territory the T2 study site. The major~~  
127 ~~towns in the other two study sites are smaller however there is a major city less than 50 km away for T3 both~~  
128 ~~of them (Montpellier and T1 Rennes respectively). Only site T1 has an important cover of protected areas~~  
129 (Natura 2000) area.

131



132  
133  
134  
135

**Figure 1 - Localisation** of the study sites in **France** administrative regions: T1 is **Brocéliande**, T2 is La Rochelle, T3 is **Lodévois-Larzac-Brocéliande**. Source: IGN, Google, 2023.

**Table 1** - Description of three study sites in France

<u>Territory</u> / <u>Study site</u>	<u>Lodévois-Larzac (T1)</u>	<u>La Rochelle (T2)</u>	<u>Brocéliande (T3)</u>	
Country / <u>Region</u> / <u>Administrative region</u>	France / Occitanie	France / Nouvelle-Aquitaine	France / Bretagne	
Main city coordinates	43°43'57"N 3°19'02"E	46°09'35"N 1°09'05"W	48°00'08"N 2°05'48"W	
Area	554 km <sup>2</sup>	327 km <sup>2</sup>	298 km <sup>2</sup>	
Climate	Mediterranean	Oceanic	Oceanic	
Major habitats / land-use	Mediterranean forest and scrubland	Intensive agricultural field, swamps, few forests	Pastures, temperate forest	
% land-use <sup>(1)</sup>	Artificial – <u>urban areas</u>	28%	7%	
	Farmland	55%	29%	
	Semi-natural	17%	64%	
Landscape and dynamics	Rural, woodland expansion	Coastal, intensive agriculture, urban expansion	Rural, agricultural intensification	
Urbanization <u>context</u>	Close to a major city, <del>across</del> <u>(Montpellier), linked by a highway</u>	Economic and tourist dynamism, <u>major city presence</u>	Close to a major city, <del>across</del> <u>(Rennes), linked by a highway</u>	
% protected area <sup>(2)</sup>	Regulatory	0%	0%	
	Land control	1618 ha – 0.6%	148 ha – 0.45%	80 ha – 0.25%
	Contractual	38096 ha – 68.8%	3534 ha – 10.7%	586 ha – 2%
	Total	38602 ha – 69.7%	3546 ha – 10.8%	592 ha – 2%

(1) source: OSO Land Cover (Inglada et al., 2019). Artificial is impermeable surfaces; crop field is annual crop and orchards and vineyards; semi-natural is hardwood and soft wood and grasslands and water.

(2) source: <https://inpn.mnhn.fr/>

## 138 2.2 – Workflow of analysis

139 A methodological framework was developed to test the influence of different database sources in mapping  
140 priority conservation areas in SEAs thanks to a SCP approach, all steps are summarized in Figure 2- and in the  
141 following text.

## 142 2.3 – Databases available for SEA stakeholders

### 143 2.3.1 – Content of available databases

144 We focused on nine taxonomic groups commonly used for naturalist inventories for environmental impact  
145 assessment studies: Amphibia, nesting Aves (hereafter name Aves), Chiroptera, Flora, Mammalia aptera  
146 (hereafter name Mammalia), Orthoptera, Odonata, Papilionidae and Reptilia (Bigard et al., 2017; Guillet et al.,  
147 2019; Iorio et al., 2022). The three spatial scales used for data collection are depicted in Figure 1: the local scale  
148 (i.e. study site with a 10km buffer around), the regional scale (i.e. French administrative regions); and the  
149 country scale (i.e. continental France). We selected three open-access databases containing these groups that  
150 can be widely used by SEA stakeholders for the assessment and hierarchy of conservation priorities (Figure 2,  
151 step 1.1). This study thus directly addresses SEA stakeholders (i.e. decision makers, environmental consultants  
152 and conservation managers) using the databases available to them.

153 The first of these databases concerns the French Natural and Landscape Information System (SINP) that is  
154 structured at the regional-scale in France-scale of French administrative regions in charge of data extraction  
155 requests. Each site has its own database, can be collected only at local scale due to the limited extent of data  
156 requests, without the need for a special request (maximum 2000 km<sup>2</sup>, i.e. nearby 10km buffer zone around  
157 the study site). This database is composed of opportunist observations and only contains presence data for  
158 taxa for which identification is confirmed by experts (Jomier et al., 2018). Access to this database requires a  
159 data extraction request to the regional administrative structure in charge of the database. Generally, a special  
160 request is necessary above a 10km buffer zone around the study site (maximum 2000 km<sup>2</sup>), the data remains  
161 accessible only locally. Each site has its own database (see Appendix- (see Appendix A.1).

162 The second database is the Global Biodiversity Information Facility (GBIF) an international platform for the  
163 provision of biodiversity data that is based on information collected from various databases (Telenius, 2011).  
164 It is composed of observation data that are not based on protocols and for which presence data and



165 identification are not subject to expert confirmation. The data downloading is autonomous from the website  
166 (see Appendix A.1).

167 The third database is a French biodiversity monitoring scheme (Vigie Nature) dedicated to assess spatio-  
168 temporal populations trends. Within this monitoring scheme data collection is based on a standardized  
169 biodiversity survey. Despite local spatial distribution heterogeneities, the sampling plan ensures a  
170 representation of the current ~~national~~ distribution of habitats and landscapes across France (Julliard and  
171 Jiguet, 2002; Mariton et al., 2022). Homogeneity in identification criteria and compliance with the protocol are  
172 ensured by offering training to volunteers. This database is composed of presence/absence and abundance  
173 data. Access to these databases requires a data extraction request to the person in charge (see Appendix A.1).

174 These three databases were combined in two ways: “All databases” (i.e. the combination of SINP, GBIF and  
175 Vigie Nature) used in section 2.3 (i.e. for assess which taxa are enough documented within each dataset) and  
176 “National databases” (i.e. the combination of GBIF and Vigie Nature, which are available at country scale) used  
177 specifically in section 2.4 (i.e. ~~for to~~ test the effect of database sources on SDM performance).

178 ~~The databases were collected for continental France except for the SINP that was collected in 10km buffer~~  
179 ~~zone around the study sites due to the access restrictions explained above. The databases were collected over~~  
180 ~~a period of 10 years (i.e. from 01/01/2010 to 31/12/2020) and data with spatial inaccuracy greater than 50~~  
181 ~~meters were not considered. We made a series of operations to standardize, correct and homogenize species~~  
182 ~~names, transform data into occurrences, limit their spatial biases and identify the presence of species into~~  
183 ~~each study site (Appendix A.1).~~

184 The databases were collected for continental France except for the SINP that was collected in 10km buffer  
185 zone around the study sites due to the access restrictions explained above. The databases were collected over  
186 a period of 11 years (i.e. from 01/01/2010 to 31/12/2020) and data with spatial inaccuracy greater than 50  
187 meters were not considered. We made a series of operations to standardize, correct and homogenize taxa  
188 names at the specific taxonomic level using the French taxonomic reference “TAXREF.V14” (Gargominy et al.,  
189 2021). We transformed data into occurrences and limited their sampling biases by geographical filtering using  
190 “spThin” package (Aiello-Lammens et al., 2015). In each study site, we identify the presence of one species at  
191 least five observations from “All databases” combined at local scale and defined as present species in France  
192 by TAXREF.V14 (Appendix A.1).

### 193 2.3.2 – Quantitative description of databases for SDM

194 For the nine taxonomic groups, four metrics were selected to quantitatively describe the amount and thus  
195 the suitability of each database (i.e. SINP, GBIF, Vigie Nature and “All databases”) for the realization of SDM  
196 in presence-only (Figure 2, step 1.2): (i) the number of species observed in the study site; (ii) the proportion of  
197 species with < 15 observations which represents the minimal threshold for the utility of SDM with more  
198 accurate predictions than in a random model (Støa et al., 2019); (iii) the proportion of species with between  
199 15 and 50 observations, i.e. the minimum number of presences recommended for SDM (Merow et al., 2014 in  
200 Guisan et al., 2017); (iv) the proportion of species with > 50 observations, i.e. highly suitable for modelling  
201 (Støa et al., 2019). These metrics were calculated at three different scales: (a) for each ~~territory~~study site  
202 including a 10km buffer zone (local scale) that is the maximum extent for a SINP data request; (b) on a regional  
203 scale that is used for the structure of biodiversity data in France; (c) for continental France. SINP database is  
204 analysed individually only at the local scale due to the previously mentioned restriction of access, nevertheless  
205 it integrated the three scales of the “All databases:”.

## 206 2.4 - Systematic conservation planning process

207 For the three study sites, we identified priority conservation areas with a Systematic Conservation Planning  
208 (SCP) tool based on Species Distribution Models (SDM) (Figure 2, steps 2.1 and 2.2). Several variants of SDM  
209 were made using different database sources (i.e. GBIF, Vigie Nature and “National databases”) and two  
210 methods of generating pseudo-absences (named individual database or mixed databases). Among the nine  
211 taxonomic groups studied above, only Aves and Papilionidae taxa were analysed to compare the tests in the  
212 SCP process. The data available in France for these groups ~~seems to be~~ sufficient in quantity to realize SDM  
213 with each database. The use of these two groups allows for a comparison between one group of highly mobile  
214 taxa with a large home range (Aves) and another group with a smaller home range and whose movement  
215 closely tracks local environmental variation (Papilionidae). These two taxonomic groups thus have different

216 biological traits associated with their dispersal and function, hence we predict differences in in terms of the  
217 spatial resolution of their distribution.

#### 218 2.4.1 - Species distribution modelling (SDM)

219 We modelled favourable habitats for birds and butterflies in the three study sites using SDM (Figure 2, step  
220 2.1). A resolution of 50m was used to meet the needs of the SEA of land-use planning. A buffer zone of 10km  
221 around each of the study sites (i.e. local scale) was used for the SDM prediction to limit any border edge effects  
222 and to increase the number of species that could be modelled and evaluated. Indeed, species with less than  
223 15 data points for the calibration (threshold explain above, Støa et al., 2019) and/or less than 10 data for  
224 performance evaluation (threshold defined by expert opinion) were not modelling.

225 Biodiversity data used for the SDM came from the databases ~~described~~described above at the  
226 national/country scale, according to the results of section 2.3 (Table 2). These data were separated into two  
227 independent datasets that allow for robust validations with independent data (Matutini et al., 2021).  
228 National/Country data without local data were used for model calibration and local data were used only for  
229 model performance evaluation. Therefore, the SINP database, which is only available at local scale, was not  
230 used for model calibration.

231 For model calibration, pseudo-absences were generated with two methods: separately for each taxonomic  
232 group. First, methods to generate pseudo-absences in the individual databases for all their data (i.e. GBIF, Vigie  
233 Nature and “National databases”). For databases with a protocol for sampling (i.e. Vigie Nature) to optimize  
234 species detection (day and year periods), the absence points were defined as all the points without the  
235 observed species. For databases without such sampling protocols for all their data (i.e. GBIF and “National  
236 databases”), pseudo-absence data were generated with the target-group (TG) approach, which infers the  
237 sampling bias from the aggregated occurrences of (TG) species, i.e. the respective taxonomic groups (Ponder  
238 et al., 2001; Anderson, 2003; Phillips et al., 2009). Second, a method to mix the presence data in the GBIF and  
239 “National databases” with the absence from Vigie Nature (named mixed databases) was applied (Hermoso et  
240 al., 2015a).

241 Three types of environmental variables were used for SDM: biogeographic/geographic, human occupancy  
242 and pollution and fragmentation (Appendix A.2A.2).

243 SDM were calibrate/calibrated by Random Forest down-sample (Valavi et al., 2021a) which according to  
244 Valavi et al. (2021b) is among the best performing models for presence-only data. Although the Random Forest  
245 is not very sensitive to the non-independence of the variables and over-parametrization (Matsuki et al., 2016;  
246 Srisa-An, 2021), in order to be parsimonious, the collinear variables were removed (Pearson >0.7, Appendix  
247 A.2A.2, Brun et al., 2020). Thirty bootstraps were performed for each SDM (Guisan et al., 2017) using a  
248 calibration for 70% of the data at the national/country scale outside of local scale. The thirty Random Forest  
249 bootstraps were combined with mean to provide an ensemble prediction of habitat suitability for all species.

250 The performance evaluation of the models was done using the Boyce’s index (CBI), the most suitable metric  
251 for model in presence-only (Boyce et al., 2002; Leroy et al., 2018), with local data (i.e. “All databases”  
252 combining SINP, GBIF and Vigie Nature). Dubos et al. (2022) reveal the CBI turns out to be misleading in some  
253 cases, thus we used a threshold of 0.3 to define good or poor model quality.

#### 254 2.4.2 – Systematic conservation planning (SCP) tool

255 Priority conservation areas in the three territories/study sites were analysed from SDMs for each database  
256 source in using a SCP tool (Figure 2, step 3.1). To meet the needs of SEAs, the study site was restricted to  
257 administrative boundaries with a buffer zone of 1km to maintain coherence between administrative entities  
258 and a resolution of 50m. The SEAs/aim of SEA biodiversity conservation strategies aim is to establish priorities  
259 for the whole territory/study site as a whole and all the cells have the same cost value of 1. The objective was  
260 a maximum coverage objective that seeks to maximize the number of features, i.e. the SDMs (Church et al.,  
261 1996). The features were only the predictions of SDM with a Boyce’s index > 0.3 to limit the influence of  
262 poor/good quality models defined previously. The priority conservation areas decisions were between 0 and 1.  
263 To obtain a priority gradient, we cumulated Ferrier importance scores (Ferrier et al., 2000) from nine targets  
264 of the total amount of each feature (from 0.1 to 0.9 every 0.1). ~~The~~We used the package “prioritizr” ~~was used~~  
265 (Hanson et al., 2021) with the open-source solver SYMPHONY (Kim et al., 2023).

## 2.5 - Comparative analysis of SDM and priority conservation areas

We analysed the influence of database sources on SDM predictions and priority conservation areas were analysed (Figure 2, step 2.2 and 3.2). SDM performance evaluations were analysed between database sources. The SDM prediction and priority conservation areas maps were compared with the Spearman's rank coefficient (Phillips et al., 2009) and the Schoener's D index as a measure of projection overlap (Schoener, 1968) which was calculated with the ENMTool R package (Warren et al., 2008; Warren and Dinnage, 2022).

## 2.6 - Species community analysis

We assessed the influence of the complete data driven workflow on the composition of species communities, i.e. differences between the original community (i.e. all species observed in study site) in the database and the final community used to identify priority conservation areas (Figure 2, step 4).

To do so, we developed a workflow resistance score for each of the methodological filters for all species. A score of 1 is allocated to species observed in the study area that did not cross any of the stepwise filters. A score of 2 is allocated to species with sufficient data to calibrate SDM, i.e. > 15 national/country observations, or evaluate to evaluate the performance of SDM, i.e. > 10 local observations. A score of 3 is allocated to species with sufficient data to calibrate and evaluate the performance of the SDM. A score of 4 is for species that were present in the final analysis as a priority species for conservation planning (i.e. with the two previous filters and a Boyce's index > 0.3). In order to assess species composition bias ultimately considered in priority conservation areas, species communities were analysed through traits that can influence species detection (mass, displacement capacity, period activity) and ecological traits (habitats, specialisation) (Appendix A.3). Missing data were completed with a trait imputation procedure generated using the R package "missForest" (Stekhoven and Bühlmann, 2012) by considering evolutionary relationships in the imputation process (see Carmona et al., 2021) using the R script of Toussaint et al. (2021). Due to the nature of the response variable (i.e. ordinal scoring including four modalities), we used ordinal regression mixed models with cumulative link using the clmm function of "ordinal" R package (Christensen, 2022). We adapted the link function to the data distribution for Aves and Papilionidae, using the "cauchit" link for Aves and a "logit" link, respectively link for Papilionidae. Species traits were used as fixed effects, while the random effects selected were the study sites for Aves and the combination for study sites and database sources for Papilionidae. Finally, we evaluated the quality of the full model by comparing to the null model with Akaike's information criterion (AIC) (Mac Nally et al., 2018).


Thus, our models were structured in the following way:

$$\begin{array}{ll} \text{For Aves} & \text{Resistance} \sim \beta_0 + \beta_i \cdot \text{trait}'_i + (1|\text{StudySites}) \\ \text{For Papilionidae} & \text{Resistance} \sim \beta_0 + \beta_i \cdot \text{traits}'_i + (1|\text{StudySites}/\text{Databases}) \end{array}$$


Where  $\beta$  is the parameter estimates,  $i$  correspond to the variables of 'traits' using in fix effect, 0 is the shift between ordinal class of resistance (i.e. 1|2, 2|3 and 3|4) and "1|" is the random effect.




## Workflow for mapping priority conservation areas



## Analysis





**Figure 2** - Methodological framework applied to test the influence of different database sources to map conservation planning areas in SEAs based on SCP. The green and blue boxes are the appendices detailing the methodology and the results of the analyses, respectively. The red boxes "R." are the methodological filters used for generate workflow resistance score.

301  
302  
303  
304  
305  
306

### 3.1 - Quantitative description of databases for SDM

The use of individual observation databases provided a limited number of observation data for distribution modelling of many species, regardless of the taxonomic group (Table 2, Appendix B.1)-B.1). None of the GBIF and SINP individual databases have more than 50 observations for all nine taxonomic groups, and there are currently no programs for three of the nine taxonomic groups in the Vigie Nature databases. Local databases and, to a lesser extent, regional databases are not equivalent in terms of the amount of data available for different taxa for the three study sites. The difference is particularly pronounced for the SINP databases, with no data available for five taxonomic groups in the T3 study site, whereas in the T1 study site, six taxonomic groups have sufficient data to model over 50% of the species. At the regional scale, the number of observations per species is highly variable among [territoriesstudy sites](#) for the GBIF databases and is more similar for the Vigie Nature databases.

The use of combined databases (i.e. ["All databases"](#)) increased the total number of species that can be studied. At the local scale, the proportion of species in each suitability class showed only small changes, while at the [nationalcountry](#) scale of France, it allowed a significant gain in species with suitable data (Table 2). Indeed, on a [nationalcountry](#) scale, the GBIF and Vigie Nature databases are complementary with each other. For example, the GBIF has few Chiroptera data, which is complemented by Vigie Nature data, and vice versa for Amphibian data.

At the [nationalcountry](#) scale, aggregation of the databases seem to provide the most suitable setup (databases and scale) for SDM analysis. Using these compiled, [nationalcountry](#) databases provides a large amount of data for a large number of species present in the three [territoriesstudy sites](#) (Table 2 and Appendix B.1).

### 3.2 - SDM and priority conservation areas analysis

The evaluation of SDM revealed differences among the database sources; none of which produced more than 87% of satisfactory models for the two studied taxonomic groups and some had less than 20% of satisfactory models (Table 3, Appendix B.2). Use of the GBIF data led to a higher proportion of well-evaluated SDM, ranging between 48 and 79% of satisfactory models for the species in the two taxonomic groups. GBIF data are also more suitable than Vigie Nature data, they produced between 11% and 37% more satisfactory models than the latter database (Table 3). Nevertheless, between 4% and 9% of species provide well-evaluated models from Vigie Nature and poorly evaluated by GBIF database. The combination of ["National databases"](#) (i.e. GBIF and Vigie Nature) decreased the performance of SDMs with GBIF data, but still yield better results than SDM based on the Vigie Nature database. The substitution in individual databases of pseudo-absences for the absences from Vigie Nature (i.e. mixed databases) reduced the performance of models based on GBIF, but increased the performance of the combined ["National databases"](#) (Figure 3, Appendix B.2)-B.2). Regardless of the database used, our analyses revealed significant differences between [territoriesstudy sites](#) (Table 3). No SDMs for butterflies in T2 could be evaluated due to insufficient local data. For the T3 study site, over 50% of the Papilionidae and Aves models perform poorly, whereas for the T1 study site poor models occur in less than 50% of the evaluations.

Although important differences in model performance between the database sources used for SDM showed a high degree of overlap, as indicated by Shoener's D index with values above 0.8, the ranking of habitat suitability was highly variable. This was in particularly the case for the GBIF and Vigie Nature databases that had median spearman's rank coefficient values between 0.2 and 0.5 and a very wide distribution (Appendix B.3). The substitution of pseudo-absence data in GBIF and ["National databases"](#) by absence data from Vigie Nature, showed a similar situation (Appendix B.4).

For priority conservation areas, whatever the individual databases used, the overlaps with Schoener's D index were above 0.72 and similar in each [territorystudy site](#). Nevertheless, Spearman's rank coefficients showed a greater difference in prioritization ranks in particular between GBIF and Vigie Nature and for the T3 study site (Table 4). Between maps of priority conservation areas, we observed similarities in overlap, despite a significant difference in the hierarchy of areas to be prioritized (Figure 4, Appendix B.5). [The list of species is presented in Appendix B.7, where it can be seen that there are no difficulties with respect to invasive species which are very few in the data sets.](#)

358 **3.3 - Species community analysis**

359 The distribution of workflow resistance scores showed that only 30% and 42% respectively of Aves and  
360 Papilionidae species were integrated in priority conservation areas maps. Among species not integrated, the  
361 workflow steps filtering the most species concern the amount of suitable data for model evaluation followed  
362 by the quality of the models (Figure 5). The analysis of the species community composition observed in each  
363 of the three study sites in comparison with the species community integrated in priority conservation  
364 areas area identification revealed significant differences for all three study sites. For Aves communities, the  
365 differences concern an under-representation of nocturnal species, large species with high dispersal capacity,  
366 and species of swamp habitats and deciduous forests in relation to the observed species community in the  
367 databases for the three study sites. Conversely, species that favour urban habitats, shrubland, grassland and  
368 coniferous forests are over-represented in the species community of the final maps, as are species with  
369 specialized diets and foraging strata (Table 5). For Papilionidae communities, common species with long flight  
370 periods are over-represented in the final community used for analysis. Species related to anthropogenic and  
371 thermo/meso Mediterranean habitats, and species that use a wide range of hostplants are over-represented  
372 in relation to the original species community, while supra-Mediterranean species and those of montane  
373 environments are under-represented (Table 5).

374  
375  
376  
377

**Table 2** - Quantitative description of the observation databases (Local SINP, GBIF, Vigie Nature) for nine taxonomic groups in the three study sites ("T1": Lodévois-Larzac, "T2": La Rochelle, "T3": Brocéliande) for "local" (study site with buffer of 10km), "Regional" (administrative region) and "FranceCountry" (continental France). "nSp" is the number of species observed at least 3 times in a given territory/study site within a 10 km buffer zone, ">15obs" is the percentage of species with more than 15 observation and ">50obs" the percentage with more than 50 observations (red < 25%, orange 25-75%, green >75%, grey - no data).

Taxa	GBIF									SINP									Vigie Nature									All databases combined												
	Local			Regional			FranceCountry			Local			Regional			FranceCountry			Local			Regional			FranceCountry			Local			Regional			FranceCountry						
	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3				
Amp	nSp	11	1	11	13	2	11	14	2	11	13	2	0	No program									14	2	11	14	2	11	14	2	11	14	2	11	14	2	11	14	2	11
	>15obs	[red]			[orange]			[green]			[grey]			[grey]									[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[grey]									[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Ave	nSp	100	73	77	198	159	120	204	170	131	197	94	82	114	155	101	182	143	120	201	170	132	205	171	132	205	166	132	205	171	132	205	171	132	205	171	132			
	>15obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Chi	nSp	4	0	5	13	13	5	19	19	21	21	6	18	25	22	24	25	22	24	25	22	25	26	22	26	26	22	26	26	22	26	26	22	26	26	22	26			
	>15obs	[red]			[red]			[red]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Flo	nSp	786	476	639	1428	1093	825	1573	1261	995	1703	1335	843	286	177	136	825	657	413	1336	1105	848	1844	1460	1093	1844	1158	1093	1844	1460	1093	1844	1460	1093						
	>15obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Mam	nSp	10	1	14	23	6	21	23	6	28	23	7	30	No program									23	7	30	23	6	30	23	7	30	23	7	30	23	7	30			
	>15obs	[red]			[orange]			[green]			[grey]			[grey]									[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[grey]									[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Odo	nSp	38	5	23	56	7	28	57	7	30	57	1	0	0	1	24	56	7	27	57	7	30	57	7	30	57	7	30	57	7	30	57	7	30	57	7	30			
	>15obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Ort	nSp	53	3	24	93	25	26	95	25	34	91	8	0	25	19	19	25	19	19	25	19	19	98	25	34	98	25	34	98	25	34	98	25	34	98	25	34			
	>15obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Pap	nSp	115	15	33	144	18	43	147	18	52	154	11	0	0	0	50	132	18	51	149	18	52	154	18	52	154	18	52	154	18	52	154	18	52						
	>15obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
Rep	nSp	18	2	7	20	4	7	21	4	7	21	4	0	No program									22	4	7	22	4	7	22	4	7	22	4	7	22	4	7			
	>15obs	[red]			[orange]			[green]			[grey]			[grey]									[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		
	>50obs	[red]			[orange]			[green]			[grey]			[grey]									[orange]			[orange]			[orange]			[orange]			[orange]			[orange]		

378


**Table 3** - Proportion of well-evaluated SDM corresponding to Boyce's index greater than 0.3. "National" combines GBIF and Vigie Nature databases.

Taxa	Study site	Number of species	Individual database			SDM with Vigie Nature absence	
			GBIF	Vigie Nature	"National"	GBIF	"National"
Aves	T1	114	79 %	61 %	70 %	87 %	82 %
	T2	87	68 %	31 %	47 %	57 %	46 %
	T3	56	48 %	38 %	29 %	52 %	43 %
Papilionidae	T1	127	75 %	52 %	64 %	75 %	73 %
	T3	25	52 %	16 %	20 %	12 %	20 %

379



380



381

382

383

384

385

386

387


388

**Figure 3** - Proportion of SDM with Boyce's index (CBI) greater than 0.3 by database source ("Individual" and "Mixed" with Vigie Nature absence), by combining territories/study sites (T1, T2, T3) and taxonomic groups (Aves, Papilionidae). "National" combines GBIF and Vigie Nature databases.

**Table 4** - Overlap of priority conservation areas between database sources using two metrics: Schoener's D index (D) and Spearman's rank coefficient (S cor). VN is Vigie Nature database and "National" combines GBIF and Vigie Nature databases.

Taxa	Study sites	Individual database						Pseudo-absence of individual database - absence from VN			
		"National" - GBIF		"National" - VN		GBIF - VN		GBIF		"National"	
		D	S cor	D	S cor	D	S cor	D	S cor	D	S cor
Aves	T1	0.82	0.56	0.81	0.53	0.78	0.36	0.75	0.88	0.79	0.38
	T2	0.82	0.74	0.76	0.49	0.72	0.42	0.76	0.64	0.79	0.64
	T3	0.77	0.34	0.81	0.51	0.78	0.29	0.73	0.19	0.79	0.39
Papilionidae	T1	0.85	0.64	0.84	0.65	0.83	0.59	0.72	0.66	0.82	0.66
	T3	0.80	0.49	0.77	0.31	0.78	0.22	0.76	0.17	0.72	0.06

389




390

391

392

**Figure 4** - Maps of priority conservation areas for Aves and Papilionidae of tree study sites (T1, T2, T3) from different individual database source (All, GBIF, Vigie Nature).



**Figure 5** - Workflow resistance scores for Papilionidae and Aves. 1 - species observed in the study area that did not cross any of the stepwise filters. 2 - species with sufficient data to calibrate SDM, i.e. > 15 national/country observations, or evaluate the performance of SDM, i.e. > 10 local observations. 3 - species with sufficient data to calibrate and evaluate the performance of the SDM. 4 - species present in the final analysis, i.e. Boyce's index > 0.3.

**Table 5** - Parameter estimates ( $\beta$ ), standard error (se) and P-values for the full model of Aves and Papilionidae species resistance to the workflow. Appendix B.6, the evaluation of the quality of the model.

	Variables	Aves			Papilionidae			
		$\beta$	se	P-value	$\beta$	se	P-value	
Full model	$\beta_0$ 1 2	-7.84	1.12	/	$\beta_0$ 1 2	-2.54	1.28	/
	$\beta_0$ 2 3	0.14	0.33	/	$\beta_0$ 2 3	1.39	1.26	/
	$\beta_0$ 3 4	1.4	0.34	/	$\beta_0$ 3 4	2.97	1.27	/
	Mass	-0.21	0.13	.	WingspanM	0.01	0.09	
	Avian hand-wing index	-0.31	0.07	***	FMoMean	0.38	0.11	***
	Nocturn	-2.28	0.59	***	Hostplant N	0.23	0.11	*
	Deciduous	-0.27	0.15	.	Hostplant Spe	0.16	0.11	
	Coniferous	0.38	0.15	*	HPG Bi	0.28	0.50	
	Woodland	0.17	0.14		HPG Th	-0.21	0.22	
	Shrub	0.55	0.13	***	HPG Sb	-0.32	0.23	
	Grassland	0.47	0.13	**	HPG Tr	0.22	0.45	
	Mountain meadows	0.23	0.24		HPG Li	0.02	0.36	
	Reed	0.08	0.33		SSI	0	0	
	Swamps	-0.84	0.28	**	AltVeg 'A'	0.06	1.12	
	Rocks	-0.14	0.21		AltVeg 'Mo'	-0.93	0.31	**
	Urban	0.83	0.14	***	AltVeg 'SupMed'	-0.79	0.46	.
	Spe. Diet	0.92	0.35	**	AltVeg 'ThMeMed'	1.37	0.37	***
	Spe. Foraging behav.	-0.08	0.4		Rarity '2'	0.86	0.29	**
	Spe. Diet strat	1.49	0.4	***	Rarity '3'	1.42	0.34	***
	Spe. Habitat	-0.14	0.38					
	Spe. Nest	-0.28	0.72					
	Spe. Mean	-2.26	1.63					

P-value: \*\*\* P < 0.001, \*\* P < 0.01, \* P < 0.05, . P < 0.1, / P-value not applicable

Trait description: Spe. is the specialization, FMoMean is duration of yearly flight period, Hostplant N is hostplant specificity, HostplantSpe is hostplant specificity index, HPG is hostplant growth form (Bi: short herb, Th: tall herb, Sb: shrub, Tr: tree, Li: Liana), SSI is Species Specialization Index, AltVeg is altitudinal vegetation (A=Anthropogenic, Co: Foothill, TheMeMed: Thermo/Meso-Mediterranean, Med: Mediterranean, SupMed: Supra-Mediterranean, Mo: Montane, Asa: Alpine and Subalpine). For more details on traits see Appendix A.3

393

394

395

396

397

398

399

400

401

402

403

404

405

406

The absence of recommendations for the use of available databases that differ in terms of their quality and pertinence by public authorities or other institutions remains a serious problem for local conservation planning (Clare et al. 2019). The goal of this study was to test the suitability of different database sources that can be used by public stakeholders to map priorities for biodiversity stakes in SEAs and SCP. We found that the compilation of databases at the national/country scale is the most suitable procedure to apply SDM to a large number of species. For Aves and Papilionidae, the GBIF database provided the highest proportion of well-assessed SDM. We detected a significant overlap in species distributions in different database sources despite significant variability in the order of habitat suitability and similar spatial predictions for priority conservation areas. Finally, we showed that the composition of the species community used for priority conservation areas in all three study sites were clearly not representative of the observed species communities in the original database (in terms of ecological traits)-species and ecological traits). Finally, despite important differences among the study sites in terms of the proportion of artificial land cover and protected areas we found no particular differences between the three study sites. Clearly, the data sources are the most important factor influencing the results.

Open-access biodiversity data provide a valuable source of information for decision makers, environmental consultants and conservation and land-use planning managers (i.e. SEA stakeholders); they contain vital information on species locations compared to expert knowledge and unshared datasets that are inaccessible to most users (Sousa-Baena et al., 2014; Meyer et al., 2015). Through their use and careful application of SDM, they contribute to the estimation of relative habitat suitability in a given study site (Baker et al., 2021). Our study showed however that SDM for a large majority of species observed locally requires their compilation on a national/country scale. Local and regional data are not suitable for model calibration but remain important for assessing the suitability of models in a local context. Indeed, this performance evaluation step is one of the most restrictive filters in the workflow we proposed, as evidenced by SDM's for butterflies in the T2 study site, where no species could be evaluted/evaluated. The spatial extent of data collection can influence distribution modelling (Meyer, 2007), and our study emphasises this importance for SDM and the use of data available on national/country scale in France.

Different types of databases are constructed in different way – with opportunistic data collection or, in some cases, as part of a scientific monitoring scheme – allowing the use of as complementary data sources (Beck et al., 2013; Shirey et al., 2021). The three individual databases examined in this study are indeed complementary in that when they are combined they provide a suitable source for modelling the distribution of many species. Nevertheless, some groups commonly have a low number/amount of data in such bases, e.g. Insecta (Troudet et al., 2017). The study of data with a fine spatial grain, as required for SEA and SCP (Guisan et al., 2013), reveals information gaps for more taxonomic groups at the global scale, e.g. Amphibians and Mammals (Witté and Touroult, 2017). The construction of an overall database at national/country scale is therefore the most appropriate way to have suitable data for SDM of different taxonomic groups. Furthermore, there is a dilemma between protocolized and opportunistic data. Although protocolized data are recommended for SDM (Guisan et al., 2017; Guillera-Arroita et al., 2015), very often the amount of such data is low, which can be detrimental at the local scale, particularly for model evaluation with data having the same sampling bias. For opportunistic data, their large number is of course a positive point, however the estimating their sampling bias can be a real challenge (Botella et al., 2018; Fithian et al., 2015; Matutini et al., 2021) to ensure the reliability of the results.

SDM of Aves and Papilionidae species clearly revealed differences between the databases used for modelling, with differential impacts on the identification of conservation priorities. Indeed, the high overlap in species distribution overlap is high between database/data sources, as indicated by the Schoener's/Schoener's D index, i.e. indicates that, regardless of the data source, species, are predicted in similar environments (Warren et al., 2008). However, Spearman's ranking of habitat suitability between data sources was highly variable, indicative that species' responses to environments are highly variable, as are the location of favourable habitats (Warren et al., 2008). Although the use of presence-absence data is advocated for SDM (Guillera-Arroita et al., 2015; Valavi et al., 2021b; Dubos et al., 2022)(Guillera-Arroita et al., 2015; Valavi et al., 2021b; Dubos et al., 2022), we showed that opportunist data from GBIF provided a greater number of well-assessed models at the local scale. Models using opportunist data with a target-group approach to generate pseudo-absences provides a sufficient quality of information on species

460 distribution (Phillips et al., 2009; Barber et al., 2022) and can be correctly used in SCP (Sofaer et al., 2019;  
461 Baker et al., 2021). The lack of data at the regional/local scale, whatever the database, does not allow us to  
462 explain a better fit of models using GBIF data. Evaluating the models with a large proportion of opportunist  
463 data could however bias the evaluation, but only independent data were used, which provides robust  
464 validation of SDM (Matutini et al., 2021). Moreover, in contrast to Hermoso et al. (2015a), we found that  
465 mixing presence-only data with absence data increased the number of misjudged models. In addition to  
466 the use of the ROC curve (AUC) as a presence-only model evaluation metric by Hermoso et al. (2015a), the  
467 different results can be explained by different sampling biases between the two data types (Baker et al.,  
468 2022; Barber et al., 2022). Finally, the GBIF data seem to be more adapted to model the distribution of a  
469 large number of species.

470 The notion of “garbage in – garbage out” emphasises the critical importance of the quality of data  
471 (Sanders and Saxe, 2017), nevertheless, the examination of data suitability for conservation planning  
472 remains rare. In addition to the above issues our study revealed the importance of attention that should  
473 be paid to the representativeness of the species communities used in the models compared with the actual  
474 species communities observed in the study sites. This is particularly important in the light of the finding  
475 that there are marked differences between conservation priorities when different database sources are  
476 employed. Indeed, the number and composition of species in the community used can influence  
477 conservation priorities. Elsewhere it has been shown the difference will decline as the number of species  
478 increases (Kujala et al., 2018). The methodology tested in our study is based on a data-driven approach  
479 that attempts to use all available biodiversity data. This approach is data intensive, but is necessary to  
480 ensure the best representation of the observed local biodiversity. We revealed that such an approach can  
481 nevertheless induce a significant bias in the species community that is ultimately studied. Indeed, the  
482 prevalence of data affects the composition of the modelled species as well as the accuracy of the models  
483 and the evaluation of the species response (Fukuda and De Baets, 2016). Particular attention should thus  
484 be paid to the representativeness of the species communities used in the models in relation to the actual  
485 species communities observed in the territory under study- site.

486 Our study presents a workflow (Figure 2) for identifying biodiversity stakes using a data-driven  
487 approach from open-access database sources. Land-use planning/SEA stakeholders can use this workflow  
488 as a step towards the rationalization of data in order to reduce the biases mentioned above. The  
489 confrontation of the limits of such a workflow with the needs of SEA stakeholders could illustrate how to  
490 precisely target new sources of database that should be collected according to the suitability of current  
491 databases for priority groups. This workflow could be compared with the data context of another country  
492 to compare our findings. Hermoso et al. (2015b) revealed that evaluation models using a new collection of  
493 field data does not necessarily reduce the problems of model uncertainty. However, other databases can  
494 be examined by SEA stakeholders as well as other monitoring schemes (e.g. “PopAmphibien” for Reptilian  
495 and Amphibian populations in France <http://lashf.org/popamphibien-2/>) or negotiate the use of databases  
496 that are not yet shared. An important issue is thus the integration of~~To overcome this data sharing~~  
497 problem, the structuring of networks of different contributors of data and users of the databases and  
498 ambitious regional policies is necessary. As evidenced by our three study sites, the quantity of local data  
499 available is correlated with the number of years the SINP has been implemented. An important issue is  
500 thus the integration of SEA stakeholders in the workflow we propose, and their appropriation of the  
501 procedure. This could be done by a form of participatory modelling (Lagabrielle et al., 2010; Lees et al.,  
502 2021), where stakeholders are consulted for issues and choices such as the species to be examined. In such  
503 participatory modelling it is important to avoid arbitrary choices that are neither reproducible nor  
504 representative of local diversity, but rather the result of administrative or political interest. Finally, it is  
505 currently recommended to use these tools to elaborate a more holistic approach to SCP (Cadotte and  
506 Tucker, 2018).

## 508 **5 – Perspectives: operational implementation by SEA stakeholders**

509 Spatially exhaustive and ecologically representative priority conservation areas are crucial for the  
510 elaboration of SEAs that aim to limit artificialization impacts as early as possible in the planning process.  
511 Empirical observations are major sources of information on biodiversity that are still rarely used by SEAs.

512 The collection of open-access databases for SEA territories provides important but incomplete knowledge  
513 on species occurrence. Furthermore, their use is particularly interesting to help strategically direct  
514 inventory campaigns (especially for under sampled taxa and areas) that go beyond the emphasis on rare,  
515 threatened and emblematic species. These additional data would clearly improve the assessment of the  
516 SDMs suitability in ~~local territories; administrative entities as our study sites.~~ What is also interesting here  
517 for SEA stakeholders is that the process of filtering species and attributing them a score allows ~~us to~~  
518 identify for the identification of different groups of species in terms of their needs for additional data in  
519 order to undertake SDM.

520 ~~The influence of database sources on the identification of priority conservation areas reveals the~~  
521 ~~importance of examining their suitability. Thus, it is necessary to be prudent in the interpretation of~~  
522 ~~biodiversity maps. The integration of local experts may help limit any misjudgements in the workflow~~  
523 ~~procedure.~~

524 The influence of database sources on the identification of priority conservation areas reveals the  
525 importance of examining their suitability. In our study this is true for three highly contrasting study areas  
526 that differ markedly in terms of the cover of protected areas and artificialisation. The problem of data  
527 sources is thus typical of many areas. Thus, it is necessary to be prudent in the interpretation of biodiversity  
528 maps. The integration of local experts may help limit any misjudgements in the workflow procedure.  
529 Indeed, the integration of "expert" knowledge and local studies is valuable information, which is important  
530 to share, and which it is important to consider in order to complete our proposal. In future studies, species  
531 conservation issues for spatial prioritization could be considered by focusing on (for example) the issues  
532 associated with threatened and/or invasive species. The multiple dimensions of biodiversity could be  
533 analyzed within a context of limited data access and the complementarity of different facets (functional  
534 and phylogenetic) in addition to a classical species-based approach (Brumm et al., 2021; Cadotte and  
535 Tucker, 2018).

536 A data-driven approach that considers as many species as possible requires a large amount of data,  
537 biases the species communities considered and does not highlight species of particular interest as their  
538 threats and regulatory protections. It is therefore necessary to rationalize this approach, by integrating the  
539 needs and issues of local SEA stakeholders.

540



541

## Acknowledgements

542 ~~We thank Nicolas Dubos and Boris Leroy for our exchanges about SDM.~~ This work would not have been  
543 possible without the help of many voluntary field workers who continually provide open-access data. We  
544 are grateful to Benoit Fontaine for his work as administrator and for providing access to the Vigie Nature  
545 data, to Annegret Nicolai for collection of all open-access data on “Communauté de commune de  
546 Brocéliande (i.e. the study site “T3”), to Paul Fromage for providing access for Fauna data, to Mathieu  
547 Largarde for OEB data access, to Solène Robert for the exchanges about SINP database and Joel Kamdoun  
548 Ngueuko for access to SINP Occitanie data. We also thank Josselin Giffard and Samuel Alleaume for access  
549 to DHI NDVI, to Karine Princé for discussion on agricultural intensification variables and to Jean-Pierre  
550 Moussus for access to Papilionidae trait databases and Jeremy Froidevaux for the preview access to the  
551 EuroBaTrait <https://doi.org/10.1038/s41597-023-02157-4>. We thank Nicolas Dubos and Boris Leroy for  
552 discussion on the use of SDM and Eric Durand for ~~helpful~~ more general discussion of this study. Finally, we  
553 also acknowledge the reviewers which improved this version.

554

## Data, scripts, code, and supplementary information availability

555 Observation data from SINP and Vigie nature databases are not available for confidentiality reasons,  
556 but their link and the request process are detailed in Appendix A1. The data of GBIF are available online:  
557 <https://doi.org/10.15468/dl.ry6uw7>.

558 The data, R scripts, outputs of steps 1.2 to 4 of figure 2 are available online:  
559 <https://doi.org/10.5281/zenodo.7883973>.

560

## Conflict of interest disclosure

561 The authors declare that they comply with the PCI rule of having no financial conflicts of interest in  
562 relation to the content of the article.

563

## Funding

564 This study was funded by “Naturalia Environnement” and “Association Nationale de la Recherche et de  
565 la Technologie” (grant number: 2020/0584).

566

## References

- 567 Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B., Anderson, R.P., 2015. spThin: an R package  
568 for spatial thinning of species occurrence records for use in ecological niche models. *Ecography* 38, 541–  
569 545. <https://doi.org/10.1111/ecog.01132>
- 570 Almenar, J.B., Bolowich, A., Elliot, T., Geneletti, D., Sonnemann, G., Rugani, B., 2019. Assessing habitat loss,  
571 fragmentation and ecological connectivity in Luxembourg to support spatial planning. *Landscape and*  
572 *Urban Planning* 189, 335–351. <https://doi.org/10.1016/j.landurbplan.2019.05.004>
- 573 Anderson, R.P., 2003. Real vs. artefactual absences in species distributions: tests for *Oryzomys albigularis*  
574 (Rodentia: Muridae) in Venezuela. *Journal of Biogeography* 30, 591–605.  
575 <https://doi.org/10.1046/j.1365-2699.2003.00867.x>
- 576 Baker, D.J., Maclean, I.M.D., Goodall, M., Gaston, K.J., 2022. Correlations between spatial sampling biases  
577 and environmental niches affect species distribution models. *Global Ecology and Biogeography* 31,  
578 1038–1050. <https://doi.org/10.1111/geb.13491>
- 579 Baker, D.J., Maclean, I.M.D., Goodall, M., Gaston, K.J., 2021. Species distribution modelling is needed to  
580 support ecological impact assessments. *Journal of Applied Ecology* 21–26.  
581 <https://doi.org/10.1111/1365-2664.13782>
- 582 Baker, M., Carter, J., Short, M., Jay, S., 2005. *Strategic Environmental Assessment and Land Use Planning:*  
583 *An International Evaluation.* Routledge.

584 Barber, R.A., Ball, S.G., Morris, R.K.A., Gilbert, F., 2022. Target-group backgrounds prove effective at  
585 correcting sampling bias in Maxent models. *Diversity and Distributions* 28, 128–141.  
586 <https://doi.org/10.1111/ddi.13442>

587 Beck, J., Ballesteros-Mejia, L., Nagel, P., Kitching, I.J., 2013. Online solutions and the ‘Wallacean shortfall’:  
588 what does GBIF contribute to our knowledge of species’ ranges? *Diversity and Distributions* 19, 1043–  
589 1050. <https://doi.org/10.1111/ddi.12083>

590 [Beck, J., Böller, M., Erhardt, A., Schwanghart, W., 2014. Spatial bias in the GBIF database and its effect on](https://doi.org/10.1016/j.ecoinf.2013.11.002)  
591 [modeling species’ geographic distributions. \*Ecological Informatics\* 19, 10–15.](https://doi.org/10.1016/j.ecoinf.2013.11.002)  
592 <https://doi.org/10.1016/j.ecoinf.2013.11.002>

593 Bezombes, L., Kerbiriou, C., Spiegelberger, T., 2019. Do biodiversity offsets achieve No Net Loss? An  
594 evaluation of offsets in a French department. *Biological Conservation* 231, 24–29.  
595 <https://doi.org/10.1016/j.biocon.2019.01.004>

596 Bigard, C., Pioch, S., Thompson, J.D., 2017. The inclusion of biodiversity in environmental impact  
597 assessment: Policy-related progress limited by gaps and semantic confusion. *Journal of Environmental*  
598 *Management* 200, 35–45. <https://doi.org/10.1016/j.jenvman.2017.05.057>

599 Bigard, C., Thiriet, P., Pioch, S., Thompson, J.D., 2020. Strategic landscape-scale planning to improve  
600 mitigation hierarchy implementation: An empirical case study in Mediterranean France. *Land Use Policy*  
601 90, 104286. <https://doi.org/10.1016/j.landusepol.2019.104286>

602 Boileau, J., Calvet, C., Pioch, S., Moulherat, S., 2022. Ecological equivalence assessment: The potential of  
603 genetic tools, remote sensing and metapopulation models to better apply the mitigation hierarchy.  
604 *Journal of Environmental Management* 305, 114415. <https://doi.org/10.1016/j.jenvman.2021.114415>

605 [Botella, C., Joly, A., Bonnet, P., Monestiez, P., Munoz, F., 2018. A Deep Learning Approach to Species](https://doi.org/10.1007/978-3-319-76445-0_10)  
606 [Distribution Modelling, in: Joly, A., Vrochidis, S., Karatzas, K., Karppinen, A., Bonnet, P. \(Eds.\),](https://doi.org/10.1007/978-3-319-76445-0_10)  
607 [Multimedia Tools and Applications for Environmental & Biodiversity Informatics. Springer International](https://doi.org/10.1007/978-3-319-76445-0_10)  
608 [Publishing, Cham, pp. 169–199. https://doi.org/10.1007/978-3-319-76445-0\\_10](https://doi.org/10.1007/978-3-319-76445-0_10)

609 Boyce, M.S., Vernier, P.R., Nielsen, S.E., Schmiegelow, F.K.A., 2002. Evaluating resource selection functions.  
610 *Ecological Modelling* 157, 281–300. [https://doi.org/10.1016/S0304-3800\(02\)00200-4](https://doi.org/10.1016/S0304-3800(02)00200-4)

611 Boyd, R.J., Stewart, G., Pescott, O., 2023. Descriptive inference using large, unrepresentative  
612 nonprobability samples: An introduction for ecologists.

613 [Brumm, K.J., Hanks, R.D., Baldwin, R.F., Peoples, B.K., 2021. Accounting for multiple dimensions of](https://doi.org/10.1016/j.ecolind.2020.107320)  
614 [biodiversity to assess surrogate performance in a freshwater conservation prioritization. \*Ecological\*](https://doi.org/10.1016/j.ecolind.2020.107320)  
615 [Indicators 122, 107320. https://doi.org/10.1016/j.ecolind.2020.107320](https://doi.org/10.1016/j.ecolind.2020.107320)

616 Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R.O., Wang, Z., Zimmermann, N.E., 2020. Model  
617 complexity affects species distribution projections under climate change. *Journal of Biogeography* 47,  
618 130–142. <https://doi.org/10.1111/jbi.13734>

619 Bull, J.W., Gordon, A., Watson, J.E.M., Maron, M., 2016. Seeking convergence on the key concepts in ‘no  
620 net loss’ policy. *Journal of Applied Ecology* 53, 1686–1693. <https://doi.org/10.1111/1365-2664.12726>

621 Cadotte, M.W., Tucker, C.M., 2018. Difficult decisions: Strategies for conservation prioritization when  
622 taxonomic, phylogenetic and functional diversity are not spatially congruent. *Biological Conservation*  
623 225, 128–133. <https://doi.org/10.1016/j.biocon.2018.06.014>

624 Canbek, G., 2022. Gaining insights in datasets in the shade of “garbage in, garbage out” rationale: Feature  
625 space distribution fitting. *WIREs Data Mining and Knowledge Discovery* 12, e1456.  
626 <https://doi.org/10.1002/widm.1456>

627 Carmona, C.P., Tamme, R., Pärtel, M., de Bello, F., Brosse, S., Capdevila, P., González-M., R., González-  
628 Suárez, M., Salguero-Gómez, R., Vásquez-Valderrama, M., Toussaint, A., 2021. Erosion of global  
629 functional diversity across the tree of life. *Science Advances* 7, eabf2675.  
630 <https://doi.org/10.1126/sciadv.abf2675>

631 CESBIO, 2022. Dynamic Habitat Indices of NDVI - Available data [WWW Document]. URL  
632 <https://files.coinduf.eu/AFSfaJ2Kynt8iYWw/> (accessed 4.3.23).

633 CESBIO, 2021. Variables for Biodiversity – Theia [WWW Document]. URL [https://www.theia-](https://www.theia-land.fr/en/ceslist/variables-for-biodiversity-sec/)  
634 [land.fr/en/ceslist/variables-for-biodiversity-sec/](https://www.theia-land.fr/en/ceslist/variables-for-biodiversity-sec/) (accessed 4.3.23).

635 Christensen, R., 2022. ordinal: Regression Models for Ordinal Data version 2022.11-16 from CRAN.

636 Church, R.L., Stoms, D.M., Davis, F.W., 1996. Reserve selection as a maximal covering location problem.  
637 *Biological Conservation* 76, 105–112. [https://doi.org/10.1016/0006-3207\(95\)00102-6](https://doi.org/10.1016/0006-3207(95)00102-6)

638 Clare, J.D.J., Townsend, P.A., Anhalt-Depies, C., Locke, C., Stenglein, J.L., Frett, S., Martin, K.J., Singh, A., Van  
639 Deelen, T.R., Zuckerberg, B., 2019. Making inference with messy (citizen science) data: when are data  
640 accurate enough and how can they be improved? *Ecological Applications* 29, e01849.  
641 <https://doi.org/10.1002/eap.1849>

642 Costello, M.J., Wieczorek, J., 2014. Best practice for biodiversity data management and publication.  
643 *Biological Conservation* 173, 68–73. <https://doi.org/10.1016/j.biocon.2013.10.018>

644 Domisch, S., Friedrichs, M., Hein, T., Borgwardt, F., Wetzig, A., Jähnig, S.C., Langhans, S.D., 2019. Spatially  
645 explicit species distribution models: A missed opportunity in conservation planning? *Diversity and*  
646 *Distributions* 25, 758–769. <https://doi.org/10.1111/ddi.12891>

647 Dubos, N., Préau, C., Lenormand, M., Papuga, G., Monsarrat, S., Denelle, P., Louarn, M.L., Heremans, S.,  
648 May, R., Roche, P., Luque, S., 2022. Assessing the effect of sample bias correction in species distribution  
649 models. *Ecological Indicators* 145, 109487. <https://doi.org/10.1016/j.ecolind.2022.109487>

650 Dupont, P., Demerges, D., Drouet, E., Luquet, G.Chr., 2013. Révision systématique, taxinomique et  
651 nomenclaturale des Rhopalocera et des Zygaenidae de France métropolitaine. Conséquences sur  
652 l'acquisition et la gestion des données d'inventaire (No. 19). MNHN-SPN.

653 Elvidge, C.D., Zhizhin, M., Ghosh, T., Hsu, F.-C., Taneja, J., 2021. Annual Time Series of Global VIIRS  
654 Nighttime Lights Derived from Monthly Averages: 2012 to 2019. *Remote Sensing* 13, 922.  
655 <https://doi.org/10.3390/rs13050922>

656 Essens, T., van Langevelde, F., Vos, R.A., Van Swaay, C.A.M., WallisDeVries, M.F., 2017. Ecological  
657 determinants of butterfly vulnerability across the European continent. *Journal of Insect Conservation*  
658 21, 439–450. <https://doi.org/10.1007/s10841-017-9972-4>

659 Fahrig, L., 1997. Relative effects of habitat loss and fragmentation on population extinction. *The Journal of*  
660 *wildlife management* 61, 603–610. <https://doi.org/10.2307/3802168>

661 Ferrier, S., Pressey, R.L., Barrett, T.W., 2000. A new predictor of the irreplaceability of areas for achieving  
662 a conservation goal, its application to real-world planning, and a research agenda for further  
663 refinement. *Biological Conservation* 93, 303–325. [https://doi.org/10.1016/S0006-3207\(99\)00149-4](https://doi.org/10.1016/S0006-3207(99)00149-4)

664 [Fithian, W., Elith, J., Hastie, T., Keith, D.A., 2015. Bias correction in species distribution models: pooling  
665 survey and collection data for multiple species. \*Methods in Ecology and Evolution\* 6, 424–438.  
666 <https://doi.org/10.1111/2041-210X.12242>](https://doi.org/10.1111/2041-210X.12242)

667 Fletcher, R.J., Hefley, T.J., Robertson, E.P., Zuckerberg, B., McCleery, R.A., Dorazio, R.M., 2019. A practical  
668 guide for combining data to model species distributions. *Ecology* 100, e02710.  
669 <https://doi.org/10.1002/ecy.2710>

670 Fukuda, S., De Baets, B., 2016. Data prevalence matters when assessing species' responses using data-  
671 driven species distribution models. *Ecological Informatics* 32, 69–78.  
672 <https://doi.org/10.1016/j.ecoinf.2016.01.005>

673 Fundingsland Tetlow, M., Hanusch, M., 2012. Strategic environmental assessment: the state of the art.  
674 *Impact Assessment and Project Appraisal* 30, 15–24. <https://doi.org/10.1080/14615517.2012.666400>

675 Gargominy, O., Terceirie, S., Régnier, C., Ramage, T., Dupont, T., Daszkiewicz, P., Poncet, L., 2021. TAXREF  
676 v14, référentiel taxonomique pour la France : méthodologie, mise en oeuvre et diffusion, Rapport  
677 Patrinat. UMS PatriNat (OFB-CNRS-MNHN), Paris, France.

678 Gontier, M., Balfors, B., Mörtberg, U., 2006. Biodiversity in environmental assessment - current practice  
679 and tools for prediction. *Environmental Impact Assessment Review* 26, 268–286.  
680 <https://doi.org/10.1016/j.eiar.2005.09.001>

681 Guillera-Arroita, G., Lahoz-Monfort, J.J., Elith, J., Gordon, A., Kujala, H., Lentini, P.E., McCarthy, M.A.,  
682 Tingley, R., Wintle, B.A., 2015. Is my species distribution model fit for purpose? Matching data and  
683 models to applications. *Global Ecology and Biogeography* 24, 276–292.  
684 <https://doi.org/10.1111/geb.12268>

685 Guillet, F., Le Floch, C., Julliard, R., 2019. Séquence Eviter-Réduire-Compenser : quelle biodiversité est visée  
686 par les mesures d'évitement? *Sciences Eaux et Territoires* 1–8. <https://doi.org/10.14758/SET-REVUE.2019.HS.07>

687

688 Guisan, A., Thuiller, W., Zimmermann, N.E., 2017. *Habitat Suitability and Distribution Models: with*  
689 *Applications in R*. Cambridge University Press.

690 Guisan, A., Tingley, R., Baumgartner, J.B., Naujokaitis-Lewis, I., Sutcliffe, P.R., Tulloch, A.I.T., Regan, T.J.,  
691 Brotons, L., McDonald-Madden, E., Mantyka-Pringle, C., Martin, T.G., Rhodes, J.R., Maggini, R.,

692 Setterfield, S.A., Elith, J., Schwartz, M.W., Wintle, B.A., Broennimann, O., Austin, M., Ferrier, S., Kearney,  
693 M.R., Possingham, H.P., Buckley, Y.M., 2013. Predicting species distributions for conservation decisions.  
694 Ecology Letters 16, 1424–1435. <https://doi.org/10.1111/ele.12189>  
695 Hanson, J.O., Schuster, R., Morrell, N., Strimas-Mackey, M., Edwards, B.P.M., Watts, M.E., Arcese, P.,  
696 Bennett, J., Possingham, H.P., 2021. prioritizr: Systematic Conservation Prioritization in R.  
697 Hermoso, V., Kennard, M.J., Linke, S., 2015a. Assessing the risks and opportunities of presence-only data  
698 for conservation planning. Journal of Biogeography 42, 218–228. <https://doi.org/10.1111/jbi.12393>  
699 Hermoso, V., Kennard, M.J., Linke, S., 2015b. Evaluating the costs and benefits of systematic data  
700 acquisition for conservation assessments. Ecography 38, 283–292. <https://doi.org/10.1111/ecog.00792>  
701 Horváth, Z., Ptacnik, R., Vad, C.F., Chase, J.M., 2019. Habitat loss over six decades accelerates regional and  
702 local biodiversity loss via changing landscape connectance. Ecology Letters 22, 1019–1027.  
703 <https://doi.org/10.1111/ele.13260>  
704 IGN, 2022. Géoservices - Téléchargement de données [WWW Document]. URL  
705 <https://geoservices.ign.fr/telechargement> (accessed 4.3.23).  
706 Inglada, J., Vincent, A., Thierion, V., 2019. Theia OSO Land Cover Map 2019 [Data set]. Zenodo.  
707 <https://doi.org/10.5281/zenodo.6538321>  
708 Iorio, É., Dusoulier, F., Soldati, F., Noël, F., Guilloton, J.-A., Doucet, G., Ponel, P., Dupont, P., Krieg-Jacquier,  
709 R., Chemin, S., Tillier, P., Touroult, J., 2022. Les Arthropodes terrestres dans les études d’impact : limites  
710 actuelles et propositions pour une meilleure prise en compte des enjeux de conservation. *Naturae* 43–  
711 99. <https://doi.org/10.5852/naturae2022a4>  
712 IPBES, 2019. Global assessment report on biodiversity and ecosystem services of the Intergovernmental  
713 Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES secretariat.  
714 <https://doi.org/10.5281/zenodo.6417333>  
715 Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K., Mooers, A.O., 2012. The global diversity of birds in space  
716 and time. *Nature* 491, 444–448. <https://doi.org/10.1038/nature11631>  
717 Jomier, R., Solène, R., Dupont, P., 2018. National Scientific Validation for Data in the Information System  
718 on Nature and Landscapes (Système d’Information sur la Nature et les Paysages - SINP). Biodiversity  
719 Information Science and Standards e25881. <https://doi.org/10.3897/biss.2.25881>  
720 Julliard, R., Jiguet, F., 2002. Un suivi intégré des populations d’oiseaux communs en France, in: *Alauda*.  
721 Presented at the Colloque Francophone Ornithologique, Limoges, France, pp. 137–147.  
722 Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann, N.E., Linder, H.P.,  
723 Kessler, M., 2021. Climatologies at high resolution for the earth’s land surface areas. *EnviDat* 170122.  
724 <https://doi.org/10.16904/envidat.228.v2.1>  
725 Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann, N.E., Linder, H.P.,  
726 Kessler, M., 2017. Climatologies at high resolution for the earth’s land surface areas. *Scientific Data* 4,  
727 170122. <https://doi.org/10.1038/sdata.2017.122>  
728 Kim, V., Ralphs, T., Guzelsoy, M., Mahajan, A., Reinhard, H., Hornik, K., Szymanski, C., Theussl, S., 2023.  
729 Ipsymphony: Symphony integer linear programming solver in R.  
730 Kujala, H., Moilanen, A., Gordon, A., 2018. Spatial characteristics of species distributions as drivers in  
731 conservation prioritization. *Methods in Ecology and Evolution* 9, 1121–1132.  
732 <https://doi.org/10.1111/2041-210X.12939>  
733 Lagabrielle, E., Botta, A., Daré, W., David, D., Aubert, S., Fabricius, C., 2010. Modelling with stakeholders to  
734 integrate biodiversity into land-use planning - Lessons learned in Réunion Island (Western Indian  
735 Ocean). *Environmental Modelling & Software* 25, 1413–1427.  
736 <https://doi.org/10.1016/j.envsoft.2010.01.011>  
737 Lees, C.M., Rutschmann, A., Santure, A.W., Beggs, J.R., 2021. Science-based, stakeholder-inclusive and  
738 participatory conservation planning helps reverse the decline of threatened species. *Biological*  
739 *Conservation* 260, 109194. <https://doi.org/10.1016/j.biocon.2021.109194>  
740 Leroy, B., Delsol, R., Hugueny, B., Meynard, C.N., Barhoumi, C., Barbet-Massin, M., Bellard, C., 2018.  
741 Without quality presence–absence data, discrimination metrics such as TSS can be misleading measures  
742 of model performance. *Journal of Biogeography* 45, 1994–2002. <https://doi.org/10.1111/jbi.13402>  
743 Lino, A., Fonseca, C., Rojas, D., Fischer, E., Pereira, M.J.R., 2019. A meta-analysis of the effects of habitat  
744 loss and fragmentation on genetic diversity in mammals. *Mammalian Biology* 94, 69–76.  
745 <https://doi.org/10.1016/j.mambio.2018.09.006>

746 Mac Nally, R., Duncan, R.P., Thomson, J.R., Yen, J.D.L., 2018. Model selection using information criteria, but  
747 is the “best” model any good? *Journal of Applied Ecology* 55, 1441–1444.  
748 <https://doi.org/10.1111/1365-2664.13060>

749 Margules, C.R., Pressey, R.L., 2000. Systematic conservation planning. *Nature* 405, 243–253.  
750 <https://doi.org/10.1038/35012251>

751 Mariton, L., Kerbiriou, C., Bas, Y., Zanda, B., Le Viol, I., 2022. Even low light pollution levels affect the spatial  
752 distribution and timing of activity of a “light tolerant” bat species. *Environmental Pollution* 305, 119267.  
753 <https://doi.org/10.1016/j.envpol.2022.119267>

754 Maron, M., Ives, C.D., Kujala, H., Bull, J.W., Maseyk, F.J.F., Bekessy, S., Gordon, A., Watson, J.E.M., Lentini,  
755 P.E., Gibbons, P., Possingham, H.P., Hobbs, R.J., Keith, D.A., Wintle, B.A., Evans, M.C., 2016. Taming a  
756 Wicked Problem: Resolving Controversies in Biodiversity Offsetting. *BioScience* 66, 489–498.  
757 <https://doi.org/10.1093/biosci/biw038>

758 Matsuki, K., Kuperman, V., Van Dyke, J.A., 2016. The Random Forests statistical technique: An examination  
759 of its value for the study of reading. *Scientific Studies of Reading* 20, 20–33.  
760 <https://doi.org/10.1080/10888438.2015.1107073>

761 Matutini, F., Baudry, J., Pain, G., Sineau, M., Pithon, J., 2021. How citizen science could improve species  
762 distribution models and their independent assessment. *Ecology and Evolution* 11, 3028–3039.  
763 <https://doi.org/10.1101/2020.06.02.129536>

764 Maxwell, S.L., Fuller, R.A., Brooks, T.M., Watson, J.E.M., 2016. Biodiversity: The ravages of guns, nets and  
765 bulldozers. *Nature News* 536, 143–145. <https://doi.org/10.1038/536143a>

766 MEDDE, GIS Sol, 2014. . Enveloppes des milieux potentiellement humides de la France métropolitaine.  
767 Notice d’accompagnement. Programme de modélisation des milieux potentiellement humides de  
768 France. Ministère d’Ecologie, du Développement Durable et de l’Energie, Groupement d’Intérêt  
769 Scientifique Sol.

770 Meyer, C., Kreft, H., Guralnick, R., Jetz, W., 2015. Global priorities for an effective information basis of  
771 biodiversity distributions. *Nature Communications* 6, 1–8. <https://doi.org/10.1038/ncomms9221>

772 Meyer, C.B., 2007. Does Scale Matter in Predicting Species Distributions? Case Study with the Marbled  
773 Murrelet. *Ecological Applications* 17, 1474–1483. <https://doi.org/10.1890/06-1410.1>

774 Middleton-Welling, J., Dapporto, L., García-Barros, E., Wiemers, M., Nowicki, P., Plazio, E., Bonelli, S.,  
775 Zaccagno, M., Šašić, M., Liparova, J., Schweiger, O., Harpke, A., Musche, M., Settele, J., Schmucki, R.,  
776 Shreeve, T., 2020. A new comprehensive trait database of European and Maghreb butterflies,  
777 Papilionoidea. *Scientific Data* 7, 351. <https://doi.org/10.1038/s41597-020-00697-7>

778 Morelli, F., Benedetti, Y., Callaghan, C.T., 2020. Ecological specialization and population trends in European  
779 breeding birds. *Global Ecology and Conservation* 22, e00996.  
780 <https://doi.org/10.1016/j.gecco.2020.e00996>

781 Moussus, J.-P., Lorin, T., Cooper, A., 2019. Guide pratique des papillons de France, Delachaux et Niestlé.  
782 ed. France.

783 Olden, J.D., Rooney, T.P., 2006. On defining and quantifying biotic homogenization. *Global Ecology and*  
784 *Biogeography* 15, 113–120. <https://doi.org/10.1111/j.1466-822X.2006.00214.x>

785 Phalan, B., Hayes, G., Brooks, S., Marsh, D., Howard, P., Costelloe, B., Vira, B., Kowalska, A., Whitaker, S.,  
786 2018. Avoiding impacts on biodiversity through strengthening the first stage of the mitigation hierarchy.  
787 *Oryx* 52, 316–324. <https://doi.org/10.1017/S0030605316001034>

788 Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J., Ferrier, S., 2009. Sample selection  
789 bias and presence-only distribution models: implications for background and pseudo-absence data.  
790 *Ecological Applications* 19, 181–197. <https://doi.org/10.1890/07-2153.1>

791 Ponder, W.F., Carter, G.A., Flemons, P., Chapman, R.R., 2001. Evaluation of Museum Collection Data for  
792 Use in Biodiversity Assessment. *Conservation Biology* 15, 648–657. <https://doi.org/10.1046/j.1523-1739.2001.015003648.x>

793

794 Pope, J., Bond, A., Morrison-Saunders, A., Retief, F., 2013. Advancing the theory and practice of impact  
795 assessment: Setting the research agenda. *Environmental Impact Assessment Review* 41, 1–9.  
796 <https://doi.org/10.1016/j.eiar.2013.01.008>

797 Pressey, R.L., Bottrill, M.C., 2008. Opportunism, Threats, and the Evolution of Systematic Conservation  
798 Planning. *Conservation Biology* 22, 1340–1345. <https://doi.org/10.1111/j.1523-1739.2008.01032.x>



799 Quétier, F., Regnery, B., Levrel, H., 2014. No net loss of biodiversity or paper offsets? A critical review of  
800 the French no net loss policy. *Environmental Science & Policy* 38, 120–131.  
801 <https://doi.org/10.1016/j.envsci.2013.11.009>

802 Rondinini, C., Wilson, K.A., Boitani, L., Grantham, H., Possingham, H.P., 2006. Tradeoffs of different types  
803 of species occurrence data for use in systematic conservation planning. *Ecology Letters* 9, 1136–1145.  
804 <https://doi.org/10.1111/j.1461-0248.2006.00970.x>

805 Sanders, H., Saxe, J., 2017. Garbage in, garbage out: how purportedly great ML models can be screwed up  
806 by bad data. Presented at the Blackhat, Las Vegas, USA, p. 6.

807 Schoener, T.W., 1968. The Anolis Lizards of Bimini: Resource Partitioning in a Complex Fauna. *Ecology* 49,  
808 704–726. <https://doi.org/10.2307/1935534>

809 Sheard, C., Neate-Clegg, M.H.C., Alioravainen, N., Jones, S.E.I., Vincent, C., MacGregor, H.E.A., Bregman,  
810 T.P., Claramunt, S., Tobias, J.A., 2020. Ecological drivers of global gradients in avian dispersal inferred  
811 from wing morphology. *Nature Communications* 11, 2463. [https://doi.org/10.1038/s41467-020-16313-](https://doi.org/10.1038/s41467-020-16313-6)  
812 6

813 Shirey, V., Belitz, M.W., Barve, V., Guralnick, R., 2021. A complete inventory of North American butterfly  
814 occurrence data: narrowing data gaps, but increasing bias. *Ecography* 44, 537–547.  
815 <https://doi.org/10.1111/ecog.05396>

816 Sofaer, H.R., Jarnevich, C.S., Pearse, I.S., Smyth, R.L., Auer, S., Cook, G.L., Edwards, T.C., Jr, Guala, G.F.,  
817 Howard, T.G., Morisette, J.T., Hamilton, H., 2019. Development and Delivery of Species Distribution  
818 Models to Inform Decision-Making. *BioScience* 69, 544–557. <https://doi.org/10.1093/biosci/biz045>

819 Sousa-Baena, M.S., Garcia, L.C., Peterson, A.T., 2014. Completeness of digital accessible knowledge of the  
820 plants of Brazil and priorities for survey and inventory. *Diversity and Distributions* 20, 369–381.  
821 <https://doi.org/10.1111/ddi.12136>

822 Srisa-An, C., 2021. Guideline of Collinearity - Avoidable Regression Models on Time-series Analysis, in: 2021  
823 2nd International Conference on Big Data Analytics and Practices (IBDAP). Presented at the 2021 2nd  
824 International Conference on Big Data Analytics and Practices (IBDAP), Bangkok, Thailand, pp. 28–32.  
825 <https://doi.org/10.1109/IBDAP52511.2021.9552165>

826 Stekhoven, D.J., Bühlmann, P., 2012. MissForest--non-parametric missing value imputation for mixed-type  
827 data. *Bioinformatics* 28, 112–118. <https://doi.org/10.1093/bioinformatics/btr597>

828 Støa, B., Halvorsen, R., Stokland, J.N., Gusarov, V.I., 2019. How much is enough? Influence of number of  
829 presence observations on the performance of species distribution models. *Sommerfeltia* 39, 1–28.  
830 <https://doi.org/10.2478/som-2019-0001>

831 Storchová, L., Hořák, D., 2018. Life-history characteristics of European birds. *Global Ecology and*  
832 *Biogeography* 27, 400–406. <https://doi.org/10.1111/geb.12709>

833 Tarabon, S., Bergès, L., Dutoit, T., Isselin-Nondedeu, F., 2019. Environmental impact assessment of  
834 development projects improved by merging species distribution and habitat connectivity modelling.  
835 *Journal of Environmental Management* 241, 439–449. <https://doi.org/10.1016/j.jenvman.2019.02.031>

836 Telenius, A., 2011. Biodiversity information goes public: GBIF at your service. *Nordic Journal of Botany* 29,  
837 378–381. <https://doi.org/10.1111/j.1756-1051.2011.01167.x>

838 Tittensor, D.P., Walpole, M., Hill, S.L.L., Boyce, D.G., Britten, G.L., Burgess, N.D., Butchart, S.H.M., Leadley,  
839 P.W., Regan, E.C., Alkemade, R., Baumung, R., Bellard, C., Bouwman, L., Bowles-Newark, N.J., Chenery,  
840 A.M., Cheung, W.W.L., Christensen, V., Cooper, H.D., Crowther, A.R., Dixon, M.J.R., Galli, A., Gaveau, V.,  
841 Gregory, R.D., Gutierrez, N.L., Hirsch, T.L., Höft, R., Januchowski-Hartley, S.R., Karmann, M., Krug, C.B.,  
842 Leverington, F.J., Loh, J., Lojenga, R.K., Malsch, K., Marques, A., Morgan, D.H.W., Mumby, P.J., Newbold,  
843 T., Noonan-Mooney, K., Pagad, S.N., Parks, B.C., Pereira, H.M., Robertson, T., Rondinini, C., Santini, L.,  
844 Scharlemann, J.P.W., Schindler, S., Sumaila, U.R., Teh, L.S.L., van Kolck, J., Visconti, P., Ye, Y., 2014. A  
845 mid-term analysis of progress toward international biodiversity targets. *Science* 346, 241–244.  
846 <https://doi.org/10.1126/science.1257484>

847 Toussaint, A., Brosse, S., Bueno, C.G., Pärtel, M., Tamme, R., Carmona, C.P., 2021. Extinction of threatened  
848 vertebrates will lead to idiosyncratic changes in functional diversity across the world. *Nature*  
849 *Communications* 12, 5162. <https://doi.org/10.1038/s41467-021-25293-0>

850 Troudet, J., Grandcolas, P., Blin, A., Vignes-Lebbe, R., Legendre, F., 2017. Taxonomic bias in biodiversity  
851 data and societal preferences. *Scientific Reports* 7, 9132. <https://doi.org/10.1038/s41598-017-09084-6>

852 Tulloch, A.I.T., Gordon, A., Runge, C.A., Rhodes, J.R., 2019. Integrating spatially realistic infrastructure  
853 impacts into conservation planning to inform strategic environmental assessment. *Conservation Letters*  
854 12, e12648. <https://doi.org/10.1111/conl.12648>

855 Tulloch, A.I.T., Sutcliffe, P., Naujokaitis-Lewis, I., Tingley, R., Brotons, L., Ferraz, K.M.P.M.B., Possingham,  
856 H., Guisan, A., Rhodes, J.R., 2016. Conservation planners tend to ignore improved accuracy of modelled  
857 species distributions to focus on multiple threats and ecological processes. *Biological Conservation* 199,  
858 157–171. <https://doi.org/10.1016/j.biocon.2016.04.023>

859 Valavi, R., Elith, J., Lahoz-Monfort, J.J., Guillera-Arroita, G., 2021a. Modelling species presence-only data  
860 with random forests. *Ecography* 44, 1731–1742. <https://doi.org/10.1111/ecog.05615>

861 Valavi, R., Guillera-Arroita, G., Lahoz-Monfort, J.J., Elith, J., 2021b. Predictive performance of presence-only  
862 species distribution models: a benchmark study with reproducible code. *Ecological Monographs*.  
863 <https://doi.org/10.1002/ecm.1486>

864 Velazco, S.J.E., Ribeiro, B.R., Laureto, L.M.O., De Marco Júnior, P., 2020. Overprediction of species  
865 distribution models in conservation planning: A still neglected issue with strong effects. *Biological*  
866 *Conservation* 252, 108822. <https://doi.org/10.1016/j.biocon.2020.108822>

867 Warren, D., Dinnage, R., 2022. ENMTools: Analysis of Niche Evolution using Niche and Distribution Models.

868 Warren, D.L., Glor, R.E., Turelli, M., 2008. Environmental Niche Equivalency Versus Conservatism:  
869 Quantitative Approaches to Niche Evolution. *Evolution* 62, 2868–2883. [https://doi.org/10.1111/j.1558-](https://doi.org/10.1111/j.1558-5646.2008.00482.x)  
870 [5646.2008.00482.x](https://doi.org/10.1111/j.1558-5646.2008.00482.x)

871 Whitehead, A.L., Kujala, H., Wintle, B.A., 2017. Dealing with Cumulative Biodiversity Impacts in Strategic  
872 Environmental Assessment: A New Frontier for Conservation Planning. *Conservation Letters* 10, 195–  
873 204. <https://doi.org/10.1111/conl.12260>

874 Wiemers, M., Chazot, N., Wheat, C.W., Schweiger, O., Wahlberg, N., 2020. A complete time-calibrated  
875 multi-gene phylogeny of the European butterflies. *ZooKeys* 938, 97–124.  
876 <https://doi.org/10.3897/zookeys.938.50878>

877 Wiemers, M., Chazot, N., Wheat, C.W., Schweiger, O., Wahlberg, N., 2019. A complete time-calibrated  
878 multi-gene phylogeny of the European butterflies - data [Data set]. Zenodo.  
879 <https://doi.org/10.5281/zenodo.3531555>

880 Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M.M., Jetz, W., 2014. EltonTraits 1.0:  
881 Species-level foraging attributes of the world's birds and mammals. *Ecology* 95, 2027–2027.  
882 <https://doi.org/10.1890/13-1917.1>

883 Witté, I., Touroult, J., 2017. Identification et cartographie des zones de méconnaissance naturaliste à  
884 l'échelle nationale (métropole) à partir des données partagées (No. 2017–6). Muséum National  
885 d'Histoire Naturel (MNHN) - Service du Patrimoine Naturel (SPN), France.

886 Zambrano, J., Garzon-Lopez, C.X., Yeager, L., Fortunel, C., Cordeiro, N.J., Beckman, N.G., 2019. The effects  
887 of habitat loss and fragmentation on plant functional traits and functional diversity: what do we know  
888 so far? *Oecologia* 191, 505–518. <https://doi.org/10.1007/s00442-019-04505-x>

889 Zuckerberg, B., Huettmann, F., Frair, J., 2011. Proper Data Management as a Scientific Foundation for  
890 Reliable Species Distribution Modeling, in: Drew, C.A., Wiersma, Y.F., Huettmann, F. (Eds.), *Predictive*  
891 *Species and Habitat Modeling in Landscape Ecology: Concepts and Applications*. Springer, New York,  
892 NY, pp. 45–70. [https://doi.org/10.1007/978-1-4419-7390-0\\_4](https://doi.org/10.1007/978-1-4419-7390-0_4)

893 Zurell, D., Franklin, J., König, C., Bouchet, P., Dormann, C., Elith, J., Fandos, G., Feng, X., Guillera-Arroita, G.,  
894 Guisan, A., Lahoz-Monfort, J.J., Leitão, P.J., Park, D., Peterson, A.T., Rapacciuolo, G., Schmatz, D.,  
895 Schröder, B., Serra-Diaz, J.M., Thuiller, W., Yates, K., Zimmermann, N., Merow, C., 2020. A standard  
896 protocol for reporting species distribution models. <https://doi.org/10.18452/22268>

897

898

900 **Appendix A - Additional material and methods**901 *A.1 - Collect and series of operation on species occurrences*

## 902 Databases collection:

## 903 - Access to source databases:

- 904     o French Natural and Landscape Information System (SINP, <https://inpn.mnhn.fr>), each of the
- 905     three study sites has its own database: <https://sinp-occitanie.fr/atlas/> for Lodévois-Larzac,
- 906     ([T1](https://obv-na.fr/)), <https://obv-na.fr/> and <https://observatoire-fauna.fr/> for ~~Brocéliande~~, [La Rochelle \(T2\)](https://data.biodiversite-bretagne.fr),
- 907     <https://data.biodiversite-bretagne.fr> for ~~La Rochelle~~ [Brocéliande \(T3\)](https://data.biodiversite-bretagne.fr).
- 908     o GBIF: GBIF.org (04 October 2022) GBIF Occurrence Download
- 909     <https://doi.org/10.15468/dl.ry6uw7>
- 910     o French biodiversity monitoring scheme (Vigie Nature): <https://www.vigienature.fr/>, the
- 911     database used are the STOC for Aves, the Vigie-Chiro for Chiroptera and Orthoptera, the
- 912     Vigie-Flore for Flora, the STELI for Odonata, the STERF for Papilionidae
- 913 - Period of ~~1011~~ years, from 01/01/2010 to 31/12/2020
- 914 - Extent: continental France (without island, Corsica and overseas regions)
- 915 - Spatial accuracy of 0 to 50m

916 Standardization of species names : French taxonomic reference TAXREF.V14 ~~(Gargominy et al.,~~

917 ~~2021)~~[\(Gargominy et al., 2021\)](#).

- 918 - Keep only the species and subspecies
- 919 - Groups data at the species level

## 920 Taxonomic filtering: from TAXREF.V14

- 921 - Keep only species in nine taxonomic groups selected: Amphibia, Aves, Chiroptera, Flora, Mammalia
- 922     aptera (Mammalia), Orthoptera, Odonata, Papilionidae and Reptilia
- 923 - Delete species not present in France so keep: "Present", "Endemic", "Subendemic", "Cryptogenic",
- 924     "Introduced", "Invasive introduced"

## 925 Data transformation and filtering:

- 926 - Transformation data into occurrence, i.e. transform data into presence only
- 927 - Geographical filtering for reduce sampling biases and delete duplicate data: spatial thinning at a
- 928     distance of 50m with "spThin" package ~~(Aiello-Lammens et al., 2015)~~[\(Aiello-Lammens et al., 2015\)](#)

929 Identify the presence of species into each study site: at least five observations from all databases in the

930 study site to certify the presence of the species

931 Standardization of the observation period between databases: for Aves filtering observations between 1

932 March and 15 July corresponding to the STOC period.

933 A.2 - Environmental variables selected for Species Distribution Models (SDM) of Aves and Papilionidae species. Variables used for SDM are the variables with  
 934 collinearity removed (Pearson >0.7).

Type of variables	Variables	Database	Year	Raw precision	Source	Variables used for SDM
Climatic	19 bioclimatic variables	Chelsa	1981-2010	1 km	Karger et al., 2021, 2017	Bio 2 (temp. range), Bio 5 (temp. warmest month), Bio 6 (min. temp. coldest month), Bio 9 (temp. driest quarter), Bio 13 (prec. wettest month), Bio 14 (prec. driest month), Bio 15(prec. seasonality)
Biogeography/Geography	Elevation	BD ALTI, IGN	2014	25 m	IGN, 2022	Elevation
	Slope, mean in a buffer of 50 meters					Slope
	Topographic Position Index (TPI), mean in a buffer of 50 meters					TPI
Wetland	Area of wetland potential: very high in a buffer of 50, 500, 2000 meters	INRAE Wetland potential	2014	50 m	Berthier et al., 2014	Wetland in buffer of 50 and 500 m
Waterway	Linear meters of waterway in a buffer of 50, 500, 2000 meters	BD TOPO, IGN	2022	Vector	IGN, 2022	Waterway in buffer of 50 and 500m
Human occupancy	Land-use	OSO Land Cover	2018	25 m	Inglada et al., 2019	Artificial, conifer, crops, deciduous, fruit culture, land, mineral, prairie, water in a buffer of 50 and 500 m
	Hedge line	BD TOPO, IGN	2022	Vector	IGN, 2022	Lawn in a buffer of 500 m, Sand in a buffer of 50, 500, 2000 m
	NDVI mean	DHI NDVI	2022	10 m	CESBIO, 2021, 2022	Hedge line in buffer of 50, 500 m
Pollution / Fragmentation	Light pollution	VIIRS night time lights	2015	500 m	Elvidge et al., 2021	DHI NDVI in buffer of 50 and 500m
	Road fragmentation	Linear meters of fragmenting of main road in a buffer of 50, 500, 2000 meters	BD TOPO, IGN	2022	Vector	IGN, 2022
						Main road in a buffer of 50, 500, 2000 m

935

936 A.3 - Species traits of Aves and Papilionidae. Missing traits were imputed by considering evolutionary  
 937 relationships following the process of Carmona et al. (2021) with the help the R scripts of Toussaint et  
 938 al. (2021).

	Traits	Description	Reference
<b>Aves</b>	Mass	Body mass using data from males, females, and/or unspecified adults in g	Storchová and Hořák, 2018
	HWI	Avian hand-wing index (HWI), an estimate of wing shape as a proxy for dispersal ability in birds	Sheard et al., 2020
	Nocturn	Nocturnal activity	Wilman et al., 2014
	Habitats (Deciduous Coniferous, Woodland, Shrub, Grassland, Mountain meadows, Reed, Swamps, Rocks, Urban)	Species occupies habitat in breeding area, 1: yes; 0: no	Storchová and Hořák, 2018
	Spe. Diet, Spe. Foraging behav., Spe. Diet strat, Spe. Hab., Spe. Nest, Spe. Mean	Specialization respectively of diet, foraging behaviour, foraging substrate, habitat, nesting site and mean of five specializations	Morelli et al., 2020
<b>Papilionidae</b>	Wingspan	Average length of male wingspan in mm	Middleton-Welling et al., 2020
	FMoMean	Average duration of yearly flight period	
	HostplantN	Hostplant specificity, number of hostplants	
	HostplantSpe	Hostplant specificity index	Essens et al., 2017
	HPG	Hostplant growth form: Short herb/grass (<1 m) (Bi), tall herb/grass (>1 m) (Th), shrub (Sb), tree (Tr), liana (Li)	
	SSI	Species Specialization Index	Dupont et al., 2013
	AltVeg	Altitudinal vegetation: A=Anthropogenic, Co=Foothill, TheMeMed=Thermo/Meso-Mediterranean, Med=Mediterranean, SupMed=Supra-Mediterranean, Mo=Montane, ASa=Alpine and Subalpine	
Rarity	Rarity in France between 1 very rare to 3 common species	In Moussus et al., 2019 from oreina.org and faune.france.org	

The phylogenetic trees used to consider evolutionary relationships were from Jetz et al. (2012) and Wiemers et al. (2019, 2020) for the Aves and Papilionidae, respectively.

939

940



941 **Appendix B - Supplementary results**

942 *B.1 - Description of observation databases (Local SINP, GBIF, Vigie Nature) available nine taxonomic groups*  
 943 *in the three study sites (T1: Lodévois-Larzac, T2: Brocéliande, T3: La Rochelle) for “Local” (study*  
 944 *site with a buffer zone 10 km), “Regional” (administrative region) and “FranceCountry” (continental*  
 945 *France) scales. Statistics: “nSp” is the number of species with at least three observations on the*  
 946 *study site within a 10 km buffer; “pSp<15obs”, “pSp15.50obs” and “pSp>50obs” are the percentage of*  
 947 *species with <15, 15 to 50 and >50 observations respectively and “-” represents no data available.*

948 *B.1.A - Local scale (10km buffer zone).*


Taxa	Statistic	Local											
		GBIF			SINP			Vigie Nature			All databases combined		
		T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3
Amphibia	nSp	11	1	11	13	2	0				14	2	11
	pSp<15obs	82%	100%	100%	23%	50%	-	No program			29%	50%	100%
	pSp15.50obs	18%	0%	0%	15%	50%	-				7%	50%	0%
	pSp>50obs	0%	0%	0%	62%	0%	-				64%	0%	0%
Aves	nSp	100	73	77	197	94	82	114	155	101	205	171	132
	pSp<15obs	87%	93%	95%	37%	96%	85%	56%	50%	54%	37%	47%	59%
	pSp15.50obs	10%	6%	4%	23%	3%	15%	32%	17%	20%	22%	22%	13%
	pSp>50obs	3%	1%	1%	40%	1%	0%	12%	33%	27%	41%	31%	28%
Chiroptera	nSp	4	0	5	21	6	18	25	22	24	26	22	26
	pSp<15obs	100%	-	100%	10%	83%	50%	12%	41%	21%	12%	41%	27%
	pSp15.50obs	0%	-	0%	38%	0%	44%	16%	36%	46%	19%	36%	34%
	pSp>50obs	0%	-	0%	52%	17%	6%	72%	23%	33%	69%	23%	39%
Flora	nSp	786	476	639	1703	1335	843	286	177	136	1844	1460	1093
	pSp<15obs	81%	85%	81%	52%	48%	74%	100%	100%	100%	52%	50%	72%
	pSp15.50obs	16%	14%	17%	30%	25%	14%	0%	0%	0%	28%	24%	17%
	pSp>50obs	3%	1%	3%	18%	27%	12%	0%	0%	0%	20%	25%	11%
Mammalia	nSp	10	1	14	23	7	30				23	7	30
	pSp<15obs	100%	100%	100%	78%	86%	60%	No program			74%	86%	57%
	pSp15.50obs	0%	0%	0%	17%	14%	20%				22%	14%	20%
	pSp>50obs	0%	0%	0%	4%	0%	20%				4%	0%	23%
Odonata	nSp	38	5	23	57	1	0	0	1	24	57	7	30
	pSp<15obs	74%	100%	100%	28%	100%	-	-	100%	100%	28%	100%	97%
	pSp15.50obs	26%	0%	0%	37%	0%	-	-	0%	0%	30%	0%	3%
	pSp>50obs	0%	0%	0%	35%	0%	-	-	0%	0%	42%	0%	0%
Orthoptera	nSp	53	3	24	91	8	0	25	19	19	98	25	34
	pSp<15obs	92%	100%	100%	71%	100%	-	24%	58%	16%	56%	68%	53%
	pSp15.50obs	8%	0%	0%	23%	0%	-	24%	32%	52%	18%	24%	29%
	pSp>50obs	0%	0%	0%	6%	0%	-	52%	10%	32%	26%	8%	18%
Papilionidae	nSp	115	15	33	154	11	0	0	0	50	154	18	52
	pSp<15obs	60%	100%	94%	21%	100%	-	-	-	88	21%	100%	73%
	pSp15.50obs	30%	0%	6%	19%	0%	-	-	-	12	18%	0%	27%
	pSp>50obs	10%	0%	0%	60%	0%	-	-	-	0	60%	0%	0%
Reptilia	nSp	18	2	7	21	4	0				22	4	7
	pSp<15obs	67%	100%	100%	24%	100%	-	No program			27%	75%	100%
	pSp15.50obs	28%	0%	0%	33%	0%	-				32%	25%	0%
	pSp>50obs	6%	0%	0%	43%	0%	-				41%	0%	0%

949 *B.1.A - Regional scale.*

Taxa	Statistic	Regional								
		GBIF			Vigie Nature			All databases combined		
		T1	T2	T3	T1	T2	T3	T1	T2	T3
Amphibia	nSp	13	2	11				14	2	11
	pSp<15obs	8%	0%	73%	No program			7%	0%	73%
	pSp15.50obs	23%	50%	18%				14%	50%	18%
	pSp>50obs	69%	50%	9%				79%	50%	9%
Aves	nSp	198	159	120	182	143	120	205	166	132
	pSp<15obs	21%	21%	40%	31%	29%	32%	12%	16%	19%
	pSp15.50obs	12%	12%	22%	21%	15%	27%	11%	10%	28%
	pSp>50obs	67%	67%	38%	48%	56%	41%	77%	74%	53%
Chiroptera	nSp	13	13	5	25	22	24	26	22	26
	pSp<15obs	100%	100%	100%	0%	0%	8%	0%	0%	15%
	pSp15.50obs	0%	0%	0%	4%	0%	17%	8%	0%	15%
	pSp>50obs	0%	0%	0%	96%	100%	75%	92%	100%	69%
Flora	nSp	1428	1093	825	825	657	413	1844	1158	1093
	pSp<15obs	43%	38%	47%	95%	94%	94%	32%	40%	47%
	pSp15.50obs	19%	19%	20%	5%	6%	6%	26%	19%	23%
	pSp>50obs	38%	49%	33%	0%	0%	0%	42%	41%	31%
Mammalia	nSp	23	6	21				23	6	30
	pSp<15obs	39%	33%	62%	No program			30%	33%	50%
	pSp15.50obs	26%	0%	33%				22%	0%	23%
	pSp>50obs	35%	67%	5%				48%	67%	27%
Odonata	nSp	56	7	28	56	7	27	57	7	30
	pSp<15obs	18%	14%	64%	77%	71%	100%	9%	14%	53%
	pSp15.50obs	30%	14%	25%	23%	29%	0%	21%	14%	37%
	pSp>50obs	52%	71%	11%	0%	0%	0%	70%	72%	10%
Orthoptera	nSp	93	25	26	25	19	19	98	25	34
	pSp<15obs	42%	24%	58%	4%	0%	11%	30%	4%	27%
	pSp15.50obs	39%	40%	42%	4%	0%	16%	22%	0%	29%
	pSp>50obs	19%	36%	0%	92%	100%	74%	48%	96%	44%
Papilionidae	nSp	144	18	43	132	18	51	154	18	52
	pSp<15obs	12%	0%	37%	42%	11%	55%	7%	0%	34%
	pSp15.50obs	19%	11%	42%	30%	22%	41%	13%	6%	31%
	pSp>50obs	69%	89%	21%	28%	67%	4%	81%	94%	35%
Reptilia	nSp	20	4	7				22	4	7
	pSp<15obs	20%	0%	14%	No program			18%	0%	14%
	pSp15.50obs	30%	25%	57%				28%	25%	57%
	pSp>50obs	50%	75%	29%				59%	75%	29%

Taxa	Statistic	GBIF			<i>FranceCountry</i> Vigie Nature			All databases combined		
		T1	T2	T3	T1	T2	T3	T1	T2	T3
Amphibia	nSp	14	2	11	No program			14	2	11
	pSp<15obs	7%	0%	0%				0%	0%	0%
	pSp15.50obs	0%	0%	9%				7%	0%	9%
	pSp.>50obs	93%	100%	91%				93%	100%	91%
Aves	nSp	204	170	131	201	170	132	205	171	132
	pSp<15obs	2%	1%	1%	8%	9%	4%	1%	0%	0%
	pSp15.50obs	3%	1%	1%	11%	9%	7%	1%	1%	0%
	pSp.>50obs	95%	98%	98%	81%	82%	89%	98%	99%	100%
Chiroptera	nSp	19	19	21	25	22	25	26	22	26
	pSp<15obs	90%	89%	90%	0%	0%	0%	0%	0%	4%
	pSp15.50obs	10%	11%	10%	0%	0%	0%	4%	0%	0%
	pSp.>50obs	0%	0%	0%	100%	100%	100%	96%	100%	96%
Flora	nSp	1573	1261	995	1336	1105	848	1844	1460	1093
	pSp<15obs	25%	20%	17%	68%	61%	53%	19%	14%	16%
	pSp15.50obs	18%	18%	16%	19%	23%	27%	21%	17%	16%
	pSp.>50obs	57%	62%	67%	13%	16%	20%	60%	69%	68%
Mammalia	nSp	23	6	28	No program			23	7	30
	pSp<15obs	13%	16%	18%				9%	14%	23%
	pSp15.50obs	17%	17%	28%				13%	29%	17%
	pSp.>50obs	70%	67%	54%				78%	57%	60%
Odonata	nSp	57	7	30	57	7	30	57	7	30
	pSp<15obs	2%	0%	0%	11%	0%	0%	0%	0%	0%
	pSp15.50obs	10%	14%	0%	26%	0%	13%	3%	0%	0%
	pSp.>50obs	88%	86%	100%	63%	100%	87%	97%	100%	100%
Orthoptera	nSp	95	25	34	25	19	19	98	25	34
	pSp<15obs	22%	12%	6%	4%	0%	0%	15%	0%	0%
	pSp15.50obs	28%	20%	21%	0%	0%	0%	25%	4%	9%
	pSp.>50obs	50%	68%	73%	96%	100%	100%	60%	96%	91%
Papilionidae	nSp	147	18	52	149	18	52	154	18	52
	pSp<15obs	4%	0%	0%	13%	5%	0%	1%	0%	0%
	pSp15.50obs	8%	0%	0%	18%	6%	0%	7%	0%	0%
	pSp.>50obs	88%	100%	100%	69%	89%	100%	92%	100%	100%
Reptilia	nSp	21	4	7	No program			22	4	7
	pSp<15obs	0%	0%	0%				4%	0%	0%
	pSp15.50obs	14%	0%	0%				5%	0%	0%
	pSp.>50obs	86%	100%	100%				91%	100%	100%


953 B.2 - SDM evaluation of Aves and Papilionidae taxonomic groups for (A) three individual database sources  
 954 and (B) between SDM using different pseudo-absence sources. Each graphic is a pair of databases-  
 955 (National combined GBIF and Vigie Nature). Each point is an individual species that is listed in both  
 956 databases, some species are assessed only in one database (missing). A Boyce's index threshold of 0.3  
 957 is used to assess the proportion of satisfactory and unsatisfactory models conjointly for each of the two  
 958 database sources. The blue quadrant contains species whose SDMs are evaluated in a satisfactory  
 959 manner in both data bases (Boyce index > 0.3 for both databases) ; the red quadrant contains the  
 960 species for which SDMs are evaluated in an unsatisfactory manner in both databases (Boyce index <  
 961 0.3 for both databases). In the top-left quadrant and the bottom-right quadrants, SDM model  
 962 evaluation was satisfactory for only one database (on the in y and x axes respectively).



963  
 964

965  
966


B.3 - Overlap between SDM from individual database sources. VN is Vigie Nature database and National combined GBIF and Vigie Nature.



967


968  
969

B.4 - Overlap between SDM from individual database sources and SDM from mixed databases (i.e. using absence from Vigie Nature). National combine of GBIF and Vigie Nature.



970

971 B.5 - Maps of conservation planning area for Aves and Papilionidae of tree study sites (T1, T2, T3) from  
 972 different individual database source (GBIF, Vigie Nature and "National databases, GBIF, Vigie Nature"  
 973 combining the two databases).



974  
 975

976 B.6 - Evaluation of the quality of the full model (Table 5) compared to null model using AIC.

Model	Aves	Papilionidae
Full	3057	1322
Null	3357	1574

977



978  
979  
980  
981

B.7 – Species list observed in each study site (T1, T2, T3) and their French status from TAXREF.V14 (Presence, Introduced, Invasive introduced) and IUCN regional red list from BD.STATUT.V16 (LC, NT, VU, EN, CR, DD, NA) for A) Aves species and B) Papilionidae Species.

B.7.A – Aves species

Species	Observed			French status	IUCN regional red list		
	T1	T2	T3		T1	T2	T3
<u><i>Accipiter gentilis</i></u>	Yes	Yes	Yes	Present	LC	VU	EN
<u><i>Accipiter nisus</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Acrocephalus arundinaceus</i></u>	Yes	Yes	-	Present	VU	CR	-
<u><i>Acrocephalus schoenobaenus</i></u>	-	Yes	Yes	Present	-	VU	LC
<u><i>Acrocephalus scirpaceus</i></u>	Yes	Yes	-	Present	NT	VU	-
<u><i>Actitis hypoleucos</i></u>	Yes	Yes	Yes	Present	EN	CR	NA
<u><i>Aegithalos caudatus</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Aegypius monachus</i></u>	Yes	-	-	Present	CR	-	-
<u><i>Aix galericulata</i></u>	Yes	-	-	Introduced	NA	-	-
<u><i>Alauda arvensis</i></u>	Yes	Yes	Yes	Present	LC	VU	LC
<u><i>Alcedo atthis</i></u>	Yes	Yes	Yes	Present	NT	NT	LC
<u><i>Alectoris rufa</i></u>	Yes	Yes	Yes	Present	DD	DD	DD
<u><i>Anas acuta</i></u>	-	-	Yes	Present	-	-	VU
<u><i>Anas crecca</i></u>	-	-	Yes	Present	-	-	LC
<u><i>Anas platyrhynchos</i></u>	Yes	Yes	Yes	Present	DD	LC	LC
<u><i>Anser anser</i></u>	-	Yes	Yes	Present	-	VU	NA
<u><i>Anthus campestris</i></u>	Yes	Yes	-	Present	VU	EN	-
<u><i>Anthus pratensis</i></u>	Yes	Yes	Yes	Present	VU	EN	VU
<u><i>Anthus spinoletta</i></u>	-	Yes	Yes	Present	-	LC	LC
<u><i>Anthus trivialis</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Apus apus</i></u>	Yes	Yes	Yes	Present	LC	NT	LC
<u><i>Aquila chrysaetos</i></u>	Yes	-	-	Present	VU	-	-
<u><i>Aquila fasciata</i></u>	Yes	-	-	Present	CR	-	-
<u><i>Ardea alba</i></u>	Yes	Yes	Yes	Present	VU	NT	EN
<u><i>Ardea cinerea</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Ardea purpurea</i></u>	Yes	Yes	-	Present	EN	VU	-
<u><i>Ardeola ralloides</i></u>	Yes	-	-	Present	VU	-	-
<u><i>Arenaria interpres</i></u>	-	Yes	-	Present	-	NA	-
<u><i>Asio otus</i></u>	Yes	Yes	-	Present	LC	LC	-
<u><i>Athene noctua</i></u>	Yes	Yes	Yes	Present	NT	NT	VU
<u><i>Aythya ferina</i></u>	-	-	Yes	Present	-	-	EN
<u><i>Aythya fuligula</i></u>	-	-	Yes	Present	-	-	LC
<u><i>Botaurus stellaris</i></u>	Yes	-	-	Present	EN	-	-
<u><i>Branta bernicla</i></u>	-	Yes	Yes	Present	-	NA	LC
<u><i>Bubo bubo</i></u>	Yes	-	-	Present	LC	-	-
<u><i>Bubulcus ibis</i></u>	Yes	Yes	Yes	Present	LC	LC	NT
<u><i>Burhinus oedicnemus</i></u>	Yes	Yes	-	Present	LC	NT	-
<u><i>Buteo buteo</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Calandrella brachydactyla</i></u>	Yes	-	-	Present	EN	-	-
<u><i>Calidris alba</i></u>	-	Yes	-	Present	-	NA	-
<u><i>Calidris alpina</i></u>	-	Yes	-	Present	-	NA	-
<u><i>Caprimulgus europaeus</i></u>	Yes	-	Yes	Present	LC	-	LC
<u><i>Carduelis carduelis</i></u>	Yes	Yes	Yes	Present	VU	NT	LC
<u><i>Cecropis daurica</i></u>	Yes	-	-	Present	VU	-	-
<u><i>Certhia brachydactyla</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Certhia familiaris</i></u>	Yes	-	-	Present	LC	-	-
<u><i>Cettia cetti</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Charadrius alexandrinus</i></u>	-	Yes	-	Present	-	EN	-
<u><i>Charadrius dubius</i></u>	Yes	Yes	-	Present	NT	VU	-
<u><i>Charadrius hiaticula</i></u>	Yes	Yes	-	Present	VU	VU	-
<u><i>Chlidonias niger</i></u>	Yes	Yes	-	Present	NA	CR	-
<u><i>Chloris chloris</i></u>	Yes	Yes	Yes	Present	NT	NT	LC
<u><i>Chroicocephalus ridibundus</i></u>	Yes	Yes	Yes	Present	LC	VU	LC
<u><i>Ciconia ciconia</i></u>	Yes	Yes	-	Present	NT	NT	-
<u><i>Ciconia nigra</i></u>	Yes	-	-	Present	EN	-	-
<u><i>Cinclus cinclus</i></u>	Yes	-	-	Present	LC	-	-
<u><i>Circaetus gallicus</i></u>	Yes	Yes	-	Present	LC	EN	-
<u><i>Circus aeruginosus</i></u>	Yes	Yes	-	Present	VU	VU	-
<u><i>Circus cyaneus</i></u>	Yes	Yes	Yes	Present	EN	NT	EN
<u><i>Circus pygargus</i></u>	Yes	Yes	-	Present	EN	NT	-
<u><i>Cisticola juncidis</i></u>	Yes	Yes	Yes	Present	LC	NT	LC
<u><i>Clamator glandarius</i></u>	Yes	-	-	Present	NT	-	-

<u>Coccothraustes coccothraustes</u>	Yes	Yes	Yes	Present	LC	NT	VU
<u>Columba livia</u>	Yes	Yes	Yes	Present	DD	DD	DD
<u>Columba oenas</u>	Yes	Yes	Yes	Present	VU	EN	LC
<u>Columba palumbus</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Coracias garrulus</u>	Yes	-	-	Present	NT	-	-
<u>Corvus corax</u>	Yes	-	-	Present	LC	-	-
<u>Corvus corone</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Corvus frugilequus</u>	-	Yes	Yes	Present	-	LC	LC
<u>Corvus monedula</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Coturnix coturnix</u>	Yes	Yes	Yes	Present	NT	VU	LC
<u>Cuculus canorus</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Cyanistes caeruleus</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Cyanus olor</u>	Yes	Yes	-	Present	NA	LC	-
<u>Delichon urbicum</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Dendrocopos major</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Dendrocopos minor</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Dryocopus martius</u>	Yes	-	Yes	Present	LC	-	LC
<u>Egretta garzetta</u>	Yes	Yes	Yes	Present	LC	LC	NT
<u>Elanus caeruleus</u>	-	Yes	-	Present	-	VU	-
<u>Emberiza calandra</u>	Yes	Yes	Yes	Present	LC	VU	EN
<u>Emberiza cirius</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Emberiza citrinella</u>	Yes	Yes	Yes	Present	NT	NT	NT
<u>Emberiza hortulana</u>	Yes	-	-	Present	VU	-	-
<u>Emberiza schoeniclus</u>	Yes	Yes	Yes	Present	EN	EN	VU
<u>Erithacus rubecula</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Falco columbarius</u>	-	Yes	-	Present	-	NA	-
<u>Falco naumanni</u>	Yes	-	-	Present	VU	-	-
<u>Falco peregrinus</u>	Yes	Yes	-	Present	VU	CR	-
<u>Falco subbuteo</u>	Yes	Yes	Yes	Present	NT	NT	NT
<u>Falco tinnunculus</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Falco vespertinus</u>	Yes	Yes	-	Present	NA	NA	-
<u>Ficedula albicollis</u>	Yes	-	-	Present	NT	-	-
<u>Ficedula hypoleuca</u>	Yes	-	-	Present	EN	-	-
<u>Fringilla coelebs</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Fringilla montifringilla</u>	-	-	Yes	Present	-	-	DD
<u>Fulica atra</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Galerida cristata</u>	Yes	Yes	-	Present	LC	LC	-
<u>Gallinago gallinago</u>	Yes	Yes	Yes	Present	CR	CR	NA
<u>Gallinula chloropus</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Garrulus glandarius</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Grus grus</u>	Yes	Yes	-	Present	CR	CR	-
<u>Gyps fulvus</u>	Yes	-	-	Present	VU	-	-
<u>Hieraetus pennatus</u>	Yes	-	-	Present	VU	-	-
<u>Himantopus himantopus</u>	Yes	Yes	-	Present	LC	NT	-
<u>Hippolais polyglotta</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Hirundo rustica</u>	Yes	Yes	Yes	Present	NT	NT	LC
<u>Ichthyaetus melanocephalus</u>	-	Yes	-	Present	-	CR	-
<u>Jynx torquilla</u>	Yes	Yes	-	Present	NT	VU	-
<u>Lanius collurio</u>	Yes	Yes	Yes	Present	NT	NT	EN
<u>Lanius excubitor</u>	-	-	Yes	Present	-	-	NA
<u>Lanius meridionalis</u>	Yes	-	-	Present	EN	-	-
<u>Lanius senator</u>	Yes	-	-	Present	NT	-	-
<u>Larus argentatus</u>	-	Yes	Yes	Present	-	VU	VU
<u>Larus canus</u>	-	Yes	-	Present	-	EN	-
<u>Larus fuscus</u>	Yes	Yes	Yes	Present	NA	LC	LC
<u>Larus marinus</u>	-	Yes	-	Present	-	EN	-
<u>Larus michahellis</u>	Yes	Yes	-	Present	LC	VU	-
<u>Limosa lapponica</u>	-	Yes	-	Present	-	NA	-
<u>Limosa limosa</u>	-	Yes	-	Present	-	CR	-
<u>Linaria cannabina</u>	Yes	Yes	Yes	Present	NT	NT	LC
<u>Locustella luscinioides</u>	-	Yes	-	Present	-	EN	-
<u>Locustella naevia</u>	Yes	Yes	Yes	Present	DD	VU	LC
<u>Lophophanes cristatus</u>	Yes	Yes	Yes	Present	LC	VU	LC
<u>Loxia curvirostra</u>	Yes	-	-	Present	LC	-	-
<u>Lullula arborea</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Luscinia megarhynchos</u>	Yes	Yes	Yes	Present	LC	LC	VU
<u>Luscinia svecica</u>	Yes	Yes	-	Present	LC	LC	-
<u>Mareca penelope</u>	-	-	Yes	Present	-	-	LC
<u>Mareca strepera</u>	-	-	Yes	Present	-	-	LC
<u>Merqus merqanser</u>	-	-	Yes	Present	-	-	NA
<u>Merops apiaster</u>	Yes	Yes	-	Present	NT	VU	-

<u>Milvus migrans</u>	Yes	Yes	-	Present	LC	LC	-
<u>Milvus milvus</u>	Yes	-	-	Present	EN	-	-
<u>Monticola saxatilis</u>	Yes	-	-	Present	VU	-	-
<u>Monticola solitarius</u>	Yes	-	-	Present	VU	-	-
<u>Motacilla alba</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Motacilla cinerea</u>	Yes	-	Yes	Present	LC	-	LC
<u>Motacilla flava</u>	Yes	Yes	-	Present	NT	LC	-
<u>Muscicapa striata</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Neophron percnopterus</u>	-	Yes	-	Present	-	EN	-
<u>Numenius arquata</u>	Yes	Yes	-	Present	CR	EN	-
<u>Numenius phaeopus</u>	-	Yes	-	Present	-	NA	-
<u>Nycticorax nycticorax</u>	Yes	Yes	-	Present	NT	VU	-
<u>Oenanthe oenanthe</u>	Yes	Yes	Yes	Present	NT	EN	EN
<u>Oriolus oriolus</u>	Yes	Yes	-	Present	LC	LC	-
<u>Otus scops</u>	Yes	-	-	Present	NT	-	-
<u>Pandion haliaetus</u>	Yes	-	-	Present	VU	-	-
<u>Parus major</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Passer domesticus</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Passer montanus</u>	Yes	Yes	Yes	Present	NT	EN	EN
<u>Perdix perdix</u>	Yes	Yes	Yes	Present	NT	DD	DD
<u>Periparus ater</u>	Yes	-	Yes	Present	LC	-	NT
<u>Pernis apivorus</u>	Yes	Yes	-	Present	LC	VU	-
<u>Petronia petronia</u>	Yes	-	-	Present	LC	-	-
<u>Phalacrocorax carbo</u>	Yes	Yes	Yes	Present	NA	VU	LC
<u>Phasianus colchicus</u>	Yes	Yes	Yes	Introduced	NA	DD	DD
<u>Phoenicurus ochrurus</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Phoenicurus phoenicurus</u>	Yes	Yes	Yes	Present	LC	LC	VU
<u>Phylloscopus bonelli</u>	Yes	Yes	-	Present	LC	NT	-
<u>Phylloscopus collybita</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Phylloscopus sibilatrix</u>	Yes	-	Yes	Present	EN	-	NT
<u>Phylloscopus trochilus</u>	Yes	Yes	Yes	Present	NA	CR	EN
<u>Pica pica</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Picus canus</u>	Yes	Yes	-	Present	EN	CR	-
<u>Picus viridis</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Platalea leucorodia</u>	-	Yes	-	Present	-	EN	-
<u>Pluvialis apricaria</u>	-	-	Yes	Present	-	-	LC
<u>Pluvialis squatarola</u>	-	Yes	-	Present	-	NA	-
<u>Podiceps cristatus</u>	Yes	-	Yes	Present	LC	-	LC
<u>Poecile palustris</u>	Yes	Yes	Yes	Present	LC	VU	NT
<u>Prunella collaris</u>	Yes	-	-	Present	EN	-	-
<u>Prunella modularis</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Ptyonoprogne rupestris</u>	Yes	-	-	Present	LC	-	-
<u>Pyrhhorcorax pyrrhhorcorax</u>	Yes	-	-	Present	VU	-	-
<u>Pyrrhula pyrrhula</u>	Yes	-	Yes	Present	VU	-	VU
<u>Rallus aquaticus</u>	Yes	Yes	-	Present	LC	VU	-
<u>Recurvirostra avosetta</u>	-	Yes	-	Present	-	VU	-
<u>Regulus ignicapilla</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Regulus regulus</u>	Yes	-	Yes	Present	LC	-	LC
<u>Riparia riparia</u>	Yes	Yes	Yes	Present	EN	NT	LC
<u>Saxicola rubetra</u>	Yes	Yes	-	Present	EN	CR	-
<u>Saxicola rubicola</u>	Yes	Yes	Yes	Present	VU	NT	LC
<u>Scolopax rusticola</u>	Yes	-	-	Present	DD	-	-
<u>Serinus serinus</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Sitta europaea</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Spatula clypeata</u>	Yes	Yes	Yes	Present	DD	VU	LC
<u>Spatula querquedula</u>	-	Yes	-	Present	-	CR	-
<u>Spinus spinus</u>	Yes	-	-	Present	VU	-	-
<u>Sterna hirundo</u>	Yes	Yes	-	Present	LC	VU	-
<u>Streptopelia decaocto</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Streptopelia turtur</u>	Yes	Yes	Yes	Present	LC	VU	LC
<u>Strix aluco</u>	Yes	Yes	Yes	Present	LC	LC	DD
<u>Sturnus vulgaris</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Sylvia atricapilla</u>	Yes	Yes	Yes	Present	LC	LC	LC
<u>Sylvia borin</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Sylvia cantillans</u>	Yes	Yes	-	Present	LC	LC	-
<u>Sylvia communis</u>	Yes	Yes	Yes	Present	LC	NT	LC
<u>Sylvia conspicillata</u>	-	Yes	-	Present	-	EN	-
<u>Sylvia hortensis</u>	Yes	-	-	Present	LC	-	-
<u>Sylvia melanocephala</u>	Yes	-	-	Present	LC	-	-
<u>Sylvia undata</u>	Yes	-	Yes	Present	VU	-	LC
<u>Tachybaptus ruficollis</u>	Yes	Yes	Yes	Present	LC	LC	LC

<u><i>Tachymarptis melba</i></u>	Yes	-	-	Present	VU	-	-
<u><i>Tadorna ferruginea</i></u>	-	Yes	-	Present	-	NA	-
<u><i>Tadorna tadorna</i></u>	Yes	Yes	-	Present	LC	LC	-
<u><i>Tetrax tetrax</i></u>	Yes	-	-	Present	NT	-	-
<u><i>Thalasseus sandvicensis</i></u>	Yes	Yes	-	Present	VU	NT	-
<u><i>Tichodroma muraria</i></u>	Yes	-	-	Present	CR	-	-
<u><i>Tringa erythropus</i></u>	-	Yes	-	Present	-	NA	-
<u><i>Tringa nebularia</i></u>	-	Yes	-	Present	-	NA	-
<u><i>Tringa ochropus</i></u>	Yes	Yes	Yes	Present	NA	NA	DD
<u><i>Tringa totanus</i></u>	Yes	Yes	-	Present	EN	VU	-
<u><i>Troglodytes troglodytes</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Turdus iliacus</i></u>	Yes	-	Yes	Present	NA	-	DD
<u><i>Turdus merula</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Turdus philomelos</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Turdus pilaris</i></u>	Yes	-	Yes	Present	VU	-	DD
<u><i>Turdus torquatus</i></u>	-	Yes	-	Present	-	LC	-
<u><i>Turdus viscivorus</i></u>	Yes	Yes	Yes	Present	LC	NT	LC
<u><i>Tyto alba</i></u>	-	Yes	Yes	Present	-	VU	DD
<u><i>Upupa epops</i></u>	Yes	Yes	Yes	Present	LC	LC	LC
<u><i>Vanellus vanellus</i></u>	-	Yes	Yes	Present	-	VU	VU

982

983 *B.7.A – Papilionidae species*

Species	Observed		French status	IUCN regional red list	
	T1	T3		T1	T3
<u><i>Aglais io</i></u>	Yes	Yes	Present	LC	LC
<u><i>Anthocharis cardamines</i></u>	Yes	Yes	Present	LC	LC
<u><i>Anthocharis euphenoides</i></u>	Yes	-	Present	LC	-
<u><i>Apatura ilia</i></u>	Yes	-	Present	LC	-
<u><i>Apatura iris</i></u>	Yes	Yes	Present	NT	NT
<u><i>Aphantopus hyperantus</i></u>	Yes	Yes	Present	LC	NT
<u><i>Aporia crataegi</i></u>	Yes	Yes	Present	LC	LC
<u><i>Araschnia levana</i></u>	Yes	Yes	Present	LC	LC
<u><i>Arethusana arethusa</i></u>	Yes	-	Present	LC	-
<u><i>Argynnis pandora</i></u>	Yes	-	Present	LC	-
<u><i>Argynnis paphia</i></u>	Yes	Yes	Present	LC	LC
<u><i>Aricia agestis</i></u>	Yes	Yes	Present	LC	LC
<u><i>Aricia montensis</i></u>	Yes	-	Present	DD	-
<u><i>Boloria dia</i></u>	Yes	Yes	Present	LC	LC
<u><i>Boloria euphrosyne</i></u>	Yes	-	Present	LC	-
<u><i>Boloria selene</i></u>	Yes	-	Present	NT	-
<u><i>Brenthis daphne</i></u>	Yes	-	Present	LC	-
<u><i>Brenthis hecate</i></u>	Yes	-	Present	VU	-
<u><i>Brintesia circe</i></u>	Yes	-	Present	LC	-
<u><i>Cacyreus marshalli</i></u>	Yes	-	Invasive introduced	NA	-
<u><i>Callophrys avis</i></u>	Yes	-	Present	LC	-
<u><i>Callophrys rubi</i></u>	Yes	Yes	Present	LC	LC
<u><i>Carcharodus alceae</i></u>	Yes	Yes	Present	LC	LC
<u><i>Celastrina argiolus</i></u>	Yes	Yes	Present	LC	LC
<u><i>Charaxes jasius</i></u>	Yes	-	Present	LC	-
<u><i>Chazara briseis</i></u>	Yes	-	Present	VU	-
<u><i>Coenonympha arcania</i></u>	Yes	Yes	Present	LC	LC
<u><i>Coenonympha dorus</i></u>	Yes	-	Present	LC	-
<u><i>Coenonympha pamphilus</i></u>	Yes	Yes	Present	LC	LC
<u><i>Colias alfacariensis</i></u>	Yes	Yes	Present	LC	LC
<u><i>Colias crocea</i></u>	Yes	Yes	Present	LC	LC
<u><i>Cupido alcetas</i></u>	Yes	-	Present	LC	-
<u><i>Cupido argiades</i></u>	-	Yes	Present	-	NT
<u><i>Cupido minimus</i></u>	Yes	-	Present	LC	-
<u><i>Cupido osiris</i></u>	Yes	-	Present	NT	-
<u><i>Cyaniris semiargus</i></u>	Yes	Yes	Present	LC	NT
<u><i>Erebia aethiops</i></u>	Yes	-	Present	NT	-
<u><i>Erebia epistyane</i></u>	Yes	-	Present	EN	-
<u><i>Erebia meolans</i></u>	Yes	-	Present	LC	-
<u><i>Erebia neoridas</i></u>	Yes	-	Present	NT	-
<u><i>Erynnis tages</i></u>	Yes	Yes	Present	LC	LC
<u><i>Euchloe crameri</i></u>	Yes	-	Present	LC	-
<u><i>Euphydryas aurinia</i></u>	Yes	-	Present	NT	-
<u><i>Fabriciana adippe</i></u>	Yes	-	Present	NT	-

<u><i>Fabriciana niobe</i></u>	Yes	-	Present	NT	-
<u><i>Glaucopsyche alexis</i></u>	Yes	-	Present	LC	-
<u><i>Glaucopsyche melanops</i></u>	Yes	-	Present	LC	-
<u><i>Gonepteryx cleopatra</i></u>	Yes	-	Present	LC	-
<u><i>Gonepteryx rhamni</i></u>	Yes	Yes	Present	LC	LC
<u><i>Hamearis lucina</i></u>	Yes	-	Present	LC	-
<u><i>Hesperia comma</i></u>	Yes	-	Present	LC	-
<u><i>Hipparchia alcyone</i></u>	Yes	-	Present	DD	-
<u><i>Hipparchia faqi</i></u>	Yes	-	Present	LC	-
<u><i>Hipparchia fidia</i></u>	Yes	Yes	Present	LC	LC
<u><i>Hipparchia qenava</i></u>	Yes	-	Present	NT	-
<u><i>Hipparchia semele</i></u>	Yes	Yes	Present	LC	EN
<u><i>Hipparchia statilinus</i></u>	Yes	Yes	Present	LC	EN
<u><i>Hyponephele lupina</i></u>	Yes	-	Present	EN	-
<u><i>Hyponephele lycaon</i></u>	Yes	-	Present	EN	-
<u><i>Iberochloe taais</i></u>	Yes	-	Present	EN	-
<u><i>Iphiclides podalirius</i></u>	Yes	-	Present	LC	-
<u><i>Isoria lathonia</i></u>	Yes	Yes	Present	LC	LC
<u><i>Laeosopis roboris</i></u>	Yes	-	Present	LC	-
<u><i>Lampides boeticus</i></u>	Yes	-	Present	LC	-
<u><i>Lasiommata maera</i></u>	Yes	Yes	Present	LC	CR
<u><i>Lasiommata megera</i></u>	Yes	Yes	Present	LC	LC
<u><i>Leptidea sinapis</i></u>	Yes	Yes	Present	LC	LC
<u><i>Leptotes pirithous</i></u>	Yes	-	Present	LC	-
<u><i>Libythea celtis</i></u>	Yes	-	Introduced	LC	-
<u><i>Limenitis camilla</i></u>	Yes	Yes	Present	LC	LC
<u><i>Limenitis reducta</i></u>	Yes	-	Present	LC	-
<u><i>Lycaena alciphron</i></u>	Yes	-	Present	LC	-
<u><i>Lycaena dispar</i></u>	Yes	-	Present	NT	-
<u><i>Lycaena phlaeas</i></u>	Yes	Yes	Present	LC	LC
<u><i>Lycaena tityrus</i></u>	Yes	Yes	Present	LC	LC
<u><i>Lysandra bellargus</i></u>	Yes	-	Present	LC	-
<u><i>Lysandra coridon</i></u>	Yes	-	Present	LC	-
<u><i>Lysandra hispana</i></u>	Yes	-	Present	LC	-
<u><i>Maniola jurtina</i></u>	Yes	Yes	Present	LC	LC
<u><i>Melanargia galathea</i></u>	Yes	Yes	Present	LC	LC
<u><i>Melanargia lachesis</i></u>	Yes	-	Present	LC	-
<u><i>Melanargia occitanica</i></u>	Yes	-	Present	LC	-
<u><i>Melanargia russiae</i></u>	Yes	-	Present	VU	-
<u><i>Melitaea athalia</i></u>	Yes	-	Present	DD	-
<u><i>Melitaea celadussa</i></u>	Yes	-	Present	LC	-
<u><i>Melitaea cinxia</i></u>	Yes	Yes	Present	LC	LC
<u><i>Melitaea deione</i></u>	Yes	-	Present	DD	-
<u><i>Melitaea diamina</i></u>	Yes	-	Present	NT	-
<u><i>Melitaea didyma</i></u>	Yes	-	Present	LC	-
<u><i>Melitaea parthenoides</i></u>	Yes	-	Present	LC	-
<u><i>Melitaea phoebe</i></u>	Yes	Yes	Present	LC	LC
<u><i>Minois dryas</i></u>	Yes	-	Present	LC	-
<u><i>Muschampia floccifera</i></u>	Yes	-	Present	NT	-
<u><i>Muschampia lavatherae</i></u>	Yes	-	Present	NT	-
<u><i>Muschampia proto</i></u>	Yes	-	Present	NT	-
<u><i>Nymphalis antiopa</i></u>	Yes	-	Present	NT	-
<u><i>Nymphalis polychloros</i></u>	Yes	Yes	Present	LC	LC
<u><i>Ochlodes sylvanus</i></u>	Yes	Yes	Present	LC	LC
<u><i>Papilio machaon</i></u>	Yes	Yes	Present	LC	LC
<u><i>Pararge aegeria</i></u>	Yes	Yes	Present	LC	LC
<u><i>Parnassius apollo</i></u>	Yes	-	Present	VU	-
<u><i>Phenacis alcon</i></u>	Yes	-	Present	VU	-
<u><i>Phenacis arion</i></u>	Yes	-	Present	NT	-
<u><i>Pieris brassicae</i></u>	Yes	Yes	Present	LC	LC
<u><i>Pieris manni</i></u>	Yes	-	Present	DD	-
<u><i>Pieris napi</i></u>	Yes	Yes	Present	LC	LC
<u><i>Pieris rapae</i></u>	Yes	Yes	Present	LC	LC
<u><i>Plebejus arqus</i></u>	Yes	Yes	Present	LC	NT
<u><i>Plebejus idas</i></u>	Yes	Yes	Present	NT	CR
<u><i>Polygonia c-album</i></u>	Yes	Yes	Present	LC	LC
<u><i>Polyommatus amandus</i></u>	Yes	-	Present	VU	-
<u><i>Polyommatus daphnis</i></u>	Yes	-	Present	VU	-
<u><i>Polyommatus dolus</i></u>	Yes	-	Present	VU	-
<u><i>Polyommatus dorylas</i></u>	Yes	-	Present	VU	-
<u><i>Polyommatus escheri</i></u>	Yes	-	Present	LC	-

<u><i>Polyommatus icarus</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Polyommatus thersites</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>DD</u>	-
<u><i>Pontia daplidice</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Pseudophilotes baton</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Pyraus alveus</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>NT</u>	-
<u><i>Pyraus armoricanus</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Pyraus carthami</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>NT</u>	-
<u><i>Pyraus cirsii</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>VU</u>	-
<u><i>Pyraus foulquieri</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>EN</u>	-
<u><i>Pyraus malvae</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>DD</u>	-
<u><i>Pyraus malvoides</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Pyraus onopordi</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>DD</u>	-
<u><i>Pyronia bathseba</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Pyronia cecilia</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Pyronia tithonus</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Quercusia quercus</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Satyrrium acaciae</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Satyrrium esculi</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Satyrrium ilicis</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Satyrrium pruni</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>DD</u>	-
<u><i>Satyrrium spini</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Satyrrium w-album</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Satyrus actaea</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>VU</u>	-
<u><i>Satyrus ferula</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>VU</u>	-
<u><i>Speyeria aqaja</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Spialia sertorius</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Thecla betulae</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Thymelicus acteon</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Thymelicus lineola</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Thymelicus sylvestris</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Vanessa atalanta</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Vanessa cardui</i></u>	<u>Yes</u>	<u>Yes</u>	<u>Present</u>	<u>LC</u>	<u>LC</u>
<u><i>Zerynthia polyxena</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-
<u><i>Zerynthia rumina</i></u>	<u>Yes</u>	-	<u>Present</u>	<u>LC</u>	-