

17 **Abstract** - Cues involved in mate seeking and recognition prevent hybridization and can be involved in
18 speciation processes. In malaria mosquitoes, females of the two sibling species *Anopheles gambiae s.s.* and *An.*
19 *coluzzii* mate in monospecific male swarms and hybrids are rare. Long-range sex pheromones driving this
20 behavior have been debated in literature but to date, no study has proven their existence or their absence. Here,
21 we attempted to bring to light their existence. To put all the odds in our favor, we used different chemical
22 ecology methods such as behavioral and electrophysiological assays as well chemical analyses, and we worked
23 with mosquitoes at their optimal physiological mating state *i.e.* with swarming males during their natural
24 swarming windows. Despite all our efforts, our results support the absence of long-range sex pheromones
25 involved in swarm detection and recognition by females. We briefly discuss the implications of this finding in
26 ecology, evolution and for control strategies.

27

28

29 **Key Words** - *Anopheles*, male swarm, mating behavior, mosquito control.

31 Cues involved in mate seeking and mate recognition prevent hybridization and can be involved in speciation
32 processes. In insects, these cues can be visual, acoustical or chemical with a short or a long-range action and are
33 highly species specific (Alexander et al. 1997; Clements 1999).

34 In mosquitoes, many species mate in flight, within individual aggregations called "mating swarms"
35 (Gibson 1945; Downes 1969; Savolainen 1978; Shelly and Whittier 1997; Sivinski and Petersson 1997). In
36 malaria mosquitoes of the *Anopheles gambiae* complex, mating takes place at sunset in monospecific swarms
37 containing a few males to thousands of them in which virgin conspecific females come to find a mate
38 (Charlwood and Jones 1980; Marchand 1984; Charlwood et al. 2002; Diabaté et al. 2003, 2009; Howell and
39 Knols 2009; Sawadogo et al. 2014). In West Africa, *An. coluzzii* and *An. gambiae s.s.* are often found in
40 sympatry but they form distinct swarms spatially segregated and hybrids are rare ($\approx 1\%$) (della Torre et al. 2001;
41 Tripet et al. 2001; della Torre et al. 2005; Diabaté et al. 2006; Costantini et al. 2009; Diabaté et al. 2009;
42 Sawadogo et al. 2013, 2014). No evidence for selection against hybrids was found and spermatheca analyses
43 showed that these species mostly mate assortatively (Persiani et al. 1986; Tripet et al. 2001; Diabaté et al. 2005,
44 2007; Hahn et al. 2012; Pombi et al. 2017). This suggests that reproductive isolation between these two sibling
45 species is achieved by strong pre-mating reproductive barriers (della Torre et al. 2001; Diabaté et al. 2007;
46 Lehmann and Diabaté 2008). In addition, since females usually only mate once in their lives (Clements 1992),
47 errors in the choice of mate should be costly and fall under negative selection. Consequently, one would expect
48 to find specific cues that lead females to conspecific male swarms. However, the way females are attracted to
49 swarms is unknown.

50 In species of the *An. gambiae* complex, several cues have been identified to play a role in bringing
51 sexes together. Acoustic cues were shown to be involved in close-range recognition, the male and the female
52 adjusting their respective wing-beat frequencies to converge on a shared harmonic frequency (Gibson et al.
53 2010; Pennetier et al. 2010). However, it was recently demonstrated that females of some *Anopheles* species are
54 able to detect swarm sounds only in a very close vicinity. Thus, they are unable to use swarm sound to locate or
55 identify swarms at long range (Feugère et al. 2021). Visual cues also play an important role in species
56 segregation with *An. coluzzii* males swarming over contrasted visual ground markers (marker, hereafter) and *An.*
57 *gambiae* males swarming over bare ground (Diabaté et al. 2009). However, a recent work showed that, like *An.*
58 *coluzzii*, *An. gambiae* males also use visual markers but rather to locate their swarm at a distance from the
59 marker (Poda et al. 2019). Moreover, females also use these markers to form swarms in the absence of males

60 suggesting that females may use these markers to join the swarm location (Poda et al. 2019). Nevertheless, the
61 distance between two heterospecific swarms using the same **visual ground** marker can be about 2 m in semi-field
62 conditions (Poda et al. 2019) and it is still unknown from which distance mosquitoes can see **such** markers,
63 suggesting that females could cross the location of heterospecific swarms by accident. Chemical cues have also
64 been under investigation in *Anopheles* species. Heptacosane, a cuticular hydrocarbon, enhances the interaction
65 between males and females (Wang et al. 2021). This compound, however, can only be perceived by contact and
66 can consequently be involved only in mate recognition during courtship and to stimulate acceptance by females.
67 Moreover, the absence of assortative mating in confined heterospecific males and females in laboratory cages or
68 indoor swarms (Dao et al. 2008), suggests that close-range mating cues, if they exist, cannot ensure total
69 reproductive isolation by themselves. Species isolation is thus likely to occur through long-range and specific
70 swarm recognition cues acting as a first barrier in pre-mating processes which prevent hybridization by limiting
71 contact between sexes of the different species.

72 To our knowledge, volatile sex pheromones in the *An. gambiae* complex have never been brought to
73 light without ambiguity. Charlwood et al. (2002) reported an absence of response of *An. gambiae* males in
74 natural swarms to squashed females on filter paper or to living females in a net cage. However, in natural
75 conditions, females are the ones attracted to male aggregation sites. Consequently, males should be the ones that
76 emit long-range pheromones and should be the attractive sex. Nevertheless, a laboratory study failed to
77 demonstrate any attractiveness of dead males to virgin females in a Y-tube olfactometer (Gomulski 1988).
78 However, the difficulties in highlighting the existence of such pheromones might be due to an emission in very
79 low quantities and/or exclusively during swarming. Indeed, in sandfly species, which also mate in large
80 aggregations of males, it was shown that the concentration of male-produced sex pheromones is greater in larger
81 swarms, and hence, the chances for individual males of mating with a conspecific female are also greater (Kelly
82 and Dye 1997; Bray et al. 2010). Similarly, Diabaté et al. (2011) showed that *An. coluzzii* females were more
83 frequently attracted to large swarms, suggesting a potential additive effect of cues released by males. Males
84 gathering in large swarms may increase their "detectability" in the female's olfactory landscape (Shelly and
85 Whittier 1997). In addition, in many insects, pheromone release and receptivity were shown to respond to diel
86 rhythms or even strict time windows of only a few minutes or hours (Bjostad et al. 1980; Merlin et al. 2007;
87 Rund et al. 2013; Levi-Zada et al. 2014) or even to be dependent of the presence of conspecifics (Andersson et
88 al. 2007; Robledo and Arzuffi 2012). Nevertheless, a recent work by Mozūraitis et al. (2020) and concomitant to
89 a first version of the present work (Poda et al. 2021), highlighted the existence of five volatile compounds

90 potentially emitted by *Anopheles* males. According to the authors, these might be involved in aggregation
91 behavior, attracting both males and females, and increasing the insemination rate. However, though an
92 interesting biological activity was described by the authors, the claim that these compounds are male swarming
93 aggregation pheromones must be considered with caution and it requires further investigations. Indeed, these
94 five compounds (acetoin, sulcatone, octanal, nonanal and decanal) are very frequently found in nature and, as
95 reported by the authors, have been shown to have a biological activity in *Anopheles* mosquitoes in very different
96 contexts from the reproductive behavior. They were repeatedly found in human and animal body odor (Verhulst
97 et al. 2010; Pandey and Kim 2011; Dormont et al. 2013b, a; Tchouassi et al. 2013; McBride et al. 2014), breath
98 (Poli et al. 2010; Calenic and Amann 2014; Filipiak et al. 2014; Cainap et al. 2020), and in host-plants (Dekel et
99 al. 2019). These represent the main sources of food (blood and sugar meals, respectively) for *Anopheles*
100 mosquitoes. As a matter of fact, octanal, nonanal and decanal are part of a blend used in some traps to mimic
101 mammalian host odor for several species of mosquito vectors (Tchouassi et al. 2013; Nyasembe et al. 2014).
102 Some of these compounds were also found associated to oviposition sites (Suh et al. 2016; Wondwosen et al.
103 2016, 2018) and to ambient air (Kostiainen 1995; Kruza et al. 2017). In addition, sulcatone is thought to be
104 responsible for discrimination between humans and animals in human-seeking mosquitoes (McBride et al. 2014).
105 Thus, it is not surprising that they trigger a strong biological/flight activity in mosquitoes.

106 Nevertheless, the discovery of sex pheromones or a highly attractive blend in malaria mosquitoes would
107 be a precious step toward the development of new control and monitoring strategies. Furthermore, sex or
108 aggregation pheromones could help in designing sexually competitive mosquitoes in sterile insect and gene drive
109 techniques. Here, we **investigated** the existence of long-range sex pheromones in *An. gambiae* and *An. coluzzii*
110 mosquitoes, that may allow the females to detect, recognize and track conspecific male swarms. On the premise
111 that such pheromones could be produced only by males during swarming windows and to put all the odds in our
112 favor, we used different chemical ecology methods, always with living swarming mosquitoes during their natural
113 swarming windows. **First, we investigated the long-range behavioral response of females exposed to the volatile**
114 **blend from male swarms in an olfactometer. Second, we collected and analyzed volatile organic compounds**
115 **(VOCs) with different methods on both laboratory-induced swarms and natural swarms. And third, we tested for**
116 **an antenna-electrophysiological response of females to male swarm VOCs.** As much as possible, we used both
117 recently colonized mosquitoes and large experimental set-ups to ensure males produced a free swarming
118 behavior. In addition, we replicated the main experiment by Mozūraitis et al. (2020) to search for the five

119 compounds. We specifically added two new controls, one to check if these compounds were male specific, and
120 the other to discard a potential environmental/laboratory pollution.

121

122

METHODS AND MATERIALS

123 Mosquitoes

124 Mosquitoes used in behavioral experiments and for both VOC extraction and electrophysiological analyses were
125 from colonies raised from wild gravid females collected in inhabited human dwellings in Burkina Faso (West
126 Africa). *Anopheles coluzzii* were collected in 2017 in Bama and *An. gambiae* in 2015 in Soumouso. Bama is a
127 village located in a rice-growing area located 30 km North of Bobo-Dioulasso (11°24'14"N; 04°24'42"W)
128 where previous studies showed that the *Anopheles* population is almost exclusively composed of *An. coluzzii*
129 (Mosqueira et al. 2015; Poda et al. 2018). Soumouso is a typical Guinean savannah village located 30 km
130 North-East of Bobo-Dioulasso (11°00'46"N, 4°02'45"W) where *An. coluzzii* and *An. gambiae* coexist, with a
131 predominance of the latter (Diabaté et al. 2004, 2006). Gravid females were placed individually in oviposition
132 cups containing tap water. After oviposition, females were identified to species by routine PCR-RFLP
133 (Santolamazza et al. 2008). The larvae were gathered according to their species and reared in tap water, fed with
134 Tetramin® Baby Fish Food (Tetrawerke, Melle, Germany) *ad libitum*. Adult mosquitoes were held in 30 × 30 ×
135 30 cm mesh-covered cages and provided with a 5% glucose solution *ad libitum*. Insectarium conditions were
136 maintained at 27±2 °C, 70±10% RH and 12L:12D. The colonies were refreshed twice a year with F1 from
137 mosquito females caught in the wild.

138 Mosquitoes used in the electrophysiological study were transferred as eggs from Burkina Faso to
139 France. On their arrival, eggs were allowed to hatch in osmosed water and the larvae were fed with Tetramin®
140 Baby Fish Food *ad libitum*. Adult mosquitoes were held in 20 × 20 × 20 cm mesh-covered cages and provided
141 with honey diluted at 5% *ad libitum*. Females were fed with rabbit blood on a PS6 Power Unit (Hemotek,
142 Blackburn, UK) for egg production. Mosquitoes were reared in a laboratory climate chamber KBF-S720
143 (BINDER Gmbh, Tuttlingen, Germany) at 27±2 °C, 70±10% RH and 12L:12D with a sunset time programmed
144 at 3:00pm (synchronization of swarming/mating time and electrophysiological test time to ensure an optimal
145 receptivity of females). Mosquitoes used for the experiments were sexed early after emergence and sexes were
146 kept in separate rearing cages to prevent mating.

147 Mosquitoes used to replicate the experiment by Mozūraitis et al. (2020) were from a 15-year-old colony
148 of *An. gambiae* (Kisumu strain) reared in the IRD laboratory of Montpellier. They were maintained at 27 ± 2 °C,
149 $80 \pm 10\%$ RH, with a photoperiod cycle of 12h light: 12h dark and reared as described above.

150

151 **Long-range behavioral response of virgin females to swarm volatile organic compounds**

152

153 **Olfactometer setup** - Bioassays were conducted in a dual-port olfactometer originally designed to study host
154 preference in the *An. gambiae* complex (Lefèvre et al. 2009, 2010; Vantaux et al. 2015; Nguyen et al. 2017). The
155 odor source container was modified to enclose male swarms in boxes made of transparent plexiglass ($L \times W \times$
156 $H: 60 \times 60 \times 120$ cm; "swarming boxes" hereafter). Each swarming box was connected by a PVC air vent hoses
157 ($L \times \text{Ø}: 600 \times 10$ cm, W3-65014-HQ4, HQ, USA) to a collecting glass box ($L \times W \times H: 32 \times 33.5 \times 44$ cm)
158 which was linked to a mesh-covered releasing cage ($L \times W \times H: 50 \times 40 \times 40$ cm) by a glass tube ($L \times \text{Ø}: 60 \times$
159 10 cm) (Fig. 1). A custom-made electric fan was located at the mid-length of each air vent hose and drew air
160 from the swarming boxes (odor-sources) to the releasing cage, providing an odor-laden air current against which
161 mosquitoes in the releasing cage were induced to fly. The air flow was controlled thanks to two mechanisms;
162 first, a power regulator (HQ-Power PS1502A, Velleman, Gavère, Belgium) connected to fans; and second, iris
163 dampers (10 cm Ø; CIR D100, France air, France) connecting the air vent hoses to the collecting boxes. The
164 openings of the air vent hoses on both the swarming and collecting boxes sides were covered with nets to prevent
165 mosquitoes from flying into the air vent hoses. The swarming boxes (odor-sources) were located side-by-side
166 outdoors and the olfactometer inside a room (Fig. 1). The air speed in the releasing cage was regulated at 18 ± 2
167 $\text{cm} \cdot \text{s}^{-1}$ using an anemometer (Model 425, Testo, Forbach, France) and the room temperature and relative
168 humidity were set at 27 ± 2 °C and $80 \pm 10\%$ RH thanks to an air conditioner and a humidifier (Defensor 505,
169 Condair, Croissy-Beaubourg, France), respectively.

170

171 **Bioassays**

172 **Test preparation:** In the morning, about 8 hours before the time of the test, 100 2- to 3-day-old virgin
173 females of *An. coluzzii* and *An. gambiae* were colored with two different colored powders (Luminous Powder
174 Kit, BioQuip, Rancho Dominguez, California, USA) according to the species. In addition, about 500 4- to 5-day-
175 old *An. coluzzii* or *An. gambiae* males were released into a swarming box to allow them to acclimate. Both males
176 and females were provided with a 10% glucose solution and were kept under insectary conditions until the time

177 of the test. To avoid biases during the test due to both humidity and odor as a result of manipulation or bacterial
178 proliferation, the empty swarming box (control box) was also provided with glucose.

179 **Test execution:** Bioassays were carried out at sunset, when males naturally swarm. About 30 min
180 before sunset time, the swarming boxes were moved outside, the cups containing the glucose pads were removed
181 and visual markers made of a 20 × 20 cm black cloth were placed either under or next to the boxes for *An.*
182 *coluzzii* or *An. gambiae*, respectively (see Poda et al. 2019). These markers allowed to trigger swarming behavior
183 and to stabilize swarms in the middle of the box. For the same purpose, we provided an artificial twilight horizon
184 made of a 40 W incandescent bulb (2,500 K) located on the floor between a white wall and a 50 cm high black
185 horizon (Facchinelli et al. 2015; Niang et al. 2019; Poda et al. 2019). Inside, about 10 min before introducing
186 females, the fans were switched on in order to purge the air vent hoses and the air flow was set at $18 \pm 2 \text{ cm.s}^{-1}$.
187 The glass tube openings on the releasing cage side were covered with nets to prevent female mosquitoes from
188 flying into the collecting boxes before the test began. The complete olfactometer, except the end of the releasing
189 cage, was covered with a white cloth to eliminate visual bias during the test and to provide a diffused light.
190 Sunset light was allowed to enter the olfactometer room until dark and an additional artificial twilight made of a
191 40 W incandescent bulb (2,500K) projected on the wall of the room on the side of the collecting boxes was also
192 provided. Thirty minutes before sunset, about 100 females of each species (N=200, 1 replicate) were released
193 simultaneously into the releasing cage of the olfactometer allowing them to acclimate.

194 When the males started to swarm (*i.e.* flying in loops at a constant location within the swarming boxes,
195 see Downes 1969; Poda et al. 2019), the nets covering the glass tube openings and preventing access of females
196 to the collecting boxes were removed thanks to threads that were attached to the nets. The females were allowed
197 to respond for 20 min (swarm duration), then the glass tube openings were covered again with nets and females
198 that reached the upwind collecting boxes were removed with a mouth aspirator, killed by freezing and identified
199 according to their coloration (species) under a binocular (LEICA S6E) and counted. The remaining mosquitoes
200 inside the releasing cage were also removed. After each test (1 per day), the olfactometer was cleaned with 95%
201 ethanol to remove odor contaminants. All materials were handled with gloves to avoid contamination with skin
202 compounds.

203 **Test combinations:** Four combinations of choice tests were performed: i) *An. coluzzii* male swarm vs.
204 empty box (hereafter, *An. coluzzii* test); ii) *An. gambiae* male swarm vs. empty box (hereafter, *An. gambiae* test),
205 iii) *An. coluzzii* vs. *An. gambiae* male swarms (hereafter, *An. coluzzii* vs. *An. coluzzii* test) and iv) empty vs.
206 empty box (hereafter, control test). We ran 12 replicates per combination and females were used only once.

207 Males were used twice in two consecutive days. In that case, a 10% glucose solution was introduced into the two
208 swarming boxes after the test and boxes were kept under insectarium conditions until the next day. The
209 swarming boxes were cleaned with ethanol between each batch of males. **To avoid biases, the matching of**
210 **species and colors was switched between each test, and we alternated the treatments (mosquitoes or control)**
211 **between the swarming boxes and the right and left arm of the olfactometer.** We assessed the instrumental and
212 **arm** bias through an empty vs. empty box choice test (control test). The olfactometer did not present any
213 symmetrical biases (Fig. S1).

214

215 **Collection of *swarm* volatile organic compounds (VOCs)**

216 Because compounds can be emitted at a given time and only produced in tiny quantities at individual scale, we
217 chose to work directly on swarms. Three different sampling methods were used; two methods were run in a
218 laboratory setup (Niang et al. 2019) and one in the field.

219

220 ***Dynamic headspace in the laboratory***

221 **Extraction setup:** The extraction of swarm VOCs was performed in a room specially designed to
222 **stimulate *Anopheles* swarming behavior thanks to a set of visual stimuli (see Niang et al. (2019) for details).** The
223 headspace volume consisted of a 50 × 50 × 50 cm transparent plexiglass box (**extraction box**). Such a volume
224 was a trade-off between a volume large enough to allow males to display a swarming behavior without too much
225 constraints and a volume small enough to be sucked up in a reasonable time. Entrance and exit flow rates were
226 maintained by two pumps and regulated by flowmeters. The entrance flow was higher than the exit flow
227 ensuring that the system was continuously purged to compensate for the inevitable leaks, and that no
228 contaminated outside air would enter the system. At the entrance side, **ambient** air was purified in a glass
229 charcoal filter and then humidified passing through deionized water. Exit flow passed through an odor trap to
230 adsorb VOCs. Three different odor traps were used for the chemical analyses and electrophysiological analyses.
231 Tenax-TA/Carbotrap sorbent stainless steel 6mm diameter tubes (Gerstell, Mülheim, Germany) and Porapak-Q
232 sorbent VCT glass tubes (ARS, Gainesville, Florida, USA) were used for chemical analyses. Porapak-Q sorbent
233 VCT tubes and home-made sorbent Micro-traps were used for electrophysiological analysis. Micro-traps were
234 constituted by ChromatoProbe® quartz microvials of Varian Inc. (length: 15 mm; inner diameter: 2 mm),
235 previously cut closed-end and filled with 3 mg of a 1:1 mix of Tenax-TA and Carbotrap® (60–80 and 20–40

236 mesh, respectively; Sigma Aldrich, Munich, Germany)(See Dormont et al. 2013b). Flow rates and extraction
237 durations were dependent on the odor trap (See Table 1).

238 **Extraction:** About 500 4- to 6-day-old males of *An. gambiae* or *An. coluzzii* were transferred into the
239 extraction box about 30 min before swarming time. Clean air was pushed into the box at a rate of 10 L.min⁻¹ to
240 purge the box from both human and mosquito container odors before connecting it to the odor trap. Five minutes
241 after males started to swarm (*i.e.* flying in loops at a constant location within the box), the air flow was adjusted
242 at the required rate and the odor trap connected. Forty minutes later, swarming behavior was stopped by turning
243 off the light and the VOC collection continued until the required collection time was reached (see Table 1). After
244 extraction, samples were stored at 4 °C until analysis. After males were removed, the box was cleaned with 95%
245 ethanol and then flushed with clean air for at least 5 hours to remove the odor contaminants left from the
246 previous extraction. A total of 30 swarm extracts were collected; 17 *An. coluzzii* swarms (5, 8 and 4 with
247 Tenax/Carbotrap tubes, Porapak-Q VCT tubes and Micro-traps, respectively) and 13 *An. gambiae* swarms (6 and
248 7 with Tenax/Carbotrap tubes, and Porapak-Q VCT tubes, respectively). Control consisted of a clean empty box.
249 Before they were sent and used in Burkina Faso, adsorbents were cleaned as follows. Tenax-TA/Carbotrap tubes
250 were heated at 250 °C for 30 min under a 30 mL.min⁻¹ flow then sealed. Porapak-Q VCT tubes were eluted with
251 3 mL of hexane then packed individually in a nalophan bag. Chromatoprobes were heated for 2 h at 270 °C by
252 100 chromatoprobes under a 100 mL.min⁻¹ nitrogen flow, then packed in a glass vial with a Teflon cap.

253
254 **VOC collection from natural swarms** - VOCs were collected from a natural swarm of *An. coluzzii* in the village
255 of Bama. Static sorptive extractions of volatiles from a non-enclosed swarm were performed with Twisters®
256 (100 µm PDMS stir bar; Gerstel, Mülheim, Germany; Bicchi et al. 2000; Tienpont et al. 2000). The swarm flew
257 about 2.5 to 3.5 m above the ground. Thus, we used a 1.5 m long glass stick (Glaswarenfabrik Karl Hecht GmbH
258 & Co, Germany) with a metallic push pin head glued at the tip and covered with nalophan to introduce the
259 magnetic twisters directly into the swarm. Control twisters were placed about 3 m away upwind from the swarm.
260 Extraction lasted for the whole swarm duration (about 20-25 min). We chose a swarm far from habitations and
261 livestock to limit odor pollutions, containing a large number of males (more than 1,000) and attractive for
262 females (observation of a large number of couples). After VOC collections, twisters were individually placed
263 into glass vials and stored at 4 °C until analysis. Five replicates were performed on the same swarm on different
264 days.

265

266 **Solvent extraction** - About 1,000 4- to 6-day-old males of *An. gambiae* or *An. coluzzii* were introduced into a 30
267 × 30 × 30 cm cage and kept under insectarium conditions. The cage was made of unpainted metallic frames
268 covered by white nets and the bottom was covered with a white paper. Thirty minutes before sunset time, the
269 cage was placed outside and mosquitoes were observed. A 5 × 5 cm black cloth was placed in or out of the cage
270 according to the species as described above to stimulate swarming behavior. About 5 min after mosquitoes
271 started to swarm, the entire cage was quickly placed into a freezer at -20° C for about 10 min. Then, mosquitoes
272 were transferred into a 20 ml glass tube and covered with a 1:1 mix of Hexane and Dichloromethane. The tube
273 was sealed and mosquitoes were kept in the solution for 24h at 27±2 °C. Then, the solution was filtered on glass
274 wool to remove mosquito scales and stored at 4° C until analysis. Three different controls were made: i) with 1-
275 to 2-day-old virgin males (young male control), ii) with 2- to 3-day-old virgin females (female control) and iii)
276 with solvent mix alone (blank control). A total of 8 swarm extracts were done; 4 of *An. coluzzii* and 4 of *An.*
277 *gambiae* and 1 replicate of each type of control per species.

278

279 **SPME extraction** – We replicated the experiment from the study by Mozūraitis et al. (2020) in which they
280 collected mosquito headspace with a SPME fiber in a 1L bottle. However, because most methods used to
281 introduce mosquitoes into a bottle may potentially result in entering pollutants as well, we added a
282 supplementary control. One way to easily introduce the mosquitoes is to blow them from a mouth aspirator into
283 the bottle. This classic method may bring breath volatiles within the recipient, which can contain the five
284 compounds described in Mozūraitis et al. (2020) (Poli et al. 2010; Calenic and Amann 2014; Filipiak et al. 2014;
285 Cainap et al. 2020). Consequently, we added a "breath control". In addition, we also replicated the experiment
286 with females to check for sex specificity. In the morning, the two glass bottles were cleaned with acetone and
287 dried under a clean air flow, then closed with nalophan. Three hours before swarming time (5pm), we collected
288 the headspace of empty bottles with polydimethylsiloxane/divinylbenzene-coated SPME fibers for 1h (step 1:
289 empty bottle). Then, we simulated the introduction of mosquitoes by blowing 3 times for 3 seconds with a mouth
290 aspirator into the bottles. These were then closed with Nalophan (6pm) and the head space of the breath of the
291 experimenter was collected for 1h with SPME fibers (step 2: breath control). We then introduced 60-70, 5- to 7-
292 day-old unmated males in each bottle with the mouth aspirator, closed the bottles with Nalophan (7pm) and let
293 the mosquitoes acclimate for 1h. During the acclimation period, we simulated a light transition from photophase
294 to scotophase to trigger crepuscular flight. Then, when males started to fly (8pm), SPME fibers were introduced
295 for 1h (step 3: "swarm" extract). Six different fibers were used and cleaned at 250 °C during 5 min in the injector

296 of a gas-chromatograph (HP 6890 Series PLUS, Agilent, Santa Clara, USA) just before use. Each fiber was
297 always used for the same type of extract and in the same bottle. Before the experiment, we made sure that the
298 fibers had similar sensitivity by exposing them to a mix of the five compounds. Only minor quantitative
299 differences were detected, and this was considered during subsequent statistics. Secondly, we performed the
300 same 3 steps but with 60-70, 3- to 4-day-old virgin females instead of males. And finally, as during the
301 introduction of mosquitoes, we blew a second time into the bottles and repeated the three steps of headspace
302 extraction but this time, instead of introducing mosquitoes in the 3rd step, we blew again into the bottles (breath
303 ×2) to check for potential accumulation of breath compounds. We kept track of the day of the experiment, the
304 bottles and fibers identity for statistical analyses.

305

306 **Chemical analyzes**

307 Samples obtained on Porapak-Q VCT tubes were eluted with 150 µl of Hexane and injected (1 µl) into a Gas
308 Chromatograph coupled with Mass Spectrometer (GC-MS). Those obtained on Tenax/Carbotrap tubes, twisters
309 or SPME fibers were directly thermo-desorbed into GC-MS. Solvent extracts were injected as is or concentrated
310 under a gentle stream of nitrogen. All the analyses were run at the “Platform for Chemical Analyses in Ecology”
311 (PACE), the technical facilities of the LabEx CeMEB (Centre Méditerranéen pour l’Environnement et la
312 Biodiversité, Montpellier, France).

313

314 *Liquid analysis (Porapak elutions and solvent extractions)* - Liquids were analysed with a GC-MS QP2010
315 Plus (Shimadzu, Kyoto, Japan). Mass spectra were recorded in electronic impact mode (EI) at 70eV over a m/z
316 mass range from 38 to 350. The temperature of the transfer line and the ion source were programmed to 250 °C
317 and 200 °C, respectively. The injections were made with an injector temperature of 250 °C, and a 1:4 split mode
318 ratio. Analyses were performed using a 30 m × 0.25 mm × 0.25 µm Optima 5-MS (Macherey-Nagel, Düren,
319 Germany) fused silica capillary column with a constant helium flow set close to 1 ml.min⁻¹. The oven
320 temperature was programmed as follows: 40 °C (held 5 min) to 250 °C at 6 °C.min⁻¹, and finally to 300 °C at 14
321 °C.min⁻¹ and held 2 min. GC-MS Solution software (Shimadzu, Kyoto, Japan) was used for data processing of
322 these analyses, with the NIST 2011 as spectrum database.

323

324 *Tenax/Carbotrap tubes and twister analysis* - These sorbents were analysed as described by Souto-Vilarós et al.
325 (2018). Samples were analysed using a gas chromatograph (GC, Trace™ 1310, Thermo Scientific™ Milan,

326 Italy) coupled with a mass spectrometer (MS, ISQ™ QD Single Quadrupole, Thermo Scientific™ Milan, Italy).
327 The column used was an Optima 5-MS, the same as for liquid analysis. Absorbent traps were handled with a
328 Multi Purpose Sampler (Gerstell, Mülheim, Germany) and desorbed with a double stage desorption system,
329 composed of a Thermal Desorption Unit (TDU) and a Cold Injection System (CIS) (Gerstell, Mülheim,
330 Germany). The sorbents were desorbed in splitless mode at 250 °C on a trap cooled at -80 °C by liquid nitrogen.
331 Then, the trap was heated to 250 °C with a 1:4 split ratio to inject the compounds in the column. The carrier gas
332 used was helium at 1 ml.min⁻¹. Oven temperature program was as follows: held at 40 °C for 3 min., then
333 increased to 220 °C at 5 °C.min⁻¹ and finally to 250 °C at 10 °C.min⁻¹, and held for 2 min. Mass spectra were
334 recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38 to 350. The temperature of the
335 transfer line and the ion source were programmed to 250 °C and 200 °C, respectively. Xcalibur™ software
336 (Thermo Scientific™, Milan, Italy) was used for data processing and the NIST 2011 as spectrum database.

337

338 **SPME fiber analysis** – We analyzed SPME fibers as described in Mozūraitis et al. (2020) with a GC-MS
339 QP2010-SE (Shimadzu, Kyoto, Japan). Volatiles were desorbed into the injector at 225 °C for 1 min in split
340 mode (1:4 ratio). Analyses were performed using a 30 m × 0.25 mm × 0.25 μm J&W DB-Wax silica capillary
341 column (Agilent, Santa Clara, USA) with a constant helium flow at 1 ml.min⁻¹. The oven temperature was
342 programmed as follows: 40 °C (held 1 min.) to 150 °C at 5 °C.min⁻¹, and finally to 220 °C at 20 °C.min⁻¹ and
343 held 9 min. Mass spectra were recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38
344 to 350. The temperature of the transfer line and the ion source were 250 °C and 200 °C, respectively. GC-MS
345 Solution software (Shimadzu, Kyoto, Japan) was used for data processing of these analyses.

346

347 **Electrophysiological analysis**

348 Electrophysiological activity of female olfactory detection, on the male swarm extracts, was tested with an
349 Electro-Antennographic Detection system, coupled with Gas Chromatography (GC-EAD) at the PACE. Thirty
350 minutes before the mosquito scotophase, both 4- to 6-day-old males and 2- to 3-day-old virgin females were
351 placed separately in cardboard cups (Ø = 75 mm, h = 100 mm), provided with a 5% honey solution and
352 transferred from the insectarium to the electrophysiology laboratory at PACE. [At mating time, mosquitoes were](#)
353 [placed in the dark. Males were checked for erected antennae as a proxy to ensure that change in the light-dark](#)
354 [cycle, transport and electrophysiology laboratory conditions did not alter mosquito motivation for mating.](#) Two

355 minutes after male antennae were erected, a female was removed from the cardboard for the antennal
356 preparation.

357 The female's head was gently excised with a pair of surgical scissors and mounted between two glass
358 capillary tubes of 76 mm in length and 1.12 mm in diameter (World Precision Instrument, Sarasota, USA),
359 pulled and cut using a vertical micropipette-puller (P-30 model, World Precision Instruments, Hertfordshire,
360 United Kingdom) filled with insect Ringer solution (6.0 g.l⁻¹ NaCl, 0.4 g.l⁻¹ KCl, 0.27 g.l⁻¹ CaCl₂ and 3.20 g.l⁻¹
361 NaC₃H₅O₃) and connected to silver wires. The indifferent electrode was inserted into the back (the foramen) of
362 the isolated head and the intact tip of one antenna was inserted into the recording electrode which was connected
363 to an Electro-Antennography Detector setup (EAD, Syntech IDAC-2, Kirchzarten, Germany). One end of the
364 glass electrodes was pulled to a fine point which was then broken to ensure a close fit with the female's head and
365 antenna.

366 The above EAD setup was linked to a Gas Chromatograph-Flame Ionization Detector (GC-FID, CP-
367 3800, Varian, Palo Alto, USA) equipped with an Optima 5-MS capillary column, the same as for the GC-MS
368 analyses. Liquid injections after Porapak solvent elution were made in a 1079 PTV injector (Programmed
369 Temperature Vaporizator) at 250 °C with a 1:4 split ratio. Microtraps were desorbed in the same injector using a
370 ChromatoProbe sample introduction device (Varian, Palo Alto, USA) with the same split ratio and temperature
371 programmed as follows: increased from 40 °C to 250 °C at 200 °C.min⁻¹ and held for 5 min. Oven temperature
372 was held at 50 °C for 0.40 min, then increased to 180 °C at 10 °C.min⁻¹, and finally to 220 °C at 12 °C.min⁻¹, for
373 a run duration less than 17 min. The carrier gas used was helium at 1 ml.min⁻¹. The effluent was split equally
374 into two deactivated fused silica capillary columns (100 cm x 0.25 mm), one leading to the FID (270 °C) and the
375 other into a heated EAD port (200 °C) (transfer line: Syntech, Kirchzarten, Germany) and led to antenna. The
376 GC effluent for antenna was mixed with charcoal-filtered, humidified airflow (300 ml.min⁻¹). StarWorkstation
377 6.41 software (Varian, Palo Alto, USA) was used for data processing.

378 Different types of male swarm extracts were tested on the antennae of several conspecific females.
379 Antennae of 19 *An. coluzzii* females were tested with solvent extracts, Porapak-Q VCT tube extracts and Micro-
380 trap extracts (8, 9 and 2 females, respectively) and 20 *An. gambiae* females were tested with solvent and
381 Porapak-Q VCT tube extracts (10 females each). Before GC-EAD analysis of the VOC extracts, we checked the
382 living antenna for good activity using a negative (solvent) and a positive (mix of many VOCs) stimulus
383 solutions. We proceeded as follows: 5µl of each solution was applied to a 1 cm × 2 cm filter paper contained in a
384 Pasteur pipette. The pipette was then placed in an air pump system and the volatiles were directly sent to the

385 antennal preparation with a pulse duration of 0.5 s and a flow rate of 890 ml.min⁻¹ regulated by a CS-55 Stimulus
386 Controller (Syntech, Hilversum, Netherlands).

387

388 **Identification of specific and active VOCs**

389 For all the chemical analysis, we compared the chromatograms of swarming mosquito extracts with
390 those of their corresponding controls to search for any qualitative or quantitative differences. When such a
391 difference was pinpointed, we matched the mass spectra of the compounds of interest with the NIST 2011 MS
392 library. A solution of n-alkanes (Alkanes standard solution, 04070, Sigma Aldrich, Darmstadt, Germany) was
393 also injected to calculate the linear retention index (LRI) of these compounds of interest (Zellner et al. 2008) and
394 their LRIs were compared with those reported in the literature.

395 In the electrophysiological analysis, we studied the chromatograms in a range of LRI between 800 and
396 1700, including compounds present at very low and trace amounts. When a depolarization was observed in the
397 electrophysiological analysis, a section of 20 points of LRI was studied around the compound's LRI on the GC-
398 MS trace. Comparison of the samples with their respective controls collected on the same day, allowed to
399 subtract potential contaminants from the samples. Non-natural compounds, such as industrial chemicals and/or
400 compounds not naturally produced by living organisms (Charpentier et al. 2012), were considered as
401 contaminants.

402 We also particularly searched for the presence of acetoin, sulcatone, octanal, nonanal and decanal
403 reported by Mozūraitis et al. (2020). With that aim, we defined a LRI window of 20 points on both sides of the
404 LRI of these compounds on the bases of injected standards (for SPME fibers; acetoin, sulcatone, octanal,
405 nonanal and decanal, ≥98%, 99%, 99%, ≥98% and ≥98% GC, respectively, Sigma-Aldrich, Darmstadt,
406 Germany) or literature (for other extraction methods). Then, we searched for their specific ions at those LRI as
407 follows: (LRI/ion) 711/88, 981/108, 998/84, 1100/98, and 1201/112 for acetoin, sulcatone, octanal, nonanal and
408 decanal, respectively. Then, when a peak corresponding to both LRI and specific ion was found, we compared its
409 spectra with the NIST 2011 MS library before quantification with integration units.

410

411 **Statistical analysis**

412 Both the activation rate and choice were analyzed using binomial Generalized Linear-Mixed Models
413 (GLMM, "glmer" function in "lme4" package). Tested combinations (four levels: *An. coluzzii*, *An. gambiae*, *An.*
414 *coluzzii* vs. *An. gambiae* and control tests), female species (two levels: *An. coluzzii* and *An. gambiae*) and their

415 interaction were considered as fixed effects, and replicates as random effects. All analyses were performed using
416 R (version 3.4.0).

417 We analyzed separately the quantities using integration units of acetoin, sulcatone, octanal, nonanal and
418 decanal. When pairing between the controls and the mosquito extracts was possible (with SPME and twister) and
419 to account for repeated measurements, we used Gaussian GLMMs. In those cases, the extract (3 levels for SPME
420 fibers: empty bottles, breath and flying mosquitoes (either male or females or breath $\times 2$) and 2 levels for
421 twisters: control and swarm) was considered as fixed effect. The replicates (i.e. the day for twisters and the
422 SPME) and fiber nested within the bottle for SPME fibers were considered random effects. Considering the
423 SPME fibers as a random effect made it possible to account for their minor differences of sensitivity in the
424 model. When pairing between the controls and the mosquito extracts was not possible (Tenax-TA/Carbotrap and
425 Porapak-Q VCT tubes), Gaussian GLMs were used. The extract (empty box, *An. coluzzii* swarm or *An. gambiae*
426 swarm) was considered as fixed effect. In some occasions, some compounds were not found in the extracts or
427 only in two samples. Consequently, it was not possible to perform statistical analyses and we considered the
428 extracts as not significantly different. No statistical analysis was performed with liquid extracts due to a small
429 number of replicates. For model selection, we used the stepwise removal of terms, followed by likelihood ratio
430 tests. Term removals that significantly reduced explanatory power ($P < 0.05$) were retained in the minimal
431 adequate model. All percentages are provided with their 95% confident interval.

432

433

RESULTS

434 Behavioral assays

435 Over 12 replicates of c.a. 200 females for each of the four combinations, a total of 8,918 females were
436 tested (4,423 *An. coluzzii* and 4,495 *An. gambiae*) among which 3,591 flew upwind into the collecting boxes
437 (activation rate: $40.2 \pm 1\%$). There was a significant interaction between female species and the four combinations
438 of swarm boxes tested ($\chi^2_3 = 22.19$, $P < 0.001$). Species subset analyses showed that the different test
439 combinations had no effect on the *An. coluzzii* activation rate ($\chi^2_3 = 5.72$, $P = 0.12$; Fig. 2). However, in *An.*
440 *gambiae*, there was a significant effect of the test combinations on the female activation rate ($\chi^2_2 = 14.94$, $P =$
441 0.001 ; Fig. 2) with a higher activation rate when females were exposed simultaneously to both *An. coluzzii* and
442 *An. gambiae* male swarms than when exposed to an *An. coluzzii* swarm alone ($z = 4.01$, $P < 0.001$) or to the
443 control (two empty boxes: $z = 2.57$, $P = 0.04$). Nonetheless, no difference was found when females were exposed
444 to an *An. gambiae* male swarm alone ($z = -1.49$, $P = 0.44$). Neither the *An. coluzzii* swarms, nor the *An. gambiae*

445 swarms were attractive for activated females (female species: $\chi_1^2 = 2.45$, $P = 0.11$; choice test combinations: χ_2^2
446 $= 1.15$, $P = 0.56$ and female species \times choice test combination interaction: $\chi_2^2 = 0.34$, $P = 0.84$; Fig. 3).

447

448 **Chemical analyses**

449 **No qualitative or quantitative differences were detected between samples of swarming males and their**
450 **relative controls for both species obtained from the Porapak-Q, Twisters® and the liquid extracts (Figs S2 & S3).**
451 **Minor quantitative differences were detected with the Tenax/Carbotrap tubes. However, mass spectra showed**
452 **that these peaks were pollutants, such as silicones, BTX (mono-aromatic compounds like benzene, toluene and**
453 **xylene), alkanes which were all emitted by the box containing the mosquitoes. Quantitative differences including**
454 **linear organic acids were also detected, but they were not reproducible.**

455 During the experiment replicating the protocol by Mozūraitis et al. (2020), both males and females
456 started to fly at the expected time (8pm) but due to the small volume it was not possible to determine if the group
457 was a swarm. Analyses of SPME fibers in search for acetoin, sulcatone, octanal, nonanal and decanal showed
458 that acetoin was absent from almost all the male samples (Fig. 4A), but was present on three occurrences in
459 female extracts (over six) (Fig. 4B). Sulcatone, octanal, nonanal and decanal were found in almost all types of
460 extracts including controls. However, we did not find significant increased quantities of the five compounds in
461 "swarming" mosquito extracts compared to the breath controls (Fig. 4). The control to test the effect of blowing
462 twice in the bottle showed that the headspaces of "breath" and "breath $\times 2$ " were very similar (Fig. S4),
463 demonstrating an absence of accumulation of breath odor between the breath control and the introduction of
464 mosquitoes.

465 A posteriori, we also quantified acetoin, sulcatone, octanal, nonanal and decanal in our previous GC-
466 MS analyses in search for similar variations as described in Mozūraitis et al. (2020). We did not find acetoin in
467 our chromatograms. Indeed, as we used a low polarity Optima 5-MS column, acetoin (if present) was eluted
468 before the LRI 800. In the samples obtained from natural swarms of *An. coluzzii* with the twister method,
469 sulcatone and octanal were absent. The quantities of both nonanal and decanal were not significantly higher in
470 the swarm than in the control (Fig. 5A). In laboratory swarm samples, sulcatone, octanal, nonanal and decanal
471 were found inconsistently. Nevertheless, swarms did not show higher quantities of these compounds compared
472 to the empty box (Fig. 5B) excepted in samples obtained with the Porapak tubes (Fig. 5C). However, this
473 difference was not statistically significant. Whatever, the method used, we can note a high quantitative
474 variability between the samples. This was true for both the control and the mosquito samples.

475

476 **Electrophysiology analyses**

477 The male swarm VOCs induced a few weak and poorly reproducible depolarizations on the antennae of
478 conspecific females. In *An. coluzzii*, 3 females out of 9 weakly responded to a single Porapak-Q extract at the
479 linear retention index (LRI) 1371 (retention time (RT) 12.85 min), and 2 females out of 8 responded to solvent
480 extracts at LRI 921, 978 and 1179 (RT 6.40, 7.24 and 10.23 min, respectively, Fig. 6). However, these responses
481 did not correspond to noticeable peaks in the GC detector of the GC-EAD apparatus. Successive GC-MS
482 analyses of the same porapak-Q and liquid extracts did not allow to detect and identify any specific peak in
483 relation to antennae responses (Figs S2b & S3b). **The peaks found at LRIs that elicited a response in *An. coluzzii***
484 **females were identified as environmental pollutants and were present in both swarming mosquitoes and control**
485 **samples.** No reproducible response was recorded from *An. coluzzii* females tested with Micro-traps extracts. In
486 *An. gambiae*, no reproducible antennae response was recorded from the 10 females tested with either Porapak-Q
487 or liquid extracts.

488

489

DISCUSSION

490 For the past decades, several authors have suggested the existence of long-range sex pheromones in the attraction
491 of *An. gambiae s.l.* females to male swarms (Charlwood et al. 2002; Tripet et al. 2004; Diabaté et al. 2011; Poda
492 et al. 2019) but to date, no reliable experimental evidence **is** available. Here, we used behavioral, physiological
493 and detailed chemical approaches to attempt to highlight the existence of such compounds. However, despite a
494 large set of methodologies and working hypotheses to maximize the chances of highlighting the presence of
495 pheromones (*i.e.* timing, physiological state, natural behavior, recently colonized mosquitoes), our study failed to
496 provide evidence of the presence of long-range male sex pheromones in both *An. coluzzii* and *An. gambiae*. We
497 also replicated an experimental setup that recently showed an increase in quantity of five compounds in
498 "swarming" males compared to controls. However, unlike in Mozūraitis et al. (2020), we did not find any
499 significant difference with our controls.

500 **During** our behavioral tests in the olfactometer, **females were not attracted to air currents passed over**
501 **male swarms regardless of the species.** During the experiment, the males released in the large plexiglass boxes
502 and exposed to natural light started to fly randomly at sunset. Then, about 150-200 males out of the 500 released
503 (<50%) gathered in the upper half of the box flying in loops without touching the box walls. They also reacted to
504 the movements of the visual marker, meaning they were indeed swarms. The other males flew randomly in the

505 box, bouncing against the walls. Females showed a good activation rate but they did not show a particular choice
506 for any arm, containing a swarm or not. Nevertheless, *An. gambiae* females showed an intriguing higher
507 activation rate when exposed simultaneously to both a conspecific and a heterospecific swarm compared to a
508 single *An. coluzzii* swarm. This was, however, probably without biological significance as no difference was
509 detected in the test providing *An. gambiae* females with a choice between the control and *An. gambiae* swarm,
510 vs. a test offering two controls.

511 Our behavioral result was consistent with our physiological study. Indeed, despite the fact that we
512 checked for both behavioral receptivity of mosquitoes and for receptivity of mounted antennae, no consistent
513 antennal depolarization was observed in females across assays when exposed to swarm extracts. [This suggests a](#)
514 [lack of response](#) by females to any volatile chemicals present in our swarm extracts.

515 Chemical analyses were also negative and no compounds specific for male swarms could be detected
516 whatever the method used. Moreover, unlike in Mozūraitis et al. (2020), the quantities of acetoin, sulcatone,
517 octanal, nonanal and decanal were found inconsistently across swarming mosquito samples and not in larger
518 quantities compared to controls, making it difficult to support the assumption according to which they could
519 actually be emitted by males. These divergent results could be explained by the fact that, unlike in the
520 experiments of Mozūraitis et al. (2020), we used a control which considered the potential introduction of
521 pollutions at the same time as mosquitoes. This showed that breath was responsible for most of the variability.
522 The most convincing result showing that sulcatone, octanal, nonanal and decanal are probably not pheromones
523 emitted by males is the one obtained with the twister method. Indeed, twisters are coated with the same
524 adsorbent phases as SPME fibers used both in this study and in the one by Mozūraitis et al. (2020). According to
525 the manufacturer, twisters are up to a thousand times more sensitive than SPME fibers partly due to a larger
526 sorbent volume. Moreover, we exposed the twisters directly in a natural swarm formed by more than 1 000
527 males (probably up to 6 000 males according to the estimation of trained technical staffs). Despite this
528 sensitivity, the number of mosquitoes and the natural biological context, sulcatone and octanal were not found
529 and both nonanal and decanal had similar quantities in the swarm and outside the swarm (control twister placed
530 3 m upwind from the swarm). In addition, the high quantitative variability found in the laboratory and the fact
531 that these compounds are frequently found in controls suggest that they could be laboratory and/or human
532 pollutions that are difficult to control for.

533 [Our results contrast with those of Mozūraitis et al. \(2020\) probably for several reasons. First, in their](#)
534 [laboratory experiments, they reported a simple flight activity instead of a swarming activity in which males](#)

535 should fly in regular loops with erected antennae (Downes 1969; Poda et al. 2019). This can be explained by the
536 absence of adequate visual stimuli such as ground markers which are mandatory to trigger swarming behavior
537 (Charlwood and Jones 1980; Marchand 1984; Gibson 1985; Facchinelli et al. 2015; Niang et al. 2019; Poda et al.
538 2019). Second, this flight activity was extraordinary long for a swarming flight, exceeding 200 min in the
539 presence of the five compounds. Indeed, swarming behavior is known to last for only 20-30 min in nature
540 (Charlwood and Jones 1980; Marchand 1984; Sawadogo et al. 2013, 2014; Bimbilé Somda et al. 2018) and only
541 up to 60 min in artificial conditions (Charlwood and Jones 1980; Poda et al. 2019). In the latter, the number of
542 males in swarms decreased over time while the others flew randomly, bouncing on the flight arena walls (Poda et
543 al. 2019). Indeed, swarming is an activity with a high energy demand, consuming 50% of sugar and glycogen
544 reserves in 25 min (Maïga et al. 2012, 2014). Mosquitoes probably switched to a more random flight to try to
545 escape and search for a sugar meal to refuel their reserves. This behavior can be stimulated by acetoin, sulcatone,
546 octanal, nonanal and decanal, as they are frequently emitted from nectar sources and fermented sugar (Goodrich
547 et al. 2006; Schiestl 2010; Dekel et al. 2019). Finally, they also reported that these five compounds increased
548 insemination rates in five different species in semi-field cages. However, instead of stopping the experiment after the
549 swarming time, they collected the females in the morning. As they left both a dark box in the arena as resting site and a
550 sugar source, it is likely that males and females also mated in or around these two resources during the night. Indeed, it
551 was shown that mating can occur indoors, outside of swarming times and locations (Dao et al. 2008). The boxes and
552 sugar sources with these five compounds were probably more attractive than the control ones gathering more
553 mosquitoes and thus inducing more inseminations.

554 Independently, our methods can also have weaknesses. **Even if the number of swarming males was**
555 **biologically relevant compared to natural swarms, a higher number of males could be needed in experimental**
556 **setups, or** the sensitivity thresholds of our chemical and electrophysiological apparatus are below the ability of
557 insects to detect low amounts of compounds. However, altogether and keeping in mind that an absence of
558 evidence is not an evidence of absence, our results support the absence of long-range sex pheromones emitted by
559 male swarms. However, further investigations are needed and complementary methodologies such as electro-
560 physiology on palps (Lu et al. 2007; Iatrou and Biessmann 2008; Pitts et al. 2011; Guidobaldi et al. 2014),
561 molecular analyses of olfactory protein expression on adequate physiological status, or real-time chemical
562 analyses of swarm volatiles with a sensitive apparatus such as a proton-transfer-reaction mass spectrometry
563 could be options.

564 The question of how *Anopheles* females seek male swarms is still open. The lack of post-mating barrier
565 (Persiani et al. 1986; Diabaté et al. 2005, 2007; Hahn et al. 2012; Pombi et al. 2017) and the low rate of hybrids
566 in this geographical area (della Torre et al. 2001, 2005; Tripet et al. 2001; Diabaté et al. 2006) necessarily
567 involve a reasonably distant process which prevents females from entering heterospecific swarms. As acoustic
568 cues have shown some limitations at long range (Feugère et al. 2021), and because long-range chemical cues are
569 still disputable, visual cues such as ground markers (Diabaté et al. 2009; Sawadogo et al. 2014; Poda et al. 2019)
570 could be good candidates, but their effective range and level of specificity is poorly known.

571 From an evolutionary and ecological point of view, **our findings are** of utmost importance. **They**
572 **suggest** that the ecological barriers that allow these two species to mate assortatively are probably not of
573 chemical origin. Regarding control strategies, thanks to our findings, research programs working on the sterile
574 insect technique (SIT) or genetically-modified mosquitoes (GMM) will avoid focusing on the eventuality that
575 induced modifications may alter long-range sex pheromones making their mosquitoes less competitive than wild
576 individuals.

577

578 **Declarations**

579 **Funding**

580 This work was funded by a grant from the Agence Nationale de la Recherche (ANR-15-CE35-0001-01) awarded
581 to O.R. S.B.P. received financial support through a doctoral fellowship from the Institut de Recherche pour le
582 Développement (IRD).

583

584 **Competing interests**

585 We have no competing interests.

586

587 **Availability of data and material**

588 The raw datasets are available from the corresponding author upon reasonable request.

589

590 **Code availability**

591 Not applicable.

592

593 **Authors' contributions**

594 O.R. conceived the study. O.R., S.B.P., B.B., B.L. and L.D. designed the chemical and electrophysiological
595 experiments. O.R. and S.B.P. performed chemical extractions and SBP and BB performed the chemical analysis.
596 S.B.P. and B.L. performed the electrophysiological experiments. O.R. and S.B.P. designed olfactometric
597 experiments and S.B.P. performed data collection. S.B.P. and O.R. performed statistical analyses. S.B.P., O.R.,
598 B.B., B.L. drafted the manuscript and L.D., O.G., A.D, and R.K.D critically revised the manuscript. All authors
599 revised the manuscript, gave final approval for publication and are accountable for the work performed therein.

600

601 **Acknowledgements**

602 We thank David Sanou for his assistance in the field, Stephane Somda, Sofan Somé, Bethsabée Scheid, Marie
603 Rossignol and Carole Ginibre for their help in mosquito rearing and Heidi Lançon for proofreading the paper.

604

605 **REFERENCES**

606 Alexander RD, Marshall DC, Cooley JR (1997) Evolutionary perspectives on insect mating. In: Choe JC, Crespi

607 BJ (eds) The evolution of mating systems in insects and Arachnids. Cambridge University Press, pp 4–31

608 Andersson J, Vongvanich N, Wiklund C, et al (2007) Male sex pheromone release and female mate choice in a
609 butterfly. *J Exp Biol* 210:964–970. <https://doi.org/10.1242/jeb.02726>

610 Bicchi C, Cordero C, Iori C, et al (2000) Headspace Sorptive Extraction (HSSE) in the headspace analysis of
611 aromatic and medicinal plants. *J High Resolut Chromatogr* 23:539–546. [https://doi.org/10.1002/1521-](https://doi.org/10.1002/1521-4168(20000901)23:9<539::AID-JHRC539>3.0.CO;2-3)

612 [4168\(20000901\)23:9<539::AID-JHRC539>3.0.CO;2-3](https://doi.org/10.1002/1521-4168(20000901)23:9<539::AID-JHRC539>3.0.CO;2-3)

613 Bimbilé Somda NS, Poda BS, Sawadogo PS, et al (2018) Ecology of reproduction of *Anopheles arabiensis* in an
614 urban area of Bobo-Dioulasso, Burkina Faso (West Africa): monthly swarming and mating frequency and
615 their relation to environmental factors. *PLoS One* 13:e0205966.

616 <https://doi.org/10.1371/journal.pone.0205966>

617 Bjostad LB, Gaston LK, Shorey HH (1980) Temporal pattern of sex pheromone release by female *Trichoplusia*
618 *ni*. *J Insect Physiol* 26:493–498. [https://doi.org/10.1016/0022-1910\(80\)90122-5](https://doi.org/10.1016/0022-1910(80)90122-5)

619 Bray DP, Alves GB, Dorval ME, et al (2010) Synthetic sex pheromone attracts the leishmaniasis vector
620 *Lutzomyia longipalpis* to experimental chicken sheds treated with insecticide. *Parasit Vectors* 3:1–11.

621 <https://doi.org/10.1186/1756-3305-3-16>

622 Cainap C, Pop LA, Balacescu O, Cainap SS (2020) Early diagnosis and screening in lung cancer. *Am J Cancer*

623 *Res* 10:1993–2009

624 Calenic B, Amann A (2014) Detection of volatile malodorous compounds in breath: current analytical
625 techniques and implications in human disease. *Bioanalysis* 6:357–376. <https://doi.org/10.4155/bio.13.306>

626 Charlwood JD, Jones MDR (1980) Mating in the mosquito, *Anopheles gambiae* s.l. II. Swarming behaviour.
627 *Physiol Entomol* 5:315–320. <https://doi.org/10.1111/j.1365-3032.1980.tb00241.x>

628 Charlwood JD, Pinto J, Sousa CA, et al (2002) The swarming and mating behaviour of *Anopheles gambiae* s.s.
629 (Diptera: Culicidae) from São Tomé Island. *J Vector Ecol* 27:178–183

630 Charpentier MJE, Barthes N, Proffit M, et al (2012) Critical thinking in the chemical ecology of mammalian
631 communication: roadmap for future studies. *Funct Ecol* 26:769–774. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2435.2012.01998.x)
632 [2435.2012.01998.x](https://doi.org/10.1111/j.1365-2435.2012.01998.x)

633 Clements AN (1999) The biology of mosquitoes. Volume 2: sensory, reception and behaviour. CABI Publishing,
634 Wallingford, UK

635 Clements AN (1992) The biology of mosquitoes: development, nutrition and reproduction. Chapman & Hall,
636 London

637 Costantini C, Ayala D, Guelbeogo MW, et al (2009) Living at the edge: biogeographic patterns of habitat
638 segregation conform to speciation by niche expansion in *Anopheles gambiae*. *BMC Ecol* 9:16.
639 <https://doi.org/10.1186/1472-6785-9-16>

640 Dao A, Adamou A, Yaro AS, et al (2008) Assessment of alternative mating strategies in *Anopheles gambiae*:
641 does mating occur indoors? *J Med Entomol* 45:643–52

642 Dekel A, Yakir E, Bohbot JD (2019) The sulcatone receptor of the strict nectar-feeding mosquito *Toxorhynchites*
643 *amboinensis*. *Insect Biochem Mol Biol* 111:103174. <https://doi.org/10.1016/j.ibmb.2019.05.009>

644 della Torre A, Fanello C, Akogbeto M, et al (2001) Molecular evidence of incipient speciation within *Anopheles*
645 *gambiae* s.s. in West Africa. *Insect Mol Biol* 10:9–18. <https://doi.org/10.1046/j.1365-2583.2001.00235.x>

646 della Torre A, Tu Z, Petrarca V (2005) On the distribution and genetic differentiation of *Anopheles gambiae* s.s.
647 molecular forms. *Insect Biochem Mol Biol* 35:755–769. <https://doi.org/10.1016/j.ibmb.2005.02.006>

648 Diabaté A, Baldet T, Brengues C, et al (2003) Natural swarming behaviour of the molecular M form of
649 *Anopheles gambiae*. *Trans R Soc Trop Med Hyg* 97:713–716.
650 <https://doi.org/10.1371/journal.pone.0004549>

651 Diabaté A, Brengues C, Baldet T, et al (2004) The spread of the Leu-Phe kdr mutation through *Anopheles*
652 *gambiae* complex in Burkina Faso: genetic introgression and de novo phenomena. *Trop Med Int Heal*
653 9:1267–1273. <https://doi.org/10.1111/j.1365-3156.2004.01336.x>

654 Diabaté A, Dabiré KR, Kim HE, et al (2005) Larval development of the molecular forms of *Anopheles gambiae*
655 (Diptera: Culicidae) in different habitats: a transplantation experiment. *J Med Entomol* 42:548–553

656 Diabaté A, Dabiré KR, Millogo N, Lehmann T (2007) Evaluating the effect of postmating isolation between
657 molecular forms of *Anopheles gambiae* (Diptera: Culicidae). *J Med Entomol* 44:60–64

658 Diabaté A, Dabiré RK, Kengné P, et al (2006) Mixed swarms of the molecular M and S forms of *Anopheles*
659 *gambiae* (Diptera: Culicidae) in sympatric area from Burkina Faso. *J Med Entomol* 43:480–483.
660 [https://doi.org/10.1603/0022-2585\(2006\)43\[480:msotmm\]2.0.co;2](https://doi.org/10.1603/0022-2585(2006)43[480:msotmm]2.0.co;2)

661 Diabaté A, Dao A, Yaro AS, et al (2009) Spatial swarm segregation and reproductive isolation between the
662 molecular forms of *Anopheles gambiae*. *Proc R Soc B Biol Sci* 276:4215–4222.
663 <https://doi.org/10.1098/rspb.2009.1167>

664 Diabaté A, Yaro AS, Dao A, et al (2011) Spatial distribution and male mating success of *Anopheles gambiae*
665 swarms. *BMC Evol Biol* 11:184. <https://doi.org/10.1186/1471-2148-11-184>

666 Dormont L, Bessière J-M, Cohuet A (2013a) Human skin volatiles: a review. *J Chem Ecol* 39:569–578.
667 <https://doi.org/10.1007/s10886-013-0286-z>

668 Dormont L, Bessiere JM, McKey D, Cohuet A (2013b) New methods for field collection of human skin volatiles
669 and perspectives for their application in the chemical ecology of human-pathogen-vector interactions. *J*
670 *Exp Biol* 216:2783–2788. <https://doi.org/10.1242/jeb.085936>

671 Downes JA (1969) The swarming and mating flight of Diptera. *Annu Rev Entomol* 14:271–298.
672 <https://doi.org/10.1146/annurev.en.14.010169.001415>

673 Facchinelli L, Valerio L, Lees RS, et al (2015) Stimulating *Anopheles gambiae* swarms in the laboratory:
674 application for behavioural and fitness studies. *Malar J* 14:271. <https://doi.org/10.1186/s12936-015-0792-2>

675 Feugère L, Gibson G, Manoukis NC, Roux O (2021) Mosquito sound communication: are male swarms loud
676 enough to attract females? *J R Soc Interface* 18:20210121. <https://doi.org/10.1098/rsif.2021.0121>

677 Filipiak W, Filipiak A, Sponring A, et al (2014) Comparative analyses of volatile organic compounds (VOCs)
678 from patients, tumors and transformed cell lines for the validation of lung cancer-derived breath markers. *J*
679 *Breath Res* 8:027111. <https://doi.org/10.1088/1752-7155/8/2/027111>

680 Gibson G (1985) Swarming behaviour of the mosquito *Culex pipiens quinquefasciatus*: a quantitative analysis.
681 *Physiol Entomol* 10:283–296. <https://doi.org/10.1111/j.1365-3032.1985.tb00049.x>

682 Gibson G, Warren B, Russell IJ (2010) Humming in tune: sex and species recognition by mosquitoes on the
683 wing. *J Assoc Res Otolaryngol* 11:527–540. <https://doi.org/10.1007/s10162-010-0243-2>

684 Gibson NHE (1945) On the mating swarms of certain Chironomidae (Diptera). *Trans R Entomol Soc London*
685 95:263–294. <https://doi.org/10.1111/j.1365-2311.1945.tb00263.x>

686 Gomulski L (1988) Aspects of mosquito mating behaviour. University of London

687 Goodrich KR, Zjhra ML, Ley CA, Raguso RA (2006) When flowers smell fermented: the chemistry and
688 ontogeny of yeasty floral scent in Pawpaw (*Asimina triloba*: Annonaceae). *Int J Plant Sci* 167:33–46.
689 <https://doi.org/10.1086/498351>

690 Guidobaldi F, May-Concha IJ, Guerenstein PG (2014) Morphology and physiology of the olfactory system of
691 blood-feeding insects. *J. Physiol. Paris* 108:96–111

692 Hahn MW, White BJ, Muir CD, Besansky NJ (2012) No evidence for biased co-transmission of speciation
693 Islands in *Anopheles gambiae*. *Philos Trans R Soc B Biol Sci* 367:374–384.
694 <https://doi.org/10.1098/rstb.2011.0188>

695 Howell PI, Knols BGJJ (2009) Male mating biology. *Malar J* 8:S8. <https://doi.org/10.1186/1475-2875-8-S2-S8>

696 Iatrou K, Biessmann H (2008) Sex-biased expression of odorant receptors in antennae and palps of the African
697 malaria vector *Anopheles gambiae*. *Insect Biochem Mol Biol* 38:268–274.
698 <https://doi.org/10.1016/j.ibmb.2007.11.008>

699 Kelly DW, Dye C (1997) Pheromones, kairomones and the aggregation dynamics of the sandfly *Lutzomyia*
700 *longipalpis*. *Anim Behav* 53:721–731. <https://doi.org/10.1006/anbe.1996.0309>

701 Kostiainen R (1995) Volatile organic compounds in the indoor air of normal and sick houses. *Atmos Environ*
702 29:693–702

703 Kruza M, Lewis AC, Morrison CG, Carslaw N (2017) Impact of surface ozone interactions on indoor air
704 chemistry: a modeling study. *Indoor Air* 27:1001–1011. <https://doi.org/10.1111/ina.12381>

705 Lefèvre T, Gouagna LC, Dabire KR, et al (2009) Evolutionary lability of odour-mediated host preference by the
706 malaria vector *Anopheles gambiae*. *Trop Med Int Heal* 14:228–236. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-3156.2009.02206.x)
707 [3156.2009.02206.x](https://doi.org/10.1111/j.1365-3156.2009.02206.x)

708 Lefèvre T, Gouagna LC, Dabiré KR, et al (2010) Beer consumption increases human attractiveness to malaria
709 mosquitoes. *PLoS One* 5:e9546. <https://doi.org/10.1371/journal.pone.0009546>

710 Lehmann T, Diabaté A (2008) The molecular forms of *Anopheles gambiae*: a phenotypic perspective. *Infect*
711 *Genet Evol* 8:737–746. <https://doi.org/10.1016/j.meegid.2008.06.003>

712 Levi-Zada A, Fefer D, David M, et al (2014) Diel periodicity of pheromone release by females of *Planococcus*
713 *citri* and *Planococcus ficus* and the temporal flight activity of their conspecific males. *Naturwissenschaften*

714 101:671–678. <https://doi.org/10.1007/s00114-014-1206-y>

715 Lu T, Qiu YT, Wang G, et al (2007) Odor coding in the maxillary palp of the malaria vector mosquito *Anopheles*
716 *gambiae*. *Curr Biol* 17:1533–1544. <https://doi.org/10.1016/j.cub.2007.07.062>

717 Maïga H, Dabiré RK, Lehmann T, et al (2012) Variation in energy reserves and role of body size in the mating
718 system of *Anopheles gambiae*. *J Vector Ecol* 37:289–297

719 Maïga H, Niang A, Sawadogo SP, et al (2014) Role of nutritional reserves and body size in *Anopheles gambiae*
720 males mating success. *Acta Trop* 132S:S102–S107. <https://doi.org/10.1016/j.actatropica.2013.08.018>

721 Marchand RP (1984) Field observations on swarming and mating in *Anopheles gambiae* mosquitoes in Tanzania.
722 Netherlands J Zool 34:367–387

723 McBride CS, Baier F, Omondi AB, et al (2014) Evolution of mosquito preference for humans linked to an
724 odorant receptor. *Nature* 515:222–227. <https://doi.org/10.1038/nature13964>. Evolution

725 Merlin C, Lucas P, Rochat D, et al (2007) An antennal circadian clock and circadian rhythms in peripheral
726 pheromone reception in the moth *Spodoptera littoralis*. *J Biol Rhythms* 22:502–514.
727 <https://doi.org/10.1177/0748730407307737>

728 Mosqueira B, Soma DD, Namountougou M, et al (2015) Pilot study on the combination of an organophosphate-
729 based insecticide paint and pyrethroid-treated long lasting nets against pyrethroid resistant malaria vectors
730 in Burkina Faso. *Acta Trop* 148:162–169. <https://doi.org/10.1016/j.actatropica.2015.04.010>

731 Mozūraitis R, Hajkazemian M, Zawada JW, et al (2020) Male swarming aggregation pheromones increase
732 female attraction and mating success among multiple African malaria vector mosquito species. *Nat Ecol*
733 *Evol* 4:1395–1401. <https://doi.org/10.1038/s41559-020-1264-9>

734 Nguyen PL, Vantaux A, Hien DFS, et al (2017) No evidence for manipulation of *Anopheles gambiae*, *An.*
735 *coluzzii* and *An. arabiensis* host preference by *Plasmodium falciparum*. *Sci Rep* 7:9415.
736 <https://doi.org/10.1038/s41598-017-09821-x>

737 Niang A, Nignan C, Poda BS, et al (2019) Semi-field and indoor setups to study malaria mosquito swarming
738 behavior. *Parasit Vectors* 12:446. <https://doi.org/10.1186/s13071-019-3688-0>

739 Nyasembe VO, Tchouassi DP, Kirwa HK, et al (2014) Development and assessment of plant-based synthetic
740 odor baits for surveillance and control of malaria vectors. *PLoS Negl Trop Dis* 9:e89818.
741 <https://doi.org/10.1371/journal.pone.0089818>

742 Pandey SK, Kim K (2011) Human body-odor components and their determination. *Trends Anal Chem* 30:784–
743 796. <https://doi.org/10.1016/j.trac.2010.12.005>

744 Pennetier C, Warren B, Dabiré KR, et al (2010) “Singing on the wing” as a mechanism for species recognition in
745 the malarial mosquito *Anopheles gambiae*. *Curr Biol* 20:131–136.
746 <https://doi.org/10.1016/j.cub.2009.11.040>

747 Persiani A, Dideco MA, Petrangeli G (1986) Osservazioni di laboratorio su polimorfismi da inversione originati
748 da incroci tra popolazioni diverse di *Anopheles gambiae* s.s. *Ann Dell’Istituto Super Di Sanita* 22:221–224

749 Pitts RJ, Rinker DC, Jones PL, et al (2011) Transcriptome profiling of chemosensory appendages in the malaria
750 vector *Anopheles gambiae* reveals tissue- and sex-specific signatures of odor coding. *BMC Genomics*
751 12:271. <https://doi.org/10.1186/1471-2164-12-271>

752 Poda SB, Buatois B, Lapeyre B, et al (2021) No evidence for long-range male sex pheromones in two malaria
753 mosquitoes. *bioRxiv Prepr.* <https://doi.org/10.1101/2020.07.05.187542>

754 Poda SB, Nignan C, Gnankiné O, et al (2019) Sex aggregation and species segregation cues in swarming
755 mosquitoes: role of ground visual markers. *Parasites and Vectors* 12:589. [https://doi.org/10.1186/s13071-](https://doi.org/10.1186/s13071-019-3845-5)
756 [019-3845-5](https://doi.org/10.1186/s13071-019-3845-5)

757 Poda SB, Soma DD, Hien A, et al (2018) Targeted application of an organophosphate-based paint applied on
758 windows and doors against *Anopheles coluzzii* resistant to pyrethroids under real life conditions in Vallée
759 du Kou, Burkina Faso (West Africa). *Malar J* 17:136. <https://doi.org/10.1186/s12936-018-2273-x>

760 Poli D, Goldoni M, Corradi M, et al (2010) Determination of aldehydes in exhaled breath of patients with lung
761 cancer by means of on-fiber-derivatisation SPME-GC/MS. *J Chromatogr B* 878:2643–2651.
762 <https://doi.org/10.1016/j.jchromb.2010.01.022>

763 Pombi M, Kengne P, Gimonneau G, et al (2017) Dissecting functional components of reproductive isolation
764 among closely related sympatric species of the *Anopheles gambiae* complex. *Evol Appl* 00:1–19.
765 <https://doi.org/10.1111/eva.12517>

766 Robledo N, Arzuffi R (2012) Influence of host fruit and conspecifics on the release of the sex pheromone by
767 *Toxotrypana curvicauda* males (Diptera: Tephritidae). *Environ Entomol* 41:387–391.
768 <https://doi.org/10.1603/en11037>

769 Rund SSC, Bonar NA, Champion MM, et al (2013) Daily rhythms in antennal protein and olfactory sensitivity in
770 the malaria mosquito *Anopheles gambiae*. *Sci Rep* 3:2494. <https://doi.org/10.1038/srep02494>

771 Santolamazza F, Mancini E, Simard F, et al (2008) Insertion polymorphisms of SINE200 retrotransposons within
772 speciation islands of *Anopheles gambiae* molecular forms. *Malar J* 7:163. [https://doi.org/10.1186/1475-](https://doi.org/10.1186/1475-2875-7-163)
773 [2875-7-163](https://doi.org/10.1186/1475-2875-7-163)

774 Savolainen E (1978) Swarming in Ephemeroptera: the mechanism of swarming and the effects of illumination
775 and weather. *Ann Zool Fennici* 15:17–52

776 Sawadogo PS, Costantini C, Pennetier C, et al (2013) Differences in timing of mating swarms in sympatric
777 populations of *Anopheles coluzzii* and *Anopheles gambiae* s.s. (formerly *An. gambiae* M and S molecular
778 forms) in Burkina Faso, West Africa. *Parasit Vectors* 6:275. <https://doi.org/10.1186/1756-3305-6-275>

779 Sawadogo PS, Namountougou M, Toé KH, et al (2014) Swarming behaviour in natural populations of *Anopheles*
780 *gambiae* and *An. coluzzii*: review of 4 years survey in rural areas of sympatry, Burkina Faso (West Africa).
781 *Acta Trop* 130:24–34. <https://doi.org/10.1016/j.actatropica.2013.10.015>

782 Schiestl FP (2010) The evolution of floral scent and insect chemical communication. *Ecol Lett* 13:643–656.
783 <https://doi.org/10.1111/j.1461-0248.2010.01451.x>

784 Shelly TE, Whittier TS (1997) Lek behavior of insects. In: Choe JA, Crespi JB (eds) *The evolution of mating*
785 *systems in insects and Arachnids*. Cambridge University Press, New York, pp 294–309

786 Sivinski JM, Petersson E (1997) Mate choice and species isolation in swarming insects. In: Choe JA, Crespi JB
787 (eds) *The evolution of mating systems in insects and Arachnids*. Cambridge University Press, New York,
788 pp 294–309

789 Souto-Vilarós D, Proffit M, Buatois B, et al (2018) Pollination along an elevational gradient mediated both by
790 floral scent and pollinator compatibility in the fig and fig-wasp mutualism. *J Ecol* 106:2256–2273.
791 <https://doi.org/10.1111/1365-2745.12995>

792 Suh E, Choe D, Saveer AM, Zwiebel LJ (2016) Suboptimal larval habitats modulate oviposition of the malaria
793 vector mosquito *Anopheles coluzzii*. *PLoS One* 11:e0149800.
794 <https://doi.org/10.1371/journal.pone.0149800>

795 Tchouassi DP, Sang R, Sole CL, et al (2013) Common host-derived chemicals increase catches of disease-
796 transmitting mosquitoes and can improve early warning systems for rift valley fever virus. *PLoS Negl*
797 *Trop Dis* 7:e2007. <https://doi.org/10.1371/journal.pntd.0002007>

798 Tienpont B, David F, Bicchi C, Sandra P (2000) High capacity headspace sorptive extraction. *J Microcolumn*
799 *Sep* 12:577–584. [https://doi.org/10.1002/1520-667X\(2000\)12:11<577::AID-MCS30>3.0.CO;2-Q](https://doi.org/10.1002/1520-667X(2000)12:11<577::AID-MCS30>3.0.CO;2-Q)

800 Tripet F, Dolo G, Traoré S, Lanzaro GC (2004) The “wingbeat hypothesis” of reproductive isolation between
801 members of the *Anopheles gambiae* complex (Diptera: Culicidae) does not fly. *J Med Entomol* 41:375–
802 384. <https://doi.org/10.1603/0022-2585-41.3.375>

803 Tripet F, Touré YT, Taylor CE, et al (2001) DNA analysis of transferred sperm reveals significant levels of gene

804 flow between molecular forms of *Anopheles gambiae*. Mol Ecol 10:1725–1732.
805 <https://doi.org/10.1046/j.0962-1083.2001.01301.x>

806 Vantaux A, Hien DFS, Yameogo B, et al (2015) Host-seeking behaviors of mosquitoes experimentally infected
807 with sympatric field isolates of the human malaria parasite *Plasmodium falciparum*: no evidence for host
808 manipulation. Front Ecol Evol 3:86. <https://doi.org/10.3389/fevo.2015.00086>

809 Verhulst NO, Andriessen R, Groenhagen U, et al (2010) Differential attraction of malaria mosquitoes to volatile
810 blends produced by human skin bacteria. PLoS One 5:e15829.
811 <https://doi.org/10.1371/journal.pone.0015829>

812 Wang G, Diabaté A, Liu J, et al (2021) Clock genes and environmental cues coordinate *Anopheles* pheromone
813 synthesis, swarming, and mating. Science (80-) 371:411–415. <https://doi.org/10.1126/science.abd4359>

814 Wondwosen B, Birgersson G, Seyoum E, et al (2016) Rice volatiles lure gravid malaria mosquitoes, *Anopheles*
815 *arabiensis*. Sci Rep 6:37930. <https://doi.org/10.1038/srep37930>

816 Wondwosen B, Birgersson G, Tekie H, et al (2018) Sweet attraction: sugarcane pollen-associated volatiles
817 attract gravid *Anopheles arabiensis*. Malar J 17:90. <https://doi.org/10.1186/s12936-018-2245-1>

818 Zellner B d'Acampora, Bicchi C, Dugo P, et al (2008) Linear retention indices in gas chromatographic analysis:
819 a review. Flavour Fragr J 23:297–314. <https://doi.org/10.1002/ffj>

820

821 **Figure legends**

822 **Fig. 1** Schematic of the olfactometer set up. S: swarm; M: marker. Not at scale.

823

824 **Fig. 2** Mosquito activation rate, expressed as the proportion of females caught in both collecting boxes out of the
825 total number released females for each of the four tested combinations. Different letters indicate difference at
826 $P < 0.05$.

827

828 **Fig. 3** Mosquito choice, expressed as the proportion of females caught in one or the other collecting box out of
829 the total number of "activated" females for each test.

830

831 **Fig. 4** Quantities (integration units) of acetoin, sulcatone, octanal, nonanal and decanal contained in the
832 headspace of the empty bottles, the breath of the manipulator and flying males (A) and females (B) of *Anopheles*
833 *gambiae* Kisumu and collected with SPME fibers. The intensity values correspond to the counts related to the
834 abundance of the specific ions representative of each molecule formed in the mass spectrometer and correspond
835 to the amount of compound analyzed. The box plots indicate the median (wide horizontal bars), the 25th and
836 75th percentiles (squares), and the minimum and maximum values (whiskers). The black dots represent outliers
837 and grey dots raw data. NS=not-significant; * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$.

838

839 **Fig. 5** Quantities (integration units) of sulcatone, octanal, nonanal and decanal contained in a natural swarm
840 sampled with twisters (A); in the headspace of the empty box, swarming *Anopheles coluzzii* males and *An.*
841 *gambiae* males sampled with Tenax/Carbotrap tubes (B); or collected with Porapak tubes (C). The intensity
842 values correspond to the counts related to the abundance of the specific ions representative of each molecule
843 formed in the mass spectrometer and correspond to the amount of compound analyzed. The box plots indicate
844 the median (wide horizontal bars), the 25th and 75th percentiles (squares), and the minimum and maximum
845 values (whiskers). The black dots represent outliers and grey dots raw data. NS=not-significant; * = $P < 0.05$; **
846 = $P < 0.01$; *** = $P < 0.001$. NA: The analytical procedure for these analyses did not allow detection of acetoin.

847

848 **Fig. 6** Mean electrophysiological responses of female *An. coluzzii* antennae to male swarm solvent extracts,
849 recorded by coupled gas chromatography – electroantennography detection (GC-EAD). Dashed lines indicate

850 retention times (and related LRI) for which the most significant EAG responses were recorded. Only 2 females
851 in 8 tested produced this pattern.

852

853

854 **Table 1:** Air inlet and outlet rates, and VOC extraction duration according to the type of trap used.

	Air flow rate ($L \cdot min^{-1}$)		VOCs extraction duration (hour)
	Air inlet	Air outlet	
Tenax/Carbotrap tube	2.5	2.0	2.0
Porapak-Q VCT tube	1.5	1.0	2.5
Micro-trap	0.9	0.4	5.0

855

856

857

858 **Fig. S1** Test for olfactometer symmetry. Female choice, expressed as the proportion of female mosquitoes
859 caught in one or the other collecting box out of the total number retrieved from both collecting boxes for the test
860 with empty boxes (control vs. control combination).

861

862 **Fig. S2 A.** Chromatograms obtained with eluted Porapak extracts from headspaces of the control, an *An. coluzzii*
863 swarm and an *An. gambiae* swarm (linear retention index (LRI) 800 to 1700). **B.** Enlarged view of the retention
864 time (RT) interval containing the LRI at which *An. coluzzii* antennal responses were detected in GC-EAD. Doted
865 white lines correspond to the theoretical RT of Sulcatone, octanal, nonanal and decanal. No difference was
866 detected between the control and the swarm extracts. All the peaks correspond to emissions born from the
867 material.

868

869 **Fig. S3 A.** Chromatograms obtained with the desorption of twisters obtained with the static sorptive extraction
870 methods under natural conditions in the field (linear retention index (LRI) 800 to 1700). The twister was either
871 located within a non-enclosed swarm of *An. coluzzii* or 3 m away upwind (control). **B.** Enlarged view of the
872 retention time (RT) interval containing the LRI at which *An. coluzzii* antennal responses were detected in GC-
873 EAD. Doted white lines correspond to the theoretical RT of sulcatone, octanal, nonanal and decanal. When in
874 italic, the compound was absent. No difference was detected between the control and the swarm extracts. All the
875 peaks correspond to environmental emissions.

876

877 **Fig. S4** Quantities (integration units) of acetoin, sulcatone, octanal, nonanal and decanal contained in the
878 headspace of the empty bottles, the breath of the manipulator and the breath of the manipulator after blowing for
879 the second time (breath $\times 2$) sampled with SPME fibers. The intensity values correspond to the counts related to
880 the abundance of the specific ions representative of each molecule formed in the mass spectrometer and
881 correspond to the amount of compound analyzed. The box plots indicate the median (wide horizontal bars), the
882 25th and 75th percentiles (squares), and the minimum and maximum values (whiskers). The black dots represent
883 outliers and grey dots raw data. NS=not-significant; * = $P < 0.05$; ** = $P < 0.01$.