

1 **Estimating abundance of a recovering transboundary brown bear population with capture-**  
2 **recapture models**

3 Cécile Vanpé<sup>1</sup>, Blaise Piédallu<sup>2</sup>, Pierre-Yves Quenette<sup>1</sup>, Jérôme Sentilles<sup>1</sup>, Guillaume Queney<sup>3</sup>,  
4 Santiago Palazón<sup>4</sup>, Ivan Afonso Jordana<sup>5</sup>, Ramón Jato<sup>6</sup>, Miguel Mari Elósegui Irurtia<sup>7</sup>, Jordi Solà de  
5 la Torre<sup>8</sup>, Olivier Gimenez<sup>2</sup>

6 1 Equipe Ours, Office Français de la Biodiversité, Impasse de la Chapelle, 31800 Villeneuve-de-  
7 Rivière, France

8 2 CEFE, Univ. Montpellier, CNRS, ~~Univ. Paul Valéry Montpellier 3~~, EPHE, IRD, Montpellier,  
9 France

10 3 ANTAGENE, 6 allée du Levant, 69890 La Tour de Salvagny, France

11 4 Fauna and Flora Service, Department of Territory and Sustainability, Dr. Roux, 80. 08017  
12 Barcelona, Spain

13 5 Medi Ambient, Conselh Generau d'Aran, Plaça dera Libertat, 16, 25530 Vielha, Spain

14 6 Departamento de Producción y Asistencias Técnicas, Sociedad Aragonesa de Gestión  
15 Agroambiental, S. L. U., Ctra HUV-Fornillos, 22195 Fornillos de Apiés, Huesca, Spain

16 7 Servicio de Biodiversidad, Sección de Espacios Naturales y Especies Amenazadas, Gobierno de  
17 Navarra-Nafarroako Gobernua, Pamplona-Iruñea, Spain

18 8 Unitat de Fauna, Departament de Medi Ambient, C/ Prat de la Creu 62-64, AD500 Andorra la Vella,  
19 Principality of Andorra

20

21 Correspondence: Cécile Vanpé, Equipe Ours - Office Français de la Biodiversité, Impasse de la  
22 Chapelle, 31800 Villeneuve-de-Rivière, France. Email : [cecile.vanpe@ofb.gouv.fr](mailto:cecile.vanpe@ofb.gouv.fr)

23

1 Short title: Abundance of the Pyrenean brown bear population

## 1 Abstract

2 ~~Abundance of Enumerating~~ small populations of large mammals ~~may be assessed~~can be carried  
3 out using censuses or complete counts of ~~the~~ different individuals detected over a time period; ~~so-~~  
4 ~~called~~ minimum detected (population) size (MDS). However, as a population is growing grows larger  
5 and its spatial distribution ~~is expanding~~ expands wider, the risk of under-estimating population size  
6 using MDS ~~is rapidly increasing~~ increases sharply due to the ~~rarely~~ rarely-fulfilled assumption of  
7 perfect detection of all individuals ~~of in~~ the population. ~~, and as a result,~~ tThe need to report  
8 uncertainty ~~in around~~ population size estimates consequently becomes crucial. We addressed these  
9 issues ~~within the framework of~~ fusing the monitoring framework of the critically endangered Pyrenean  
10 brown bear population that was on the edge of extinction in the mid-1990s, with only five individuals  
11 remaining, but was ~~reinforced~~ subsequently bolstered by the translocation of 11 bears ~~originated~~ from  
12 Slovenia ~~since then~~. Each year since 1996, the abundance of the population has been assessed using  
13 MDS and minimum retained (population) size (MRS), which corresponded to a reassessment of the  
14 MDS in the light of the information collected in subsequent years. We used Pollock's closed robust  
15 design (PCRD) capture-recapture models applied to the cross-border non-invasive sampling data  
16 from France, Spain and Andorra to provide the first published annual abundance estimates of the  
17 Pyrenean brown bear population, and ~~its~~ trends over time, since 2008. Annual population size  
18 increased ~~and displayed a~~ fivefold ~~rise~~ between 2008 and 2020, reaching > 60 individuals (PCRD  
19 estimate = 66.2 with 95% Credibility Interval (CI) = [64.8, 67.8]) in 2020. PCRD estimates were  
20 globally close to MRS counts over the years and had reasonably narrow associated 95% CI. We  
21 noticed that even in cases where sampling effort is large compared to population size, the PCRD  
22 estimates of population size can diverge from the MDS counts. We found individual~~Detection~~  
23 heterogeneity in detection among individuals ~~may that might~~ stem from intraspecific home range size  
24 ~~disparities~~ variation making it more likely to ~~find signs of~~ detect individuals ~~who that~~ move ~~more~~ most.  
25 We also found a lower survival rate in cubs than in adults and subadults, ~~since~~ due to the formers  
26 cubs suffering from ~~more~~ higher mortality risks ~~(such as)~~ (from infanticides by males, predations,

1 ~~mother-maternal~~ death, or abandonment)s) than ~~the latter~~ other age classes. Our study provides  
2 ~~evidence that~~ Overall, the PCRD capture-recapture modelling approach ~~can~~ provides reliable  
3 estimates of abundance the size of and trend in large mammal and demographic rates of the Pyrenean  
4 brown bear populations, together with associated uncertainty, while minimizing bias due to inter-  
5 individual heterogeneity in detection probabilities ~~and allowing the quantification of sampling~~  
6 ~~uncertainty surrounding these estimates~~. We strongly encourage wildlife ecologists and managers to  
7 use such a similar robust approach for monitoring large mammal populations. Such information is  
8 vital for informing management decision-making and assessing population conservation status.

9

10 **Keywords:** abundance estimation, capture-recapture models, non-invasive monitoring, Pyrenees,  
11 *Ursus arctos*

## 1 Introduction

2 ~~Estimating accurately~~ Accurately and precisely estimating animal population size and ~~its~~ trends  
3 over time is essential to ~~monitor~~ inform conservation status and ~~to inform~~ management decision-  
4 making (Nichols & Williams 2006). However, when animals, such as most large carnivores, are rare,  
5 elusive, solitary, largely nocturnal, highly mobile, and/or inhabiting wide-large home ranges in remote  
6 and/or rugged habitats (~~such as most large carnivores~~), population monitoring can be particularly  
7 challenging (Thompson 2013). Invasive physical tagging-based methods are ~~almost~~  
8 ~~impossible~~ difficult to implement and, so population monitoring ~~thus consequently~~ often needs to  
9 rely on non-invasive sampling methods (Long et al. 2008; Thompson 2013). Among them, ~~such~~  
10 ~~as~~ molecular tools ~~or~~ and camera trapping are commonly used methods now (e.g., Forsyth et al. 2022;  
11 Piel et al. 2022; Proctor et al. 2022) (~~Long et al. 2008; Thompson 2013~~). For species lacking unique  
12 natural individual patterns that can be identified in photos, non-invasive genotyping of DNA extracted  
13 from animal hair or scat often remains the ~~sole~~ most practical solution to estimate population  
14 abundance (Waits & Paetkau 2005).

15 Abundance of small populations of large mammals may be assessed using censuses or complete  
16 counts of ~~the different~~ unique individuals detected over a time period (Wilson & Delahay 2001;  
17 Keating et al. 2002), ~~so-called~~ known as the minimum population detected size (Solberg et al. 2006;  
18 Miotto et al. 2007; Morin et al. 2022) and abbreviated here (MDS for minimum detected (population)  
19 size). In the case of genetic identification, MDS is ~~then~~ defined as the number of unique genotypes  
20 identified among the genetic samples inside the study area (e.g., Creel et al. 2003; Solberg et al. 2006).  
21 ~~However,~~ Obtaining a MDS through exhaustive counts, such as molecular tools or camera trapping,  
22 MDS are is often expensive, time consuming, and logistically demanding (Balme, Hunter & Slotow  
23 2009; Blanc et al. 2013). In addition, as populations is growing larger and spatial distributions is  
24 expanding wider, the risk of under-estimating population size using MDS is increasing sharply due  
25 to the ~~rarely~~ rarely-fulfilled assumption of perfect detection of all individuals ~~of~~ in the population  
26 (Solberg et al. 2006; Denes et al. 2015; Staton et al. 2022; Tourani 2022), ~~and~~ the need to report

1 uncertainty ~~in~~around population estimates consequently becomes crucial (e.g., Forney 2000;  
2 McGowan, Runge & Larson 2011). To address these issues, capture-recapture (CR) ~~surveys~~models  
3 are often used to estimate population abundance while accounting for the impossibility ~~of~~to~~detecting~~  
4 ~~exhaustively~~ all individuals in a population (Otis et al. 1978). ~~While~~Whereas ~~CR models were~~  
5 originally limited to live-trapping studies, ~~CR models~~they have been specifically adapted for use with  
6 non-invasive DNA-based sampling (Lukacs 2005; Lukacs & Burnham 2005). In particular, non-  
7 invasive genetic CR models were specifically designed to account for issues such as, ~~which implies~~  
8 individual identification errors due to genotyping errors, uncertainty in the date of individual  
9 detection, and the possibility of collecting multiple samples ~~of~~from the same individual across space  
10 within a single sampling occasion (~~Lukaes 2005;~~ Lukacs & Burnham 2005; Petit & Valière 2006;  
11 Lampa et al. 2013).

12 In standard closed-population CR models (whether or not they have been adapted to non-invasive  
13 genetic sampling), the population is assumed to be closed to changes in abundance both  
14 geographically (no immigration nor emigration) and demographically (no births nor deaths).  
15 Additionally, ~~and~~ all individuals are ~~supposed~~assumed to have identical detection probabilities  
16 ~~whatever~~regardless of their individual attributes (e.g., age, body mass, social status) and habitat  
17 features (home-range location and composition) (Otis et al. 1978). ~~But~~However, these conditions are  
18 rarely fulfilled in real populations of wild mammals (e.g., Bellemain et al. 2005; Solbert et al. 2006).

19 ~~For~~Over the last decades, considerable advances to these standard models have been developed  
20 to help alleviate issues linked to closure violation and detection~~probability~~ heterogeneity (~~see a~~  
21 ~~review by~~ Lukacs & Burnham 2005). In particular, ~~the~~ Pollock's closed robust design (PCRD) CR  
22 modelling (PCRD; Pollock 1982; Kendall, Nichols & Hines 1997) was developed in a maximum-  
23 likelihood (ML) framework to study-estimate survival, temporary emigration, and animal abundance  
24 while minimizing bias due to heterogeneity in detection ~~probabilities~~ among individuals. PCRD CR  
25 models rely on several so-called primary sampling occasions, each being composed of secondary  
26 occasions. The time interval between secondary sampling occasions must be short enough to meet

1 the population closure assumption, while consecutive primary occasions should be sufficiently  
2 separated in time to allow the population to change.

3 In Bayesian statistics, past knowledge of similar experiments is encoded into a statistical device  
4 known as a prior, and this prior is combined with current experiment data to make a conclusion on  
5 the test at hand, contrary to the Frequentist approach which makes predictions on the underlying  
6 truths of the experiment using only data from the current experiment. PCRDR models were ~~also~~  
7 recently formulated in a Bayesian framework (Schofield & Barker 2011; Rankin et al. 2016), offering  
8 several advantages over the Frequentist approach, including improved estimation ~~under low~~when  
9 sample sizes are low, access to full posterior conditional probabilities of model parameters and use  
10 of prior information. However, ~~this-it~~ is only ~~for-in~~ the few last years that ~~the-a~~ Bayesian  
11 implementation of PCRDR models has been made possible without ecologists having to code  
12 ~~themselves custom-made~~their own complex sampling algorithms (Rankin et al. 2016; Riecke et al.  
13 2018).

14 In the mid-1990s after decades of persecution, the brown bear (*Ursus arctos*) population in the  
15 Pyrenees Mountains at the border of France, Spain and Andorra,~~the brown bear (*Ursus arctos*)~~  
16 ~~population, (Fig. 1) after decades of persecution,~~ was on the edge of extinction ~~in the mid-90s~~ with  
17 only five ~~relict~~ individuals remaining (Taberlet et al. 1997). Since then, the successful translocation  
18 of 11 bears ~~originating~~ from Slovenia (Quenette et al. 2019) has allowed the population to slowly  
19 demographically recover ~~slowly~~. However, the fate of this critically endangered population (UICN  
20 France et al. 2017), isolated from the nearest Cantabrian brown bear population in north-western  
21 Spain by ~~about~~ approximately 300 km, is still uncertain (Le Maho et al. 2013) with a MDS estimated  
22 at ~~64-70~~ individuals in ~~2020-2021~~ (Sentilles et al. ~~2021a~~2022) and a high consanguinity ~~rate~~coefficient  
23 F estimated in average among individuals at 0.132 in 2020 (Beaumelle 2016; Bassi 2021). ~~In this~~  
24 ~~context,~~Thus, ~~implement~~developing reliable methods to accurately estimate ~~annual~~ population  
25 abundance and its trend over time is crucial to monitor the conservation status of this brown bear  
26 population ~~threatened with extinction~~ and implement successful management plans.

1 Currently, non-invasive monitoring of the Pyrenean brown bear population relies on both  
2 systematic and opportunistic collections of bear presence signs (e.g., scats, hair, tracks, photos/videos,  
3 visual observations, damages on livestock) in the Pyrenees Mountains combined with genetic or  
4 visual individual identifications ~~non-invasive sampling of all bear presence signs collected in the~~  
5 ~~Pyrenees, either opportunistically (i.e. collection of bear data or samples by any mountain users with~~  
6 ~~no specific sampling design) or using a systematic sampling approach (i.e. specific planned operations~~  
7 ~~following a standardized procedure) (Sentilles et al. 2021a; (Sentilles, Vanpé & Quenette 2021;~~  
8 Sentilles et al. 2022). ~~Importantly, as~~ Similar to many large carnivore populations in Europe (e.g.,  
9 Bischof, Brøseth & Gimenez 2016), the Pyrenean brown bear population is transboundary and  
10 occupies a highly politically and administratively fragmented landscape, ranging across the  
11 Principality of Andorra, two administrative regions, divided ~~across~~ in six ~~different~~ counties in France,  
12 and three autonomous regions (Catalonia, Aragon and Navarra) and one Catalonian county with  
13 specific autonomous status (Val d’Aran) in Spain (Fig. 1). As such, cross-border multi-scale  
14 population monitoring cooperation (from national to local scales) is implemented to avoid population  
15 ~~size~~-overestimation, ~~due to~~ as individuals with home ranges overlapping borders may be detected in  
16 several political jurisdictions (Bischof et al. 2016; Gervasi et al. 2016).

17 To date, the size of the Pyrenean brown bear population was annually assessed using the MDS  
18 index (Sentilles, Vanpé & Quenette 2021; Sentilles et al. 2022). However, this method assumes that  
19 all individuals present in the population have a detection probability of one. Because the population  
20 size was very small compared to the intensive sampling effort (Tables S1 and S2), the number of  
21 undetected individuals was assumed to be small. As the population was assumed to be geographically  
22 closed, the MDS of the current year was used each year to correct the MDS of previous years (e.g.,  
23 to add bears which were not detected the previous years but detected the current year) and defined  
24 what we called the minimum retained (population) size, or MRS (Sentilles, Vanpé & Quenette 2021;  
25 Sentilles et al. 2022). MRS thus corresponded to a reassessment of the MDS in the light of the  
26 information collected in subsequent years. But although MRS could be regarded so far as a precise



1 and accurate estimate of the true annual brown bear population size in the Pyrenees, it does not allow  
2 uncertainty assessment and MRS for year n is only available in year n+1 and sometimes needs a  
3 reassessment on year n+2 or n+3 (Sentilles et al. 2022). In addition, with increasing Pyrenean brown  
4 bear population size and range area, the number of undetected individuals over a year increases.  
5 Finally, the outputs of demographic analyses of the Pyrenean brown bear population are used to  
6 inform management decision-making and policies (e.g., regulation, translocation, compensation). In  
7 this context, the reporting of abundance estimates and trends can be particularly prone to political  
8 influence (Darimont et al. 2018) and stakeholder skepticism. Therefore, implementing sound  
9 population monitoring tools and robust statistical methods to convey the uncertainty around  
10 abundance estimates is crucial. According to Lukacs and Burnham (2005), DNA-based CR methods  
11 provide the most useful methods to estimate abundance from small populations up to a few thousand  
12 individuals, as in the Pyrenean brown bear population.

13 The aim of this study was therefore to use cross-border non-invasive sampling data collected from  
14 2008 to 2020 in France, Spain and Andorra, ~~and~~ for which individual identification was possible  
15 through genetic analyses or visual evidence combined with PCR-D CR modeling to provide the first  
16 published estimates of annual abundance of the Pyrenean brown bear population, while minimizing  
17 bias due to heterogeneity in detection probabilities among individuals, ~~based on a robust design CR~~  
18 ~~modelling approach.~~ The development of new methods to estimate population abundance is timely,  
19 since it gives the possibility to compare the estimates obtained with the PCR-D CR modeling approach  
20 with those from census approaches (MRS and MDS counts).

21

## 22 **Material and Methods**

### 23 *Brown bear biology*

24 ~~The brown bear is part of the few species among members of the Carnivora order with an omnivorous~~  
25 ~~diet (Wroe & Milne 2007). In the Pyrenees, 70 to 80% of the diet are composed of plants (including~~

1 bilberries, cranberries, nuts, acorns, beechnuts, raspberries, ferns, sorbs, apples and rosehips), and 20  
2 to 30% are of animal origin (mainly ant larvae, bee broods, carrion, small mammals, wild and  
3 domestic ungulates) (Couturier 1954). Mating occurs in May-June, births (with litter size ranging  
4 from 1 to 3 cubs and interbirth interval being most frequently 2 years) from January to March, and  
5 hibernation between November and March (Chapron et al. 2003). Cubs remain with their mother  
6 generally for 1.4 years, but in some rare cases for 2.4 years, before dispersing (Swenson et al. 2000).  
7 Brown bears are mostly solitary animals (except for females accompanied by their cubs and rutting  
8 period), with a promiscuous mating system (Schwartz et al. 2003). Males have larger home ranges  
9 than females, with possibilities of both intra- and inter-sexually overlap (Dahle & Swenson 2003).  
10 Dispersal is sex-biased towards males, with philopatric females establishing their home ranges in or  
11 adjacent to their mothers' home ranges (Støen et al. 2005). In Europe, female and male brown bears  
12 reach sexual maturity between 3.5 to 5 years old (Chapron et al. 2003).

13

#### 14 *Brown bear population monitoring and bear sign collection*

15 ~~This~~ We carried out this study ~~was carried out~~ in the Pyrenees Mountains in ~~South-~~  
16 ~~Western~~southwestern Europe; where the cross-border population of brown bears is present in the  
17 major part of the mountain range in France, Spain and Andorra and ranges over > 10,000 km<sup>2</sup> in 2020  
18 (Sentilles et al. ~~2021a~~2022; Fig. 1).

19 We used ~~f~~Four different non-invasive methods ~~were used~~ to monitor the brown bear population in  
20 the French Pyrenees over the study period from 2008 to 2020 (Table S1):

21 ~~-1)~~ Systematic ~~by trails~~ walking (ST) ~~corresponded to walking, equivalent to transects~~ transect  
22 surveys (from 8 to 10 km long), spread homogeneously over the area of known, regular bear presence,  
23 which covers about 3,000 km<sup>2</sup> in France (~~Sentilles et al. 2021a;~~ Vanpé et al. 2021; Sentilles et al.  
24 2022; see Fig. S1). These transects were surveyed ten times (at least once per month) between May  
25 and November each year in search of bear signs by teams of two members of the Brown Bear Network

1 (i.e. > 400 professionals and volunteers trained and managed by the bear team of the French  
2 Biodiversity Agency (OFB), who is in charge by the French Minister of Ecology of brown bear  
3 monitoring in the French Pyrenees; Sentilles, Vanpé & Quenette 2021; Sentilles et al. 2022). Trails  
4 To optimize bear detection, we set transects were set in the most favourable bear areas function of  
5 available bear in terms of habitat qualitys and in bear passage areas detected using VHF and GPS  
6 collars or ~~bear~~ presence signs. Transect staff ~~accompanied occasionally by a scat detection dog~~  
7 ~~(Sentilles et al. 2021b)~~ searched for bear hair and scats on trails and in their immediate surroundings  
8 (see De Barba et al. 2010 for a similar approach). To improve the chances of getting hair samples,  
9 between five and seven hair traps were scattered along each ~~itinerary~~ trail. Each hair trap consisted  
10 ~~ofn~~ three small barbed wires fixed at three different heights onto ~~the a~~ tree and where an attractive  
11 product (i.e. turpentine until 2016, beechwood tar called “smola”smola since 2017) was applied to  
12 encourage bear rubbing ~~behaviour~~ behavior (Berezowska-Cnota et al. 2017). Some of these hair traps  
13 were associated with a facing camera trap (similar to the systematic by camera traps method described  
14 below) to help detecting females with cubs and assessing ~~the~~ age class and number of individuals that  
15 rubbed on the ~~focused focal~~ tree, as well as the date of hair deposition (Parres et al. 2020).

16 –2) Systematic by baited hair traps (SBHT) (only from 2008 to 2011), correspondinged to  
17 enclosures of about 20-30 m<sup>2</sup> delimited by a strand of barbed wire fixed at a height of 50 cm (Woods  
18 et al. 1999; Kendall & McKelvey 2008; Quinn et al. 2022) and stretched around several trees. Bait  
19 consisting ~~in of about~~ ~ 1-L mixture of rotten blood and fish was poured into the center of the area,  
20 with a reward ~~consisting in of~~ corn grains to increase recapture probability (see Woods et al. 1999;  
21 Castro Arrellano et al. 2008; Gervasi et al. 2010). ~~We used a 4 x 4 km grid cell size on the known~~  
22 ~~female range area and a 8 x 8 km grid cell size on the remaining part of the study area and placed one~~  
23 ~~baited station on each grid cell.~~ The trapping grid was established following designs and guidelines  
24 outlined in previous DNA-based inventories in North America (Mowat & Strobeck 2000; Boulanger  
25 et al. 2002) and ~~considering the~~ average adult female home ranges of brown bears in the Pyrenees.  
26 The average home range size (Kernel 85%) of brown bears in the Pyrenees (excluding recently

1 translocated individuals) was 84 km<sup>2</sup> in adult females (N = 6) and 1,551 km<sup>2</sup> in adult males (N = 6)  
2 (Halotel et al. unpubl. data; similar to the average home range of radio-collared adult bears in similar  
3 Eurasian regions: Huber & Roth 1993; Mertzanis et al. 2005; Gavrillov et al. 2015). We used a 4 x 4  
4 km grid cell size based on the known female range areas and a 8 x 8 km grid cell size on for the  
5 remaining part of the study area, and placed with one baited station placed in each grid cell. Hair  
6 traps were placed in the best predicted bear habitat, considering topography and accessibility by 4-  
7 wheel drive vehicles, a maximum of 10 min walk from the vehicle and ~~taking into account~~ bear expert  
8 opinion (tree types or tree species, with characteristics that make them more conspicuous for rubbing;  
9 González-Bernardo et al. 2021; Proctor et al. 2022). Sites were visited once every 15 days from May  
10 to September for sample collection and lure replacement.

11 ~~—3)~~ Systematic by camera traps (SCT), ~~corresponded~~ corresponding to ~~automatic triggered~~  
12 cameras (~~essentially~~ Leaf river Outdoor, HCO Soutguard SG 550 and Uway Nicht Trakker until 2013,  
13 and Bushnell Trophy Cam or NatureView HD and Reconyx HC600 or XR6 after 2013) equipped  
14 with movement detection that were fixed on trees in areas with frequent animal passages ~~outside away~~  
15 from the walking transects and that were associated ~~closed~~ near by with hair traps similar to the ones  
16 used for the systematic by trails method (Burton et al. 2015; Parres et al. 2020; see Fig. S1). Frequent  
17 animal passages were defined here as animals' trails from all large mammals, which are visible in the  
18 vegetation and on the ground and that are often used by bears, as well as bear passage areas detected  
19 using VHF and GPS collars or bear presence signs. Each camera trap - hair trap station was visited  
20 once per month from April to November each year to collect samples and maintain cameras (Sentilles,  
21 Vanpé & Quenette 2021; Sentilles et al. 2022). We followed the same layout as above for SBHT  
22 protocol used a 4 x 4 km grid cell size on the known female range area and a 8 x 8 km grid cell size  
23 on the remaining part of the study area and placed one camera trap - hair trap station per cell. When  
24 hair samples could non-ambiguously be associated with photographs or videos, we analysed ~~collected~~  
25 pictures in an attempt to individually identify bears based on natural ~~(e.g., coat marks)~~ or artificial

1 ~~(markings, ear tags, radio-or collars)-marks~~, in order to avoid genetic analyses and decrease sampling  
2 costs.

3 ~~-4)~~ Opportunistic monitoring (OM), ~~corresponded~~ corresponding to the opportunistic collection  
4 (with no specific sampling design) throughout ~~the bear~~ potential bear range (covering > 10,000 km<sup>2</sup>)  
5 of all bear presence signs (such as hair, scats, tracks, claw marks on trees~~scratches~~, eating-feeding  
6 clues, visual observations...) gathered by ~~various-any~~ mountain users (e.g., hikers, foresters, hunters,  
7 skiers, fishermen, shepherds), as well as all putative bear damages on livestock and beehives, ~~after~~  
8 ~~examination and approval of an expert agent~~ (De Barba et al. 2010). Potential bear range is defined  
9 here as the areas surrounding bear presence, allowing random locations (for bear absences) to fall  
10 where bears could have visited (15 km from the edge of presence), as defined in Martin et al. 2012.  
11 Feeding clues are carcasses of wild or domestic preys, overturning of a large stone, and anthill and  
12 bee or wasp swarms burst open. Mountain users report their observations to the bear team of the OFB.  
13 Testimonies are examined and approved by an expert from OFB. A conclusion as to its validity as  
14 bear evidence, "confirmed," "probable," "doubtful," or "false," is given to each putative bear presence  
15 sign that could be verified, on the same day or a few days after its transmission, according to the  
16 elements necessary for their verification (Sentilles et al. 2022). Bear observations are validated only  
17 if an indirect bear clue (scats, hair, footprints) is found at the sighting site or if a photo or video is  
18 provided by the observer. To confirm that eating clues are from brown bears, we specifically look for  
19 evidence of associated bear clues close by (e.g., footprints, claw marks, hair, scats). If the elements  
20 are not sufficient to make a decision or if the observer could not be found for the statement of his/her  
21 testimony, the evidence is classified as "impossible expertise". Only confirmed bear clues are  
22 included in our analyses. Since 2014, verification of testimonies and damage reports have been  
23 occasionally carried out with the help of a scat-detection dog trained to search for brown bear scats  
24 (Sentilles, Vanpé & Quenette et al. 2021b). Only hair and scat samples collected during the same  
25 period (from May to November) as the ST systematic monitoring were included. ~~in this study.~~

26

1 While all the four ~~monitoring protocols methods~~ (ST, SBHT, SCT, OM) were used in France,  
2 brown bear monitoring consisted ~~of only~~in the ST ~~method combined with~~and OM ~~protocols~~ in  
3 Catalonia (Spain) and Andorra, and ~~only the~~in OM ~~protocol~~only in Aragon and Navarra (Spain). ~~But~~  
4 ~~note that this should not affect bear detection and population abundance estimation, since the choice~~  
5 ~~of the monitoring methods was not dictated by the country or administrative unit but rather by the~~  
6 ~~regularity of bear presence in the area (ST was implemented only in areas of known, regular bear~~  
7 ~~presence in France, Spain and Andorra, while OM was implemented everywhere within the potential~~  
8 ~~brown bear presence area).~~ Although few individuals (mostly translocated animals and problematic  
9 bears) were temporally equipped with either VHF and/or GPS collars or ear tags over the study period,  
10 we ~~analysed only~~~~focused here on~~ the ~~sole~~ non-invasive sampling data. ~~For all the four protocols, w~~We  
11 paid ~~a~~ particular attention ~~when to evaluate~~ ~~evaluating~~ the date when the signs were left by the bears  
12 and discarded any sign for which uncertainty in ~~this the~~ date was too high to define ~~precisely~~ which  
13 month the bear was present (see Supplementary Materials). ~~This study complies with the standards,~~  
14 ~~laws, and procedures concerning animal research ethics of the countries, in which it was performed.~~

15

#### 16 *Individual identification of bear signs*

17 We used all validated non-invasive brown bear signs collected in the Pyrenees from 2008 to 2020  
18 (Table S2) ~~and~~ for which individual identification was possible. Individual identification of bears was  
19 ~~mainly~~ ~~primarily~~ based on genetic analyses of hair (stored dry in envelopes) and scats (stored in  
20 microtubes filled with 96% ethanol) non-invasively collected in the field, as well as visual evidence  
21 (colouration, scars, GPS collars, or VHF ear tag transmitters) obtained by remote cameras ~~when~~  
22 ~~available~~ (Sentilles et al. 2021b). ~~This study complies with the standards, laws and procedures~~  
23 ~~concerning animal research ethics of the countries, in which it was performed.~~ ~~This visual~~  
24 ~~identification was performed by bear experts from OFB and was validated only if a consensus was~~  
25 ~~released among all those experts without any doubt.~~

1 Due to financial constraints, only a subset of all collected hair and scat samples were genetically  
2 analysed to identify individuals each year (mean  $\pm$  SD = 35.16  $\pm$  12.29, min = 17.5 in 2015 and max  
3 = 59.5 in 2008; Table S2). Samples that were sent to the lab each year were carefully selected so that  
4 we optimised the detection of individuals (e.g., we favoured samples from cubs of the year or  
5 subadults, as well samples that were collected in the expansion front of the population) and the  
6 genotyping success (e.g., freshest scats, avoidance of hair coming from different individuals).

7 Genetic samples were analyzed at the Laboratoire d'Ecologie Alpine (LECA) joint research unit  
8 ~~LECA-CNRS laboratory~~ from 2008 to 2012 using a multiple-tubes Polymerase Chain Reaction  
9 (PCR) approach (consisting in repeating each DNA amplification independently for each locus  
10 several times; Taberlet et al. 1996, 1997) and from 2013 to 2016 using high-throughput microsatellite  
11 genotyping on ~~the an-~~ Illumina platform (De Barba et al. 2017). From 2017 to 2020, samples were  
12 analyzed in our laboratory at ANTAGENE Company using a new multiple-tubes PCR approach (see  
13 the methods and Table S3 provided in Supplementary information ~~for method details~~). In all cases, a  
14 minimum of four repeats for each sample was carried out to avoid genotyping errors associated with  
15 low quantities of DNA (Miquel et al. 2006). A total of 13 microsatellites markers and one (for LECA-  
16 ~~CNRS~~) to three (for our laboratory) sex markers ~~targeting~~ were targeted by the multiplex PCR in  
17 order to identify ~~the bear~~ individuals and assign gender (De Barba et al. 2017; ~~see and~~ the methods  
18 and Table S4 provided in Supplementary information). Further information on genotyping error rate  
19 and ~~probabilities-probability~~ of identity-by-descent can be found in De Barba et al. (2017), Beaumelle  
20 (2017), Bassi (2021) and Table ~~S2S4~~.

## 21 22 *Population abundance estimation using capture-recapture models*

23 The results from all sources of individual identifications (genetic analyses and tracking of natural  
24 or artificial marks) of all bear signs for which the month when the bear left the sign was known were

1 then aggregated to compile ~~the a~~ monthly detection history ~~of for~~ each bear ~~of in~~ the population  
2 ~~through time~~ from January 2008 to December 2020 (see Supplementary Materials).

3 We used ~~the a~~ PCRD CR model (Pollock 1982; Kendall, Pollock & Brownie 1995; Kendall et al.  
4 1997; see also in Williams, Nichols & Conry 2022) to estimate population abundance. This method  
5 has been applied on a number of bear populations (Stetz et al. 2010; Pederson et al. 2012; McCall et  
6 al. 2013; Tosoni et al. 2017see also in-). PCRD CR models use a hierarchical sampling strategy,  
7 including widely-spaced “primary occasions”, between which the population is considered as open  
8 (i.e. with births, deaths and temporary emigration), and repeated captures in a short timeframe (called  
9 “secondary occasions”) between which the population is assumed to be closed to population changes.  
10 Data from secondary samples within each primary period are analyzed using closed models to derive  
11 estimates of detection probability and population size. Apparent survival and temporary emigration  
12 are estimated using open models by collapsing data from the secondary periods. Here, temporary  
13 emigration refers to some individuals that might temporarily emigrate to areas where foraging  
14 conditions or breeding success are better, or that might be temporarily unavailable for capture because  
15 they are ~~dormant, in torpor, or hibernating~~in dens (Henle & Gruber 2017). ~~while accounting for~~  
16 ~~imperfect detection of individuals and the temporary absence of some individuals from a sampling~~  
17 ~~site (e.g., individuals may temporarily emigrate to areas where foraging conditions or breeding~~  
18 ~~success are better, or may be temporarily unavailable for capture because they are dormant, in torpor,~~  
19 ~~or hibernat~~hibernating; Henle & Gruber 2017). PCRD CR models use a hierarchical sampling  
20 strategy, including widely spaced “primary occasions,” between which the population is considered  
21 as open (i.e. with births, deaths and temporary emigration), and repeated captures in a short timeframe  
22 (called “secondary occasions”) between which the population is assumed to be closed to population  
23 changes.

24 The population was assumed geographically closed, i.e. no emigration or immigration could occur  
25 between this population and another one outside the Pyrenees. We used years from 2008 to 2020 as  
26 primary occasions of capture (N = 13) and months from May to September as secondary occasions



1 (N = 5), that is 65 occasions of capture in total. We chose these secondary occasions because no births  
2 occur in this time interval. We excluded months from October to April because of low activity of  
3 bears during hibernation and high mortality risks of cubs of the year during their first months of life  
4 (bear cubs are born in the den during January-February).

5 PCRD CR models allow estimating population abundance, detection probability and apparent  
6 survival while accounting for temporary emigration (Pollock 1982; Kendall, Nichols & Hines 1997).  
7 We accounted for temporary emigration with two parameters. First we used the probability of an  
8 individual being a temporary emigrant, given it was alive and present in the study area in the previous  
9 primary sampling occasion. The other temporary emigration parameter is the probability of an  
10 individual being a temporary emigrant given it was a temporary emigrant in the previous sampling  
11 occasion. There is no temporary emigration when both parameters are 0,– random temporary  
12 emigration when both parameters are set and estimated equal (and the probability of an individual  
13 being present in the study area is not dependent on whether or not it was present in the study area in  
14 the previous sampling period) and Markovian temporary emigration when both parameters are set  
15 and estimated distinct (and the probability of an individual being present in the study area is  
16 conditional on whether it was present in the study area before). Apparent survival rate is the  
17 probability of surviving and staying in the study area, and is the product of true survival and fidelity  
18 to the study area. We used a ~~classical~~-frequentist approach fitting 24 different models in total to  
19 explore effects on survival, detection and temporary emigration structure (Murray & Sandercock  
20 2020). We ~~considered fitted 24 different models in total, with~~ four detection structures (constant,  
21 time-dependent considering variation between and within primary occasions and heterogeneous using  
22 finite mixtures, in which individuals may belong to one class of animals with a some detection  
23 probability in some proportion  $\pi$  or to another class of animals with a different detection probability  
24 in proportion  $1 - \pi$ ), two survival structures (constant and age-dependent using three age classes: i.e.  
25 cubs < 2 year old, subadults = 2-3 years old and adults > 3 years old) and three emigration structures  
26 (constant, random and Markovian) (see Table 1). We used the Akaike Information Criterion corrected

1 for small sample size (AICc) to perform model selection (Burnham & Anderson 2002). These  
2 analyses were performed with the ‘RMark’ package (Laake 2013) that allows calling the Mark  
3 program (White & Burnham 1999) from R software (RCoreTeam 2013). Because we ~~run~~ran into  
4 boundary ~~estimates~~estimation issues, we used a Bayesian approach to estimate annual population  
5 abundance, relying on the best supported model from the frequentist approach. These analyses were  
6 performed using program Jags (Plummer 2003; and Riecke et al. 2018 for PCRD models in  
7 particular). The rationale in considering both frequentist and Bayesian frameworks was to use the  
8 advantages of each of them: tThe Frequentist framework allows model selection via AICc without  
9 prohibitive computation time, and the Bayesian framework allows for obtaining interpretable  
10 estimates. Data and codes are available at [https://github.com/oliviergimenez/pyrenean-brown-bear-](https://github.com/oliviergimenez/pyrenean-brown-bear-abundance)  
11 [abundance](https://github.com/oliviergimenez/pyrenean-brown-bear-abundance).

12 We compared PCRD estimates of the annual Pyrenean brown bear population abundance with  
13 both MDS and MRS counts. Note that MRS for 2020 is provisional and will be reassessed in the  
14 future (see above).

## 16 **Results**

### 17 *Individual identification*

18 From 2008 to 2020, we had ~~in total~~ 10,019 validated brown bear signs (e.g., hair, scats, tracks,  
19 visual observations, damages, photos / videos) collected ~~in the whole~~throughout the Pyrenees year-  
20 round (Table S2). Among the 2,524 hair and scat samples, which were sent ~~for~~to genetic analyses in  
21 France over this period, 1,648 (~~about~~ 65%) allowed individual identification (Table S2). From 2008  
22 to 2020, 98 different individuals (44 females, 41 males and 13 individuals with undetermined sex)  
23 were identified in the ~~whole~~ Pyrenees from May to September. Those individuals have been detected  
24 from 1 to 61 different capture occasions (median = 5.5, mean  $\pm$  SD = 10.25  $\pm$  12.23) over the study  
25 period from 2008 to 2020 (which include 65 occasions of capture in total).

1

## 2 *Model selection*

3 The two top ranked models best supported by the data (with  $\Delta\text{AICc} < 2$ ) among the 24 fitted  
4 models both included age-dependent survival, heterogeneous detection, and either random or  
5 Markovian emigration ~~effects~~ (Table 1). All other models had much higher AICc ( $\Delta\text{AICc} > 6$ ; Table  
6 1). Survival estimates of cubs, subadults and adults were nearly identical for both top ranked models  
7 around (mean  $\pm$  SE = 84.4  $\pm$  3.8%, 95.4  $\pm$  2.8% and 96.2  $\pm$  1.5%, respectively, except that the SE of  
8 the Markovian model is 2.9% instead of 2.8% as for the random model for subadults; for both top  
9 ranked models (Table 2). ~~Regarding the heterogeneous detection, 0.72%~~ of individuals had a low  
10 detection probability of 42%, whereas ~~0.28%~~ of individuals had a high detectable probability of 85%  
11 (Table 2). The probability of leaving the study area was  $< 10\%$  for both models, whereas the  
12 probability of remaining outside the study area was ~~about~~ 22% (Table 2).

13

## 14 *Abundance estimation*

15 Based on the best-supported model from the frequentist analysis (Table 2), we ran a Bayesian  
16 PCRD CR model, in which temporary emigration is random, survival is age-dependent survival and  
17 there is heterogeneity in the detection process. We used this model (see Table S5 for estimated  
18 parameters) to estimate annual abundance of the Pyrenean brown bear population. Bayesian PCRD  
19 estimates of the Pyrenean brown bear annual population abundance ranged from 13.0 with 95%  
20 credible interval (95% CI) = [12.8, 13.3] in 2008 to 66.2 with 95% CI ~~95% credible interval~~ = [64.8,  
21 67.8] in 2020 (Fig. 2 and Table 4S6). We observed an increasing trend, with annual abundance  
22 displaying a fivefold rise between the beginning and the end of the study, with reasonably narrow  
23 95% CI (Fig. 2 and Table S6).

24 Differences in the estimates of the annual abundance of the Pyrenean brown bear population  
25 between the Bayesian PCRD CR modelling approach and census methods remained relatively small

1 over the years (except from 2017 to 2019), with globally closer values between PCR-D and MDS than  
2 between PCR-D and MRS counts (mean difference  $\pm$  SD =  $-1.02 \pm 2.27$  and  $-3.79 \pm 3.95$ , respectively  
3 ; Fig. 2 and Table S6). While PCR-D estimates were either higher or smaller than MDS depending on  
4 the year, they were consistently smaller than MRS over the years except in 2009 and 2016 ( $+1.36$  and  
5  $+0.24$ , respectively; Fig. 2 and Table S6).

6 While MRS and MDS counts remained very close to each other before 2017 (mean difference  $\pm$   
7 SD =  $0.89 \pm 1.45$ ), differences between MRS and MDS as well as between MRS and PCR-D became  
8 much larger from 2017 ( $7.00 \pm 3.56$  and  $7.65 \pm 4.65$ , respectively; Fig. 2 and Table S6), considering  
9 that MRS for 2020 is provisional and will probably be reassessed upwards.

## 13 Discussion

14 ~~Based on the combination of non-invasive genetic sampling of hair and scats and corresponding~~  
15 ~~track size data,~~ The Pyrenean brown bear population was shown to be composed at least of at least  
16 five individuals in 1995, indicating that population was then at the edge of close to extinction (Taberlet  
17 et al. 1997). To attempt preserve-preserving the remaining Pyrenean gene pool, and increase genetic  
18 diversity and revive the population dynamics, the translocation of a total of 11 bears originating from  
19 Slovenia was performed from 1996 to 2018 (Quenette et al. 2019). To assess the effectiveness of  
20 these conservation efforts and the current conservation status of the Pyrenean brown bear population,  
21 it is important to evaluate how the population size has evolved since the first translocations. We used  
22 PCR-D CR models applied to the cross-border non-invasive sampling data from France, Spain and  
23 Andorra to provide the first published annual abundance estimates and trend of the critically  
24 endangered Pyrenean brown bear population and its trends over time from 2008 to 2020 since the  
25 first translocations that occurred in 1996.

1 Our results suggest that ~~annual-the~~ size of the Pyrenean brown bear population ~~showed rapid~~  
2 ~~population growthincreased-,\_and-displayed-displaying~~ a fivefold rise between 2008 and 2020,  
3 reaching > 60 individuals (~~PCRD estimate = 66.2 with 95% CI = [64.8, 67.8]~~) in 2020. ~~Most of the~~  
4 ~~11 translocations occurred before 2008 (2 females in 1996, 1 male in 1997, 4 females and 1 male in~~  
5 ~~2006). Hence, the increase we observed in annual population size from 2008 to 2020 is not due~~  
6 ~~essentially to the translocation of new individuals in the population during the study period (which~~  
7 ~~concerns only 1 male in 2016 and 2 females in 2018), but mainly to the reproduction of an increasing~~  
8 ~~number of individuals (Bassi 2021; Sentilles, Vanpé & Quenette 2021). Note that the important~~  
9 ~~increase in the population abundance from 2018 cannot be explained by a sex ratio biased towards~~  
10 ~~adult females, since the sex ratio among adults has been systematically biased towards females since~~  
11 ~~2012 (see Table S7). While this demographic success is encouraging for the short-term viability of~~  
12 ~~the population, the fate of this critically endangered population is still uncertain due to high~~  
13 ~~consanguinity, geographic isolation, fragmentation and small population size, which makes it~~  
14 ~~particularly vulnerable to demographic, environmental and genetic stochasticity (Chapron et al. 2009;~~  
15 ~~Le Maho et al. 2013; Beaumelle 2016; Bassi 2021).~~

16 To date, the size of the Pyrenean brown bear population was annually estimated using the MDS  
17 ~~index, defined as the minimum number of different individuals detected inside the study area over~~  
18 ~~the year (Table S4). This method assumes that all individuals present in the population have a~~  
19 ~~detection probability of one. Because the population size was so far very small compared to the~~  
20 ~~intensive sampling effort (Table S1), the number of undetected individuals was considered each year~~  
21 ~~as very small. As the population was assumed geographically closed, the MDS of the current year~~  
22 ~~was used every year to correct the MDS of previous years (e.g., to add bears which were not detected~~  
23 ~~the previous years but detect the current year) and defined what we called the Minimum Retained~~  
24 ~~Size (MRS; Sentilles et al. 2021a,b). MRS thus corresponded to a reassessment of the MDS in the~~  
25 ~~light of the information newly collected in the following years. However, note that MRS estimation~~  
26 ~~can be subject to sampling bias if some specific individual types (e.g., more detectable individuals or~~

1 individuals still alive) are more prone to be detected a posteriori. While from 2008 to 2016, the MRS  
2 and MDS of the Pyrenean brown bear population remained very close from each other (mean  
3 difference  $\pm$  SD =  $0.9 \pm 1.5$ ), the difference between the two estimates becomes much larger from  
4 2017 to 2020 (mean difference  $\pm$  SD =  $7.3 \pm 3.2$ ; Table S4 and Fig. 2). This suggests that the size and  
5 distribution range of the Pyrenean brown bear population have now reached a point that we cannot  
6 anymore neglect the risk of failing at detecting all individuals of the population over a year using  
7 MDS, especially for years, during which a limited number of samples can be sent to genetic analyses  
8 due to funding restrictions (such as in 2017 and 2018; see Table S1 for details). Consequently, it  
9 becomes crucial for the monitoring of the Pyrenean brown bear population to estimate population  
10 size using a method that account for individual heterogeneity in detection probabilities and to report  
11 uncertainty on estimates. This is why implementing a new reliable method of estimation of annual  
12 population abundance combining capture-recapture modelling and non-invasive sampling was  
13 particularly relevant for our study population at this stage.

14 Differences between PCRD estimates of the annual abundance of the Pyrenean brown bear  
15 populationWe observed that PCRD estimates of the annual abundance of the Pyrenean brown bear  
16 population were close to and MRS or MDS values counts over the years (except from 2017 to 2019)  
17 and had reasonably narrow associated 95% CI (Fig. 2 and Table S6). were relatively small (mean  
18 difference  $\pm$  SD =  $-3.79 \pm 3.77$  and  $-1.02 \pm 2.27$ , respectively), with PCRD estimates being either  
19 higher or smaller than MDS and MRS values depending on the year. The fact that PCRD estimates  
20 are usually lower than MRS counts over the years (and to a lesser extent, MDS counts) Those  
21 differences could be explained by the fact that our PCRD CR framework includes temporary  
22 emigration, which means that a bear that is not found during an entire year will not be included in the  
23 total population size estimate. Moreover, to use the PCRD CR framework, we excluded signs that  
24 were difficult to date, and those that fell outside of the secondary occasions (May to September),  
25 which left some individuals identified by MDS and MRS out of our database. Furthermore, MDS and  
26 MRS estimates counts performed so far always included the individuals that were found dead in their

1 yearly counts, while a PCRD CR model would only include them if the death occurred after the end  
2 of the primary occasion from October to December. Despite these limitations, our results suggest that  
3 the PCRD CR method provides reliable estimates of the size and trend of the Pyrenean brown bear  
4 population, while minimizing bias due to inter-individual heterogeneity in detection probabilities and  
5 quantifying sampling uncertainty surrounding these estimates.

6 The larger differences between MRS counts and both PCRD estimates and MDS counts in 2017  
7 and 2018 (Fig. 2 and Table S6) may be partly explained by the fact that a limited number of DNA  
8 samples could be collected during these two years (N = 569 and 601, respectively) due to intensive  
9 translocation preparation efforts, compared for instance to 2015 and 2016 (N > 800; Table S2). This  
10 could result in a higher proportion of undetected individuals over the year, that could have been  
11 redetected during the following years. However, a large difference between MRS counts and both  
12 PCRD estimates and MDS counts was also observed in 2019 (9.08 and 7.00, respectively; Fig. 2 and  
13 Table S6), even though >800 DNA samples were collected over the year, among which 38% were  
14 analysed and 25% could be successfully genotyped (compared to  $35.16 \pm 12.29$  % and  $22.17 \pm 7.22$   
15 %, respectively, in average from 2008 to 2020; Table S2). In addition, the difference between MRS  
16 counts and PCRD estimates was not positively correlated to the proportion of collected samples that  
17 were genetically analysed ( $F_{1,11} = 0.436$ ,  $P = 0.52$ ). The accentuation of the differences between MRS  
18 and MDS counts at the end of the study period (2020 excluded due to provisional MRS) thus likely  
19 indicates that we have now reach a point for which it becomes more and more difficult to detect all  
20 individuals over a year, even with intensive sampling and genotyping efforts. As a consequence, the  
21 development of new metrics using capture-recapture methods to replace the MDS census approach to  
22 estimate the abundance of the Pyrenean brown bear populations is timely.

23 The model selection results highlighted two classes of individuals with significantly different  
24 detection probabilities (Table 2). A previous study on wolves highlighted the importance of  
25 accounting for individual heterogeneity in detection when estimating abundance of large carnivore  
26 populations (Cubaynes et al. 2010). Heterogeneity in the Pyrenean brown bears might stem from

1 intraspecific home range disparities (McLoughlin, Ferguson & Messier 2000) making it more likely  
2 to find signs of individuals ~~whothat~~ move a lot, as well as from the fact that few bears were more  
3 easily visually identified due to their specific natural and/or artificial marks. The ~~four~~-three  
4 individuals with long detection history ( $N > 20$  occasions) that were detected more frequently over  
5 the study period ( $> 85\%$  of occasions) were indeed all large-sized adult big-males with particularly  
6 large home ranges and which were easily visually identified thanks to natural or artificial marks: Néré  
7 (detected at 61 of the 65 occasions during which it was present), Pyros (detected at 41 of the 45  
8 occasions during which it was present), Goiat (detected at 22 of the 24 occasions during which it was  
9 present) and Balou (detected at 28 of the 32 occasions during which it was present). Conversely,  
10 among the 10 individuals with long detection history ( $N > 20$  occasions) that had the lowest detection  
11 probability ( $< 30\%$  of occasions), we had both males and females and we did not observe any age  
12 effect. Natural and/or artificial marks (colouration, scars, GPS collars, or VHF ear tag transmitters)  
13 may have helped temporally or permanently identifying some of the individuals of the population on  
14 photos or videos, causing potentially a bias in detection probabilities among individuals each month.  
15 However, this issue concerned only a few individuals each year (for natural marks: between 0 and 3  
16 individuals according to years; for artificial marks: 2 individuals in 2008-2009, 0 in 2010-2015, 1 in  
17 2016-2018, 4 in 2019 and 1 in 2020) and a few indices per individual (since natural marks are cryptic  
18 and not always visible on photos and videos). And in the vast majority of cases, these individuals  
19 have also been detected independently each month through genetics on scats and hair. So we are  
20 confident this should not have significantly affected individual capture histories.

21 Another factor that might have caused heterogeneity in detection and might have affected the  
22 abundance estimate is the efficiency of human agents when looking for bear signs. Some Pyrenean  
23 bears (e.g., dominant adult males and few adult females ~~such as Caramelles and Nheu~~) displayed a  
24 stable spatial behavior over the years (Camarra et al. 2015), making their movements predictable in  
25 time and allowing the agents to become better at finding their signs (Fagen & Fagen 1996). Extending  
26 our approach to spatial capture-recapture (SCR) models that account for individual heterogeneity in



1 the detection process by estimating individual-specific activity could help ~~alleviating~~alleviate those  
2 issues (Royle et al. 2014; Borchers & Fewster 2016).

3  
4 Interestingly, PCRD CR modelling approach provides not only estimates of abundance but also  
5 estimates of demographic rates that cannot be provided by census approaches (MDS and MRS). We  
6 found an age-dependent effect on survival, with cubs surviving less well (84%) than subadults (95%)  
7 and adults (96%; Table 2). These results are consistent with previous estimates from Chapron et al.  
8 (2009) in the same population ( $0.77 \pm 0.11$  for cubs,  $0.90 \pm 0.09$  for yearlings, 1.00 for sub-adults,  
9 and  $0.97 \pm 0.03$  for adults in the Central sub-population between 1993 and 2005) and from cub  
10 survival estimates from most brown bear populations around the world (e.g., in British Columbia,  
11 Canada: 0.86 (0.74–0.96); McLellan 2015; in the Southern Scandinavian populations: 0.72; Swenson  
12 et al. 1997). In contrast, our cub survival estimate in the Pyrenees is much smaller than what was  
13 found in Northern Scandinavia (0.98; Swenson et al. 1997). However, cub mortality is known to vary  
14 widely among populations according to food availability, human disturbance and hunting  
15 management, with bear hunting affecting either positively or negatively cub survival depending on  
16 populations (Swenson et al. 2001). In the Pyrenees, bear hunting is prohibited and food availability  
17 is considered as good, but human disturbance can occur through various human activities including  
18 mountain outdoor activities, forestry, livestock farming, road traffic and hunting (Martin et al. 2012).  
19 The lower survival rate ~~in~~of cubs compared to other age classes was expected, since cubs are known  
20 to suffer from many mortality risks such as infanticides, predations, ~~mother~~maternal death, or  
21 abandonments (Bunnell & Tait 1985) during their first year of life. In Scandinavia, about 80% of all  
22 cub mortality occurs during the mating season and is due to infanticide by males (Frank et al. 2017).  
23 While only a few infanticide, mother death and abandonnement cases were reported in the Pyrenees,  
24 their importance are probably greatly underestimated, since bear monitoring in the Pyrenees is mostly  
25 based on non-invasive methods. In addition, our estimate of cub survival is likely to be overestimated  
26 since our analyses do not take into account cub mortality at a very early age (< 4 months old) as we

1 considered months from May to September as secondary occasions and births occur in the dens in  
2 January-February (Spady et al. 2007). As a result, some cubs may have died before we could even  
3 detect them for the first time. But These cub mortality risks are not restricted to their first four or

4 five months of their life (which were excluded from our analyses as we considered months from

5 May to September as secondary occasions) but and can also occur after April during late spring and

6 summer.

7 ~~The outputs of demographic analyses of the Pyrenean brown bear population are used to inform~~  
8 ~~management decision-making and policies (e.g., regulation, reinforcements, compensation). In this~~  
9 ~~context, the reporting of abundance estimates and trends can be particularly prone to political~~  
10 ~~influence (Darimont et al. 2018) and stakeholder skepticism. Therefore, implementing sound~~  
11 ~~population monitoring tools and robust statistical methods to convey the uncertainty associated to~~  
12 ~~abundance estimates is crucial. Our results suggest that annual size of the Pyrenean brown bear~~  
13 ~~population displayed a fivefold rise between 2008 and 2020, reaching > 60 individuals in 2020. This~~  
14 ~~increase is mainly due to translocations of bears originated from Slovenia (1 male in 2016 and 2~~  
15 ~~females in 2018) combined with regular reproduction events during the study period (Sentilles et al.~~  
16 ~~2021b). While this is encouraging for the short term viability of the population, the fate of this~~  
17 ~~critically endangered population (UICN France et al. 2017) is still uncertain due to high~~  
18 ~~consanguinity, geographic isolation, fragmentation and small population size, which makes it~~  
19 ~~particularly vulnerable to demographic, environmental and genetic aleas (Chapron et al. 2009; Le~~  
20 ~~Maho et al. 2013; Beaumelle 2016).~~

21 Although the number of individuals within a population is commonly considered as a fundamental  
22 ecological indicator, the trend in population abundance can be a poor predictor of population viability,  
23 especially when strong inbreeding occurs and total population size is much higher than the effective  
24 population size, as it is the case in the Pyrenean brown bear population (Beaumelle 2016; Bassi 2021).  
25 Brown bear females in Europe usually start reproducing at the age of four or five with an interbirth  
26 interval of at least two years (Schwartz et al. 2003, Swenson et al. 2007). Therefore, to improve the

1 ~~assessment of the conservation status and of the demo-genetic viability of this critically endangered~~  
2 ~~population, using a set of indicators by monitoring the annual number of females with cubs of the~~  
3 ~~year (e.g., Palomero et al. 2007), the annual total number of  $\geq 4$  year-old females in the population,~~  
4 ~~or the effective population size (Frankham 1995; Bassi 2021), in addition to PCR-D estimates of the~~  
5 ~~total population abundance, would be particularly relevant (Beissinger & Westphal 1998).~~

6 In conclusion, Our study shows provides evidence that the PCR-D ~~capture-recapture-CR~~  
7 modelling approach allows correcting for imperfect detection to provides reliable estimates of  
8 abundance the size of and demographic rates trend in large mammal of the critically endangered  
9 Pyrenean brown bear populations, while ~~minimizing bias due to inter-individual heterogeneity in~~  
10 ~~detection probabilities and~~ quantifying sampling uncertainty surrounding these estimates. Even in  
11 cases where sampling effort is large compared to population size, the PCR-D CR abundance estimates  
12 can diverge from the minimum number known to be alive (MRS). In addition, MRS is obtained with  
13 at least one year's delay, and the census approach is logistically and financially demanding. In the  
14 context of the demographic growth and geographical expansion of the Pyrenean brown bear  
15 population, we therefore recommend using our PCR-D CR method rather than the former MDS metric  
16 to estimate the annual abundance and monitor the trend of this critically endangered population. Such  
17 ~~information is vital for informing management decision-making and assessing population~~  
18 ~~conservation status. We recommend for monitoring the size of the Pyrenean brown bear population~~  
19 ~~using this PCR-D capture-recapture modelling approach in place of the former MDS metric, which~~  
20 ~~increasingly failed over the last few years to detect all individuals of the population.~~

## 23 Acknowledgements

24 We would like to acknowledge all field co-workers from the Brown Bear Network (ROB) and from  
25 Spanish and Andorran teams. We thank the editors and the three reviewers, Tim Coulson, Romain

1 Pigeault and Susannah Woodruff, for their constructive comments and questions pertaining to the  
2 manuscript. We are grateful to Tim Coulson and Marci Johnson for thorough English language editing  
3 and checking. Financial and logistical support for this study was provided by the ONCFS / OFB in  
4 France. OG was supported by a grant from ‘Mission pour l’interdisciplinarité’ of CNRS, through  
5 its ‘Osez l’interdisciplinarité’ call.

6

7 **Conflict of interest disclosure**

8 The authors of this article declare that they have no financial conflict of interest with the content of  
9 this article.

## 1 References

- 2 [Balme, G.A., Hunter, L.T. & Slotow, R.O.B. \(2009\). Evaluating methods for counting cryptic](#)  
3 [carnivores. \*J. Wildl. Manag.\* 73\(3\), 433-441.](#)
- 4 Bassi, C. (2021). *Etude de la taille efficace et de la diversité génétique de la population d'ours brun,*  
5 *Ursus arctos, des Pyrénées.* Rapport de stage de Master 2<sup>e</sup> année, Université Aix-Marseille,  
6 France.
- 7 Beaumelle, C. (2016). *Evolution génétique et généalogie de la population des ours bruns (Ursus*  
8 *arctos) dans les Pyrénées.* Rapport de stage de Master 1<sup>er</sup> année, Université Grenoble Alpes,  
9 France.
- 10 Beissinger, S.R. & Westphal, I. (1998). On the use of demographic models of population viability in  
11 endangered species management. *J. Wildl. Manag.* 62, 821-841.
- 12 [Bellemain, E.V.A., Swenson, J.E., Tallmon, D., Brunberg, S. & Taberlet, P. \(2005\). Estimating](#)  
13 [population size of elusive animals with DNA from hunter-collected feces: four methods for](#)  
14 [brown bears. \*Cons. Biol.\* 19\(1\), 150-161.](#)
- 15 [Berezowska-Cnota, T., Luque-Márquez, I., Elguero-Claramunt, I., Bojarska, K., Okarma, H. & Selva,](#)  
16 [N. \(2017\). Effectiveness of different types of hair traps for brown bear research and](#)  
17 [monitoring. \*PLoS One\* 12\(10\), e0186605.](#)
- 18 Bischof, R., Brøseth, H. & Gimenez, O. (2016). Wildlife in a politically divided world: Insularism  
19 inflates estimates of brown bear abundance. *Conserv. Lett.* 9, 122-130.
- 20 Blanc, L., Marboutin, E., Gatti, S. & Gimenez, O. (2013). Abundance of rare and elusive species:  
21 empirical investigation of closed versus spatially explicit capture–recapture models with lynx  
22 as a case study. *J. Wildl. Manag.* 77(2), 372-378.
- 23 [Borchers, D. & Fewster, R. \(2016\). Spatial capture–recapture models. \*Statistical Science\* 219-232.](#)

- 1 Boulanger, J., White, G.C., McLellan, B.N., Woods, J., Proctor, M. & Himmer, S. (2002). A meta-  
2 analysis of grizzly bear DNA mark-recapture projects in British Columbia, Canada. *Ursus*  
3 137-152.
- 4 Bunnell, F.L. & Tait, D.E.N. (1985). Mortality rates of North American bears. *Arctic* 38, 316-323
- 5 Burnham, K.P. & Anderson, D.R. (2002). *Model Selection and Multimodel Inference: A Practical*  
6 *Information-Theoretic Approach*. 2nd edition. New York: Springer-Verlag.
- 7 ~~Burton, A.C., Neilson, E., Moreira, D., Ladle, A., Steenweg, R., Fisher, J.T., Bayne, E. & Boutin, S.~~  
8 ~~(2015). Wildlife camera trapping: a review and recommendations for linking surveys to~~  
9 ~~ecological processes. *J. Appl. Eco.* 52(3), 675-685.~~
- 10 Camarra, J.J., Sentilles, J., Bombillon, N., Gastineau, A. & Quenette, P.Y. (2015). *Suivi de l'ours*  
11 *brun dans les Pyrénées françaises (Sous-populations occidentale et centrale) - Rapport*  
12 *annuel, Année 2015*. Vincennes, France: Office National de la Chasse et de la Faune Sauvage.
- 13 Castro-Arellano, I., Madrid-Luna, C., Lacher Jr, T. E. & Léon-Paniagua, L. (2008). Hair-trap efficacy  
14 for detecting mammalian carnivores in the tropics. *J. Wildl. Manag.* 72(6), 1405-1412.
- 15 ~~Chapron, G., Quenette, P.Y., Legendre, S. & Clobert, J. (2003). Which future for the French Pyrenean~~  
16 ~~brown bear (*Ursus arctos*) population? An approach using stage-structured deterministic and~~  
17 ~~stochastic models. *C. R. Biol.* 326, 174-182.~~
- 18 Chapron, G., Wielgus, R., Quenette, P.Y. & Camarra, J.J. (2009). Diagnosing mechanisms of decline  
19 and planning for recovery of an endangered brown bear (*Ursus arctos*) population. *PloS One*  
20 4, e7568.
- 21 ~~Couturier, M.A.J. (1954). *L'ours brun: Ursus arctos* L. (Vol. 1). Grenoble, France: Arthaud.~~
- 22 Creel, S., Spong, G., Sands, J.L., Rotella, J., Zeigle, J., Joe, L., Murphy, K.M. & Smith, D. (2003).  
23 Population size estimation in Yellowstone wolves with error-prone noninvasive microsatellite  
24 genotypes. *Mol. Ecol.* 12(7), 2003-2009.

- 1 Cubaynes, S., Pradel, R., Choquet, R., Duchamp, C., Gaillard, J.M., Lebreton, J.D., Marboutin, E.,  
2 Miquel, C., Reboulet, A.M., Poillot, C., Taberlet, P. & Gimenez, O. (2010). Importance of  
3 accounting for detection heterogeneity when estimating abundance: the case of French  
4 wolves. *Conserv. Biol.* 24(2), 621-626.
- 5 ~~Dahle, B. & Swenson, J.E. (2003). Home ranges in adult Scandinavian brown bears (*Ursus arctos*):~~  
6 ~~effect of mass, sex, reproductive category, population density and habitat type. *J. Zool.* 260(4),~~  
7 ~~329-335.~~
- 8
- 9 Darimont, C.T., Paquet, P.C., Treves, A. & Artelle, K.A. (2018). Political populations of large  
10 carnivores. *Conserv. Biol.* 32(3), 747-749.
- 11 De Barba, M., Miquel, C., Lobréaux, S., Quenette, P.Y., Swenson, J.E. & Taberlet, P. (2017). High-  
12 throughput microsatellite genotyping in ecology: Improved accuracy, efficiency,  
13 standardization and success with low-quantity and degraded DNA. *Mol. Ecol. Res.* 17(3), 492-  
14 507.
- 15 De Barba, M., Waits, L.P., Garton, E.O., Genovesi, P., Randi, E., Mustoni, A. & Groff, C. (2010).  
16 The power of genetic monitoring for studying demography, ecology and genetics of a  
17 reintroduced brown bear population. *Mol. Ecol.* 19, 3938-3951.
- 18 Denes, F.V., Silveira, L.F. & Beissinger, S.R. (2015). Estimating abundance of unmarked animal  
19 populations: accounting for imperfect detection and other sources of zero inflation. *Methods*  
20 *Ecol. Evol.* 6(5), 543-556.
- 21 Fagen, R. & Fagen, J.M. (1996). Individual distinctiveness in brown bears, *Ursus arctos* L. *Ethology*  
22 102, 212-226.
- 23 Forney, K.A. (2000). Environmental models of cetacean abundance: reducing uncertainty in  
24 population trends. *Conserv. Biol.* 14(5) 1271-1286.

- 1 [Forsyth, D.M., Comte, S., Davis, N.E., Bengsen, A.J., Côté, S.D., Hewitt, D.G., Morellet, N. &](#)  
2 [Mysterud, A. \(2022\). Methodology matters when estimating deer abundance: a global](#)  
3 [systematic review and recommendations for improvements. \*J. Wildl. Manag.\* e22207.](#)
- 4 [Frank, S.C., Ordiz, A., Gosselin, J., Hertel, A., Kindberg, J., Leclerc, M., Pelletier, F., Steyaert, S.M.](#)  
5 [J. G., Støen, O.-G., Van de Walle, J., Zedrosser, A. & Swenson, J.E. \(2017\). Indirect effects](#)  
6 [of bear hunting: a review from Scandinavia. \*Ursus\* 28\(2\), 150-164.](#)
- 7 ~~[Frankham, R. \(1995\). Effective population size/adult population size ratios in wildlife: a review.](#)~~  
8 ~~[\*Genet. Res.\* 66\(2\), 95-107.](#)~~
- 9
- 10 Gervasi, V., Brøseth, H., Gimenez, O., Nilsen, E.-B., Odden, J., Flagstad, Ø. & Linnell, J. D. (2016).  
11 Sharing data improves monitoring of trans-boundary populations: the case of wolverines in  
12 central Scandinavia. *Wildl. Biol.* 22(3), 95-106.
- 13 Gervasi, V., Ciucci, P., Davoli, F., Boulanger, J., Boitani, L. & Randi, E. (2010). Addressing  
14 challenges in non invasive capture-recapture based estimates of small populations: a pilot  
15 study on the Apennine brown bear. *Conserv. Genet.* 11(6), 2299-2310.
- 16 [Gavrilov G.V., Zlatanova D.P., Spasova V.V., Valchev K.D. & Dutsov A.A. \(2015\). Home range and](#)  
17 [habitat use of brown bear in Bulgaria: The first data based on GPS-telemetry. \*Acta Zool. Bulg.\*](#)  
18 [67: 493–499.](#)
- 19 Henle, K. & Gruber, B. (2018). Performance of multistate mark–recapture models for temporary  
20 emigration in the presence of survival costs. *Methods Ecol. Evol.* 9(3), 657-667.
- 21 [Huber D. & Roth H.U. \(1993\). Movements of European brown bears in Croatia. \*Acta Theriol.\* 38:](#)  
22 [151–159.](#)
- 23 UICN France, MNHN, SFEPM & ONCFS (2017). *La Liste rouge des espèces menacées en France*  
24 *– Chapitre Mammifères de France métropolitaine.* Paris, France.



- 1 Keating, K.A., Schwartz, C.C., Haroldson, M.A. & Moody, D. (2002). Estimating numbers of females  
2 with cubs-of-the-year in the Yellowstone grizzly bear population. *Ursus* 13, 161-174.
- 3 Kendall, K.C. & McKelvey, K.S. (2008). Hair collection. In *Noninvasive survey methods for North*  
4 *American carnivores: 35-176.* Long, R.A., MacKay, P., Ray, J.C. & Zielinski, W.J. (Eds.).  
5 *Washington, DC: Island Press.*
- 6 Kendall, W.L., Pollock, K.H. & Brownie C. (1995). Likelihood-based approach to capture-recapture  
7 *estimation of demographic parameters under the robust design. Biometrics* 51, 293–308.
- 8 Kendall, W.L., Nichols, J.D. & Hines, J.E. (1997). Estimating temporary emigration using capture–  
9 recapture data with Pollock's robust design. *Ecology* 78, 563-578.
- 10
- 11 Laake, J.L. (2013). *RMark: An R interface for analysis of capture-recapture data with MARK*. AFSC  
12 Processed Rep. 2013-01, 25 p. Alaska Fish. Sci. Cent., NOAA, Natl. Mar. Fish. Serv., 7600  
13 Sand Point Way NE, Seattle WA 98115.
- 14 Lampa, S., Henle, K., Klenke, R., Hoehn, M. & Gruber, B. (2013). How to overcome genotyping  
15 *errors in non-invasive genetic mark-recapture population size estimation—A review of*  
16 *available methods illustrated by a case study. J. Wildl. Manag.* 77(8), 1490-1511.
- 17 Le Maho, Y., Boitani, L., Clobert, J., Quenette, P.Y., Sarrazin, F. & Sauré-Soubelet, A. (2013).  
18 *Expertise collective scientifique « L'Ours brun dans les Pyrénées »*. Paris, France: Muséum  
19 National d'Histoire Naturelle.
- 20 Long, R.A., Zielinski, W.J., Long, R., MacKay, P., Zielinski, W. & Ray, J. (2008). Designing  
21 effective noninvasive carnivore surveys. In *Noninvasive survey methods for carnivores: 8-44.*  
22 Long, R.A., MacKay, P., Ray, J. & Zielinski, W. (Eds.). Washington, USA: Island Press.
- 23 Lukacs, P.M. (2005). *Statistical aspects of using genetic markers for individual identification in*  
24 *capture-recapture studies*. Ph.D. thesis, Colorado State University, USA.

- 1 Lukacs, P.M. & Burnham, K.P. (2005). Review of capture–recapture methods applicable to  
2 noninvasive genetic sampling. *Mol. Ecol.* 14(13), 3909-3919.
- 3 Martin, J., Revilla, E., Quenette, P.Y., Naves, J., Allainé, D. & Swenson, J.E. (2012). Brown bear  
4 habitat suitability in the Pyrenees: transferability across sites and linking scales to make the  
5 most of scarce data. *J. Appl. Ecol.* 49(3), 621-631.
- 6 McCall, B.S., Mitchell, M.S., Schwartz, M.K., Hayden, J., Cushman, S.A., Zager, P. & Kasworm,  
7 W.F. (2013). Combined use of mark-recapture and genetic analyses reveals response of a  
8 black bear population to changes in food productivity. *J. Wildl. Manag.* 77(8), 1572-1582.
- 9 McGowan, C.P., Runge, M.C. & Larson, M.A. (2011). Incorporating parametric uncertainty into  
10 population viability analysis models. *Biol. Conserv.* 144(5), 1400-1408.
- 11 McLellan, B.N. (2015). Some mechanisms underlying variation in vital rates of grizzly bears on a  
12 multiple use landscape. *J. Wildl. Manag.* 79(5), 749-765.
- 13 McLoughlin, P.D., Ferguson, S.H. & Messier, F. (2000). Intraspecific variation in home range  
14 overlap with habitat quality: a comparison among brown bear populations. *Evol. Ecol.* 14(1),  
15 39-60.
- 16 Mertzanis Y., Ioannis I., Mavridis A., Nikolaou O., Riegler S., Riegler A. & Tragos A. (2005).  
17 Movements, activity patterns and home range of a female brown bear (*Ursus arctos*, L.) in  
18 the Rodopi Mountain Range, Greece. *Belgian J. Zool.* 135: 217–221.
- 19 Miotto, R.A., Rodrigues, F.P., Ciocheti, G. & Galetti Jr, P.M. (2007). Determination of the minimum  
20 population size of pumas (*Puma concolor*) through fecal DNA analysis in two protected  
21 Cerrado areas in the Brazilian southeast. *Biotropica* 39(5), 647-654.
- 22 Miquel, C., Bellemain, E., Poillot, C., Bessiere, J., Durand, A. & Taberlet, P. (2006). Quality indexes  
23 to assess the reliability of genotypes in studies using noninvasive sampling and multiple-tube  
24 approach. *Mol. Ecol. Notes* 6(4), 985-988.

- 1 [Morin, D.J., Boulanger, J., Bischof, R., Lee, D.C., Ngoprasert, D., Fuller, A.K., McLellan B.,](#)  
2 [Steinmetz R., Sharma S., Garshelis D.L., Gopalaswamy A.M., Nawaz M.A., & Karanth, U.](#)  
3 [\(2022\). Comparison of methods for estimating density and population trends for low-density](#)  
4 [Asian bears. \*Glob. Ecol. Biogeogr.\* 35, e02058.](#)
- 5 Mowat, G. & Strobeck, C. (2000). Estimating population size of grizzly bears using hair capture,  
6 DNA profiling, and mark-recapture analysis. *J. Wildl. Manag.* 183-193.
- 7 [Murray, D.L. & Sandercock, B.K. \(2020\). \*Population ecology in practice\*. John Wiley & Sons.](#)
- 8 Nichols, J.D. & Williams, B.K. (2006). Monitoring for conservation. *Trends Ecol. Evol.* 21(12), 668-  
9 673.
- 10 Otis, D.L., Burnham, K.P., White, G.C. & Anderson, D.R. (1978). Statistical inference from capture  
11 data on closed animal populations. *Wildl. Monogr.* (62), 3-135.
- 12 ~~[Palomero, G., Ballesteros, F., Nores, C., Blanco, J.C., Herrero, J. & García Serrano, A. \(2007\).](#)~~  
13 ~~[Trends in number and distribution of brown bear females with cubs of the year in the](#)~~  
14 ~~[Cantabrian Mountains, Spain. \*Ursus\* 18\(2\), 145-157.](#)~~ ~~[Parres, A., Palazón, S., Afonso, I.,](#)~~  
15 ~~[Quenette, P.Y., Batet, A., Camarra, J.J., Garreta, X., Gonçalves, S., Guillen, J., Mir, S., Jato,](#)~~  
16 ~~[R., Rodríguez, J., Sentilles, J., Xicola, L. & Melero, Y. \(2020\). Activity patterns in the](#)~~  
17 ~~[reintroduced Pyrenean brown bear population. \*Mamm. Res.\* 65\(3\), 435-444.](#)~~
- 18 [Pederson, J.C., Bunnell, K.D., Conner, M.M. & McLaughlin, C.R. \(2012\). A robust-design analysis](#)  
19 [to estimate American black bear population parameters in Utah. \*Ursus\* 23\(1\), 104-116.](#)
- 20 [Petit, E. & Valiere, N. \(2006\). Estimating population size with noninvasive capture-mark-recapture](#)  
21 [data. \*Cons. Biol.\* 20\(4\), 1062-1073.](#)
- 22 [Piel, A.K., Crunchant, A., Knot, I.E., Chalmers, C., Fergus, P., Mulero-Pázmány, M. & Wich, S.A.](#)  
23 [\(2022\). Noninvasive technologies for primate conservation in the 21st century. \*Int. J.\*](#)  
24 [\*Primatol.\* 43\(1\), 133-167.](#)

- 1 Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs  
2 sampling. In *Proceedings of the 3rd international workshop on distributed statistical*  
3 *computing*: 1-10. Hornik, K., Leisch, F. & Zeileis, A. (Eds.). Vol. 124, No. 125.10. Vienna,  
4 Austria.
- 5 [Pollock, K.H. \(1982\). A capture-recapture design robust to unequal probability of capture. \*J. Wildl.\*](#)  
6 [\*Manag.\* 46\(3\), 752-757.](#)
- 7 [Proctor, M.F., Garshelis, D.L., Thatte, P., Steinmetz, R., Crudge, B., McLellan, B.N., McShea, W.J.,](#)  
8 [Ngoprasert, D., Nawaz, M.A., Wong, S.T., Sharma, S., Fuller, A., Dharaiya, N., Pigeon, K.,](#)  
9 [Hwang, M.-H., Fredriksson, G., Wang, D. & Li, S. \(2022\). Review of field methods for](#)  
10 [monitoring Asian bears. \*Glob. Ecol. Conserv.\* e02080.](#)
- 11 Quenette, P.Y., Jonozovic, M., & Marinsic, A., Camarra, J.J., Guinot-Ghestem, M., Sentilles, J.,  
12 Steinmetzn J., Vanpé, C. & Alban, N. (2019). First translocation of females in the French  
13 Western Pyrenees: a new step in the long process of brown bear restoration in the Pyrenees  
14 Mountains. *Int. Bear News* 28, 11-13.
- 15 [Quinn, T.P., Wirsing, A.J. & Proctor, M. \(2022\). Optimal barbed wire height for brown bear hair](#)  
16 [sample collection. \*Ursus\* 2022\(33e2\), 1-6.](#)
- 17 Rankin, R.W., Nicholson, K.E., Allen, S.J., Krützen, M., Bejder, L. & Pollock, K.H. (2016). A full-  
18 capture hierarchical Bayesian model of Pollock's Closed Robust Design and application to  
19 dolphins. *Front. Mar. Sci.* 3, 25.
- 20 RCoreTeam (2013). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R  
21 Foundation for Statistical Computing.
- 22 Riecke, T.V., Leach, A.G., Gibson, D. & Sedinger, J.S. (2018). Parameterizing the robust design in  
23 the BUGS language: Lifetime carry-over effects of environmental conditions during growth  
24 on a long-lived bird. *Methods Ecol. Evol.* 9(11), 2294-2305.

- 1 [Royle, J.A., Chandler, R.B., Sollmann, R. & Gardner, B. \(2013\). \*Spatial capture-recapture\*.  
2 \*Academic Press\*.](#)
- 3 Schofield, M.R. & Barker, R.J. (2011). Full open population capture–recapture models with  
4 individual covariates. *J. Agric. Biol. Environ. Stat.* 16(2), 253-268.
- 5 ~~[Schwartz, C.C., Keating, K.A., Reynolds III, H.V., Barnes Jr, V.G., Sellers, R.A., Swenson, J.E.,  
6 Miller, S.D., McLellan, B.N., Keay, J., McCann, R., Gibeau, M., Wakkenin, W.F., Mace, R.  
7 D., Kasworm, W. & Smith, R. \(2003\). Reproductive maturation and senescence in the female  
8 brown bear. \*Ursus\* 109-119.](#)~~
- 9
- 10 Sentilles J., Lemaître P.L., Vanpé C. & Quenette P.Y. (~~2021a~~2022). *Rapport annuel du réseau ours  
11 brun 20202021*. Paris, France: Office Français de la Biodiversité.
- 12 Sentilles, J., Vanpé, C. & Quenette, P.Y. (2021~~b~~). Benefits of incorporating a scat-detection dog into  
13 wildlife monitoring: a case study of Pyrenean brown bear. *J. Vertebr. Biol.* 69(3), 20096-1.
- 14 [Sergiel, A., Barja, I., Navarro-Castilla, Á., Zwijacz-Kozica, T. & Selva, N. \(2020\). Losing seasonal  
15 patterns in a hibernating omnivore? Diet quality proxies and faecal cortisol metabolites in  
16 brown bears in areas with and without artificial feeding. \*PLoS One\* 15\(11\), e0242341.](#)
- 17 Solberg, K.H., Bellemain, E., Drageset, O.M., Taberlet, P. & Swenson, J.E. (2006). An evaluation of  
18 field and non-invasive genetic methods to estimate brown bear (*Ursus arctos*) population size.  
19 *Biol. Conserv.* 128(2), 158-168.
- 20 [Spady, T.J., Lindburg, D.G. & Durrant, B.S. \(2007\). Evolution of reproductive seasonality in bears.  
21 \*Mammal Review\* Volume 37\(1\), 21–53.](#)
- 22 [Staton, B.A., Justice, C., White, S., Sedell, E.R., Burns, L.A. & Kaylor, M.J. \(2022\). Accounting for  
23 uncertainty when estimating drivers of imperfect detection: An integrated approach illustrated  
24 with snorkel surveys for riverine fishes. \*Fish. Res.\* 249, 106209.](#)

- 1 Stoen, O.G., Bellemain, E., Sæbø, S. & Swenson, J.E. (2005). Kin related spatial structure in brown  
2 bears *Ursus arctos*. *Behav. Ecol. Sociobiol.* 59(2), 191-197.
- 3 Stetz, J. B., Kendall, K. C., & Servheen, C. (2010). Evaluation of bear rub surveys to monitor grizzly  
4 bear population trends. *J. Wildl. Manag.* 74(4), 860-870.Swenson, J.E., Adamič, M., Huber,  
5 D. & Stokke, S. (2007). Brown bear body mass and growth in northern and southern Europe.  
6 *Oecologia* 153(1), 37-47.
- 7
- 8 Swenson, J.E., Sandegren, F., Brunberg, S., Segerström, P. & Segerström, P. (2001). Factors  
9 associated with loss of brown bear cubs in Sweden. *Ursus* 69-80.
- 10 Swenson, J.E., Sandegren, F., Söderberg, A., Bjärvall, A., Franzén, R. & Wabakken, P. (1997).  
11 Infanticide caused by hunting of male bears. *Nature* 386(6624), 450-451.
- 12 Swenson, J.E., Gerstl, N., Dahle, B. & Zedrosser, A. (2000). Action plan for the conservation of the  
13 brown bear (*Ursus arctos*) in Europe. *Nat. Environ.* 114, 1-69.
- 14 Taberlet, P., Camarra, J.J., Griffin, S., Uhres, E., Hanotte, O., Waits, L.P., Dubois-Paganon, C.,  
15 Burke, T. & Bouvet, J. (1997). Noninvasive genetic tracking of the endangered Pyrenean  
16 brown bear population. *Mol. Ecol.* 6(9), 869-876.
- 17 Taberlet, P., Griffin, S., Goossens, B., Questiau, S., Manceau, V., Escaravage, N., Waits, L.P. &  
18 Bouvet, J. (1996). Reliable genotyping of samples with very low DNA quantities using PCR.  
19 *Nucleic Acids Res.* 24(16), 3189-3194.
- 20 Thompson, W. (2013). *Sampling rare or elusive species: concepts, designs, and techniques for*  
21 *estimating population parameters*. Washington, D.C., USA: Island Press.
- 22 Tosoni, E., Boitani, L., Gentile, L., Gervasi, V., Latini, R. & Ciucci, P. (2017). Assessment of key  
23 reproductive traits in the Apennine brown bear population. *Ursus* 28(1), 105-116.

- 1 Tourani, M. (2022). A review of spatial capture–recapture: Ecological insights, limitations, and  
2 prospects. *Ecol. Evol.* 12(1), e8468.
- 3 Vanpé, C., Sentilles, J., Lemaître, P.L., Desbarax, R., Bibal, D., Darmani, G., Labal, P. & Quenette,  
4 P.Y. (2021). *Cartographie quinquennale (2016-2020) de l'aire de répartition de l'ours brun*  
5 *dans les Pyrénées françaises*. Vincennes, France: Office Français de la Biodiversité.
- 6 Waits, L.P. & Paetkau, D. (2005). Noninvasive genetic sampling tools for wildlife biologists: a review  
7 of applications and recommendations for accurate data collection. *J. Wildl. Manag.* 69(4),  
8 1419-1433.
- 9 White, G.C. & Burnham, K.P. (1999). Program MARK: survival estimation from populations of  
10 marked animals. *Bird Study* 46(sup1), S120-S139.
- 11 Williams, B.K., Nichols, J.D. & Conroy, M.J. (2002). *Analysis and management of animal*  
12 *populations*. Academic press.
- 13 Wilson, G.J. & Delahay, R.J. (2001). A review of methods to estimate the abundance of terrestrial  
14 carnivores using field signs and observation. *Wildl. Res.* 28(2), 151-164.
- 15 Woods, J.G., Paetkau, D., Lewis, D., McLellan, B.N., Proctor, M. & Strobeck, C. (1999). Genetic  
16 tagging of free-ranging black and brown bears. *Wildl. Soc. Bull.* 616-627.
- 17 ~~Wroe, S. & Milne, N. (2007). Convergence and remarkably consistent constraint in the evolution of~~  
18 ~~carnivore skull shape. *Evol. Int. J. Org. Evol.* 61(5), 1251-1260.~~

1 **Figure captions**

2


3 **Figure 1.** Map of the transboundary range area (on squares of 10 x 10 km) of the Pyrenean brown  
4 bear population for the year 2020.

5

6 **Figure 2.** Variation in the annual population size of the Pyrenean brown bear from 2008 to 2020,  
7 estimated from ~~the~~ Bayesian Pollock's robust design capture-recapture approach (PCRD, black full  
8 circles and black full line, with the associated 97.5% credible interval in grey), compared to the  
9 Minimum Retained population Size (MRS, grey open squares and grey full line) and Minimum  
10 Detected population Size (MDS, black open circles and black dashed line) values.


11 Note: MRS estimate for 2020 is provisional and probably slightly underestimated.






1

2 **Figure 1.**



1



2

3 **Figure 2.**

4 **Table 1.** Model selection from the frequentist capture-recapture approach using a-Pollock's robust design (PCRD) capture-recapture (CR) modelling  
5 approach.

Model	Survival structure	Detection structure	Emigration structure	AICc	$\Delta AICc$
1	Age-dependent	Heterogeneous	Random	1496.43	0.00
2	Age-dependent	Heterogeneous	Markovian	1496.90	0.47
3	Constant	Heterogeneous	Random	1503.48	7.04
4	Constant	Heterogeneous	Markovian	1503.76	7.33
5	Age-dependent	Heterogeneous	No	1520.68	24.25
6	Constant	Heterogeneous	No	1528.73	32.30
7	Age-dependent	Time-dependent (within primary occasions)	Random	1548.00	51.57
8	Age-dependent	Time-dependent (between primary occasions)	Random	1548.28	51.85
9	Age-dependent	Time-dependent (between primary occasions)	Markovian	1548.58	52.14
10	Age-dependent	Time-dependent (within primary occasions)	Markovian	1549.03	52.60
11	Constant	Time-dependent (within primary occasions)	Random	1555.00	58.56
12	Constant	Time-dependent (between primary occasions)	Random	1555.31	58.87
13	Constant	Time-dependent (between primary occasions)	Markovian	1555.41	58.98
14	Constant	Time-dependent (within primary occasions)	Markovian	1555.85	59.42
15	Age-dependent	Constant	Random	1562.58	66.15
16	Age-dependent	Constant	Markovian	1562.86	66.43
17	Constant	Constant	Random	1569.64	73.21
18	Constant	Constant	Markovian	1569.73	73.30
19	Age-dependent	Time-dependent (within primary occasions)	No	1611.84	115.40
20	Constant	Time-dependent (within primary occasions)	No	1619.97	123.54
21	Age-dependent	Time-dependent (between primary occasions)	No	1625.67	129.23
22	Constant	Time-dependent (between primary occasions)	No	1634.10	137.66
23	Age-dependent	Constant	No	1637.98	141.54
24	Constant	Constant	No	1646.44	150.01

7 **Table 2.** Parameter estimates (estimates ± SE) for the two best-supported models from the frequentist capture-recapture (CR) approach (see models 1  
 8 and 2 from Table 1) using a robust design, in which temporary emigration is either random (first column) or Markovian (second column).

9

	<u>Random temporary emigration</u>	<u>Markovian temporary emigration</u>
-		
<u>Cub survival probability</u>	<u>0.844 ± 0.038</u>	<u>0.844 ± 0.038</u>
<u>Subadult survival probability</u>	<u>0.954 ± 0.028</u>	<u>0.955 ± 0.029</u>
<u>Adult survival probability</u>	<u>0.962 ± 0.015</u>	<u>0.962 ± 0.015</u>
<u>Probability of leaving the study area</u>	<u>0.105 ± 0.023</u>	-
<u>Probability of leaving the study area given presence in the study area at the previous sampling occasion</u>	=	<u>0.097 ± 0.023</u>
<u>Probability of leaving the study area given absence in the study area at the previous sampling occasion</u>	=	<u>0.217 ± 0.103</u>
<u>Proportion of individuals in class 1 of mixture</u>	<u>0.722 ± 0.053</u>	<u>0.723 ± 0.053</u>
<u>Detection probability of class 1 individuals</u>	<u>0.421 ± 0.023</u>	<u>0.422 ± 0.023</u>
<u>Detection probability of class 2 individuals</u>	<u>0.850 ± 0.034</u>	<u>0.850 ± 0.034</u>

10

	<b>Temporary emigration</b>			
	<b>Random</b>		<b>Markovian</b>	
	<b>Estimate</b>	<b>SE</b>	<b>Estimate</b>	<b>SE</b>
Cub survival probability	0.844	0.038	0.844	0.038
Subadult survival probability	0.954	0.028	0.955	0.029
Adult survival probability	0.962	0.015	0.962	0.015
Probability of leaving the study area	0.105	0.023	-	-
Probability of leaving the study area given presence in the study area at the previous sampling occasion	-	-	0.097	0.023
Probability of leaving the study area given absence in the study area at the previous sampling occasion	-	-	0.217	0.103
Proportion of individuals in class 1 of mixture	0.722	0.053	0.723	0.053
Detection probability of class 1 individuals	0.421	0.023	0.422	0.023
Detection probability of class 2 individuals	0.850	0.034	0.850	0.034

## 12 **Supplementary Information**

### 13 *Genetic analyses from 2017 to 2020*

14 From 2017 to 2020, genetic analyses were conducted in our laboratory at ANTAGENE  
15 (<https://www.antagene.com/en>). DNA extraction was conducted according to a sterile process in a  
16 designated extraction room free of DNA. For each sample, disposable sterile tools were used and the  
17 bench was cleaned with bleach to avoid cross-contamination. Each sample was transferred to a sterile  
18 labelled microtube to proceed to DNA extraction. Sample tubes were surrounded by positive and  
19 negative extraction controls and lysed overnight at 56°C according to the manufacturer's instructions  
20 (Nucleospin 96 Tissue Kit, Macherey-Nagel, Düren, Germany). DNA was isolated and purified using  
21 purification columns and vacuum filtration (Nucleospin 96 Tissue Kit, Macherey-Nagel, Düren,  
22 Germany). DNA was eluted with 100 µL of elution buffer to obtain final concentrations between 20-  
23 100 ng/µl. Extracts were stored in labelled 96-tube strips plates in a -20°C freezer.

24 For each DNA sample, 13 microsatellites and 3 sex identification markers (ZFX, 318.2 and  
25 SMCY) were amplified by two multiplex PCRs (polymerase chain reaction) four times and analyzed  
26 in two runs (one for each multiplex) with an automated sequencer (Table S5S6). Because the genetic  
27 sex marker described in the scientific publication De Barba et al. (2017) proved to be not very  
28 reproducible, the ANTAGENE laboratory uses a system of three pairs of primers allowing the  
29 amplification by PCR of two specific regions of the Y chromosome and one specific region of the X  
30 chromosome, according to a method developed and validated in all bear species (Bidon 2013). This  
31 system provides an internal positive control for all individuals, with the amplification of a region of  
32 the X chromosome present in males (XY) and in females (XX) and to amplify in duplicate a specific  
33 region of the Y chromosome present only in males (XY). This triple amplification guarantees an  
34 excellent recognition of the Y chromosome and therefore of males, and increases the reliability of  
35 characterization of the genetic sex, especially on DNA from degraded samples (hair, scats, etc.).

36 PCR reactions were prepared step-by-step according to a unidirectional workflow starting in a  
37 clean room with positive air pressure to prepare sensitive reagents (enzymes and DNA primers) and

38 ~~continuing-continued~~ in a pre-PCR room for combining DNA and reagents using filtered tips. Three  
39 negative and positive controls were included per PCR reaction. PCR amplifications were then  
40 performed in a dedicated post-PCR area in 96-well microplates at 10 µl final volumes containing 5  
41 µl of mastermix Taq Polymerase (Type-It Microsatellite PCR Kit, Qiagen, Hilden, Germany), and  
42 either 0.80 µL of a first pool of 8 pairs of primers or 0.36 µl of a second pool of 8 pairs of primers at  
43 a concentration from 0.08 to 0.60 µM each, and a mean of 30 ng of genomic DNA (Table S5S6).  
44 Each pair of primers was coupled with a fluorescent dye (Table S5S6). Our PCR thermal protocol  
45 consisted of 95°C for 15 min, followed by 8 touchdown cycles of 95°C for 30 s, 62°C to 55°C for 90  
46 s (decreasing 1°C per cycle), and 72°C for 30 s, then followed by 35 cycles of 95°C for 30 s, 55°C  
47 for 90 s, and 72°C for 30 s, ending with an extension of 60°C for 30 min. PCR products were resolved  
48 on an ABI PRISM 3130 XL capillary sequencer (ThermoFisher Scientific, Waltham, Massachusetts)  
49 under denaturing conditions (Hi-Di™ Formamide, ThermoFisher Scientific, Waltham,  
50 Massachusetts) with an internal size marker prepared once and dispatched equally in all sample wells  
51 of each multiplex run. The four electropherograms for each sample were analyzed using  
52 GENEMAPPER 4.1 (ThermoFisher Scientific, Waltham, Massachusetts) and analyzed  
53 independently by two analysts to determine the allele sizes for each marker of each individual. When  
54 the genotypes determined by each analyst did not agree, the electropherograms were read again,  
55 reading errors were resolved, and in case of persistent disagreement, ambiguous results were  
56 considered as missing data.

57

### 58 *Dating of bear signs*

59 For photos and videos, we used the metadata from the automatically triggered camera traps or  
60 cameras to define accurately the date of bear presence. For hair collected on baited hair traps, we used  
61 photo data collected on camera traps set up in front of baited hair traps when available to identify date  
62 when hair were left. From those specific bear signs, month of bear presence could be determined  
63 accurately based on the date when signs were left.

64 For other types of bear signs, we could not know precisely the date when signs were left and we  
65 relied on an evaluation of the time period when sign could have been left by the bear. More  
66 specifically, when hair collected on baited hair trap were not associated with any photo or video, we  
67 considered that the bear had left the hair during the time period included between the date of the last  
68 visit of the hair trap when barbed wire was cleaned and the date of the visit when hair were collected.  
69 If this time period was larger than 2 months, we discarded the hair sample from our analyses. We also  
70 discarded hair samples collected spontaneously outside systematic monitoring design, because the  
71 time interval during which they might have been left by the bear could not be evaluated precisely  
72 (bear hair deteriorates very slowly in the field), except in the case hair were associated with a damage  
73 ~~on~~ to livestock or beehives, in which case the estimated date of the damage provided the estimated  
74 date of hair deposition. Finally, we estimated the time interval when scats were dropped (≤2 weeks)  
75 by evaluating the freshness of the scat when collected in the field, using expert judgement in relation  
76 to the color and appearance of the scat, recent weather conditions (rain, sunshine, snow, temperature,  
77 etc.) and type of habitat (directly exposed to sun, under vegetation cover, etc.) (e.g., Sergiel et al.  
78 2020 for a similar approach). When the time period during which hair or scat could have been left  
79 overlapped two different months, we considered as a proxy the month of the median date between  
80 maximum and minimum date of the time period as the month of bear presence, since this should not  
81 affect much our estimation of population size with capture-recapture analyses. Note that we selected  
82 preferentially fresher scats (with less DNA degradation) to send to the molecular laboratory, allowing  
83 a better genotyping success and identifying more individuals genetically (Sentilles, Vanpé &  
84 Quenette 2021). In France, we collected in total 4,022 hair or scat samples from 2008 to 2020, among  
85 which about 5.5% were excluded from our analyses due to inaccurate dating.

86

### 87 *Compilation of monthly detection history of bears*

88 Matching genotypes were considered to arise from the same individual and classified as recaptures  
89 as the combined non-exclusion probability of the 13 microsatellites for independent individuals and



90 for sibships were negligible (Lukacs & Burnham 2005). Importantly, we did not consider location  
91 data from GPS collar or VHF transmitters to compile detection history to avoid large inter-individual  
92 differences in monitoring pressure between bears, since it concerns respectively 5 bears and 1 bear  
93 for a period ranging from several months to a few years. Orphan cubs that were captured in the field  
94 and kept in captivity for a while for care before being released in the wild were considered as still  
95 present and detected in the population during the months of captivity (this concerns only 1 orphan  
96 cub during two months of captivity $N=1$ ). For individuals for which we knew the date of death ( $N =$   
97 9), we used this information and right censored them in the corresponding detection histories. For  
98 translocated bears originating from Slovenia ( $N = 3$ ), the first month of potential detection was the  
99 month of release in the Pyrenees.

100

## 101 References

- 102 Bidon, T., Frosch, C., Eiken, H.G., Kutschera, V.E., Hagen, S.B., Aarnes, S.G., ... & Hailer, F. (2013).  
103 A sensitive and specific multiplex PCR approach for sex identification of ursine and tremarctine bears  
104 suitable for non-invasive samples. *Mol. Ecol. Res.* 13(3), 362-368.
- 105 De Barba, M., Miquel, C., Lobréaux, S., Quenette, P. Y., Swenson, J. E., & Taberlet, P. (2017). High-  
106 throughput microsatellite genotyping in ecology: Improved accuracy, efficiency, standardization and  
107 success with low-quantity and degraded DNA. *Mol. Ecol. Res.* 17(3), 492-507.

108 **Table S1.** Systematic monitoring effort in the French Pyrenees in terms of number of transects (including 6 hair traps per transect in average), total length  
 109 of transects (km), number of camera traps, number of baited hair traps ~~and number of genetically analysed samples~~ per year between 2008 and 2020.  
 110

<u>Year</u>	<u># transects</u>	<u>total length of transects (km)</u>	<u># camera traps</u>	<u># baited hair traps</u>
<u>2008</u>	<u>30</u>	<u>300*</u>	<u>7</u>	<u>59</u>
<u>2009</u>	<u>36</u>	<u>360*</u>	<u>10</u>	<u>73</u>
<u>2010</u>	<u>60</u>	<u>600*</u>	<u>12</u>	<u>86</u>
<u>2011</u>	<u>68</u>	<u>615</u>	<u>26</u>	<u>90</u>
<u>2012</u>	<u>68</u>	<u>615</u>	<u>18</u>	<u>0</u>
<u>2013</u>	<u>48</u>	<u>426</u>	<u>49</u>	<u>0</u>
<u>2014</u>	<u>50</u>	<u>411</u>	<u>39</u>	<u>0</u>
<u>2015</u>	<u>44</u>	<u>358</u>	<u>40</u>	<u>0</u>
<u>2016</u>	<u>47</u>	<u>376</u>	<u>48</u>	<u>0</u>
<u>2017</u>	<u>53</u>	<u>414</u>	<u>45</u>	<u>0</u>
<u>2018</u>	<u>57</u>	<u>441</u>	<u>45</u>	<u>0</u>
<u>2019</u>	<u>56</u>	<u>424</u>	<u>59</u>	<u>0</u>
<u>2020</u>	<u>58</u>	<u>428</u>	<u>60</u>	<u>0</u>

<u>Year</u>	<u># transects</u>	<u>total length of transects (km)</u>	<u># camera traps</u>	<u># baited hair traps</u>	<u># analysed samples</u>
<u>2008</u>	<u>30</u>	<u>300*</u>	<u>7</u>	<u>59</u>	<u>125</u>
<u>2009</u>	<u>36</u>	<u>360*</u>	<u>10</u>	<u>73</u>	<u>84</u>
<u>2010</u>	<u>60</u>	<u>600*</u>	<u>12</u>	<u>86</u>	<u>167</u>
<u>2011</u>	<u>68</u>	<u>615</u>	<u>26</u>	<u>90</u>	<u>209</u>
<u>2012</u>	<u>68</u>	<u>615</u>	<u>18</u>	<u>0</u>	<u>224</u>

2013	48	426	49	0	137
2014	50	411	39	0	193
2015	44	358	40	0	152
2016	47	376	48	0	179
2017	53	414	45	0	134
2018	57	441	45	0	158
2019	56	424	59	0	314
2020	58	428	60	0	448

---

112

113 Note: \* Estimated based on an average transect length of 10 km. ~~The number of analysed samples corresponds to the number of scat or hair samples~~  
 114 ~~(collected in France, Spain or Andorra) analysed by the French molecular laboratory (LECA or our laboratory) per year.~~

115

116 **Table S2.** Total number of validated non-invasive brown bear signs (e.g., scats, hair, tracks, visual observations, damages, photos / videos) collected in  
 117 the Pyrenees, total number of validated brown bear samples (i.e. scats and hair) collected in the Pyrenees, number of samples (among collected sampled)  
 118 genetically analysed by the French molecular laboratory LECA or ANTAGENE, number of brown bear samples (among analysed samples) successfully  
 119 genotyped and number of different brown bear genotypes identified (among successfully genotyped samples) per year between 2008 and 2020.

<u>Year</u>	<u># validated bear signs collected</u>	<u># bear samples</u>	<u># genetically analysed samples</u>	<u># successfully genotyped samples</u>	<u># different genotypes identified</u>
<u>2008</u>	<u>743</u>	<u>210</u>	<u>125</u>	<u>73</u>	<u>11</u>
<u>2009</u>	<u>712</u>	<u>229</u>	<u>84</u>	<u>42</u>	<u>12</u>
<u>2010</u>	<u>939</u>	<u>323</u>	<u>167</u>	<u>106</u>	<u>15</u>
<u>2011</u>	<u>1152</u>	<u>518</u>	<u>209</u>	<u>122</u>	<u>15</u>
<u>2012</u>	<u>1239</u>	<u>521</u>	<u>224</u>	<u>153</u>	<u>15</u>
<u>2013</u>	<u>1318</u>	<u>521</u>	<u>137</u>	<u>77</u>	<u>14</u>
<u>2014</u>	<u>1243</u>	<u>571</u>	<u>193</u>	<u>96</u>	<u>21</u>
<u>2015</u>	<u>1567</u>	<u>870</u>	<u>152</u>	<u>110</u>	<u>24</u>
<u>2016</u>	<u>1854</u>	<u>874</u>	<u>179</u>	<u>137</u>	<u>32</u>
<u>2017</u>	<u>1394</u>	<u>569</u>	<u>134</u>	<u>105</u>	<u>34</u>
<u>2018</u>	<u>1625</u>	<u>601</u>	<u>158</u>	<u>109</u>	<u>35</u>
<u>2019</u>	<u>2450</u>	<u>830</u>	<u>314</u>	<u>209</u>	<u>41</u>
<u>2020</u>	<u>2783</u>	<u>1116</u>	<u>448</u>	<u>309</u>	<u>45</u>
<u>TOTAL</u>	<u>19019</u>	<u>7753</u>	<u>2524</u>	<u>1648</u>	<u>314</u>

121

122

123

124 **Table S3.** Combination of microsatellite markers used in each PCR mix and type of fluorescent dye  
 125 used for each microsatellite marker from 2017 to 2020.

126

<b>Mix</b>	<b>Locus name</b>	<b>Dye</b>	<b>Publication</b>
<u>A</u>	<u>UA03</u>	<u>6FAM</u>	<u>De Barba et al. 2017</u>
<u>A</u>	<u>UA06</u>	<u>6FAM</u>	<u>De Barba et al. 2017</u>
<u>A</u>	<u>UA25</u>	<u>NED™</u>	<u>De Barba et al. 2017</u>
<u>A</u>	<u>UA67</u>	<u>NED™</u>	<u>De Barba et al. 2017</u>
<u>A</u>	<u>UA64</u>	<u>PET™</u>	<u>De Barba et al. 2017</u>
<u>A</u>	<u>UA63</u>	<u>PET™</u>	<u>De Barba et al. 2017</u>
<u>A</u>	<u>UA16</u>	<u>VIC™</u>	<u>De Barba et al. 2017</u>
<u>A</u>	<u>UA14</u>	<u>VIC™</u>	<u>De Barba et al. 2017</u>
<u>B</u>	<u>UA17</u>	<u>6FAM</u>	<u>De Barba et al. 2017</u>
<u>B</u>	<u>UA57</u>	<u>6FAM</u>	<u>De Barba et al. 2017</u>
<u>B</u>	<u>UA51</u>	<u>NED™</u>	<u>De Barba et al. 2017</u>
<u>B</u>	<u>UA65</u>	<u>PET™</u>	<u>De Barba et al. 2017</u>
<u>B</u>	<u>UA68</u>	<u>VIC™</u>	<u>De Barba et al. 2017</u>
<u>B</u>	<u>Our-ZFX</u>	<u>6FAM</u>	<u>Bidon et al. 2013</u>
<u>B</u>	<u>Our-318</u>	<u>6FAM</u>	<u>Bidon et al. 2013</u>
<u>B</u>	<u>Our-SMCY</u>	<u>6FAM</u>	<u>Bidon et al. 2013</u>

127

128 References:

129 Bidon, T., Frosch, C., Eiken, H. G., Kutschera, V. E., Hagen, S. B., Aarnes, S. G., Fain, S.R., Janke,  
 130 A. & Hailer, F. (2013). A sensitive and specific multiplex PCR approach for sex identification of  
 131 ursine and tremarctine bears suitable for non-invasive samples. *Mol.Ecol. Res.* 13(3), 362-368.

132 De Barba, M., Miquel, C., Lobréaux, S., Quenette, P. Y., Swenson, J. E., & Taberlet, P. (2017). High-  
 133 throughput microsatellite genotyping in ecology: Improved accuracy, efficiency, standardization and  
 134 success with low-quantity and degraded DNA. *Mol.Ecol. Res.* 17(3), 492-507.

135

136 **Table S2S4.** Summary statistics of the 58 different genotypes found in the Pyrenean brown bear  
 137 population in 2020 for each of the 13 microsatellite loci provided by the allele frequency analysis of  
 138 CERVUS software (Marshall et al. 1998).

Locus	N	k	HObs	HExp	NE-I	NE-SI	PIC	F(Null)
UA03	58	4	0.655	0.622	0.215	0.495	0.549	-0.0218
UA06	58	4	0.724	0.657	0.173	0.467	0.6	-0.0653
UA14	58	4	0.759	0.705	0.144	0.437	0.645	-0.0458
UA16	58	6	0.414	0.461	0.328	0.604	0.424	0.0908
UA17	58	3	0.517	0.497	0.308	0.581	0.442	-0.0189
UA25	58	5	0.483	0.427	0.364	0.629	0.392	-0.0989
UA51	58	4	0.603	0.537	0.269	0.551	0.483	-0.0608
UA57	58	3	0.552	0.45	0.399	0.627	0.354	-0.1089
UA63	57	6	0.719	0.694	0.146	0.442	0.639	-0.0178
UA64	58	2	0.534	0.492	0.381	0.601	0.369	-0.0455
UA65	58	4	0.621	0.595	0.246	0.516	0.513	-0.0358
UA67	58	3	0.517	0.571	0.266	0.533	0.488	0.0589
UA68	58	5	0.724	0.734	0.121	0.417	0.68	-0.0054
MEAN		4.08	0.602	0.572	0.258	0.531	0.506	-0.0289

139 Note: N: number of individuals typed, k: the number of alleles, Hobs: observed heterozygosity, Hexp:  
 140 expected heterozygosity, NE-I: average exclusion probabilities for each locus for identity, NE-SI:  
 141 average exclusion probabilities for each locus for sib identity, PIC: polymorphic information content,  
 142 F(Null): the frequency of null alleles. The combined non-exclusion probabilities for identity and sib  
 143 identity were  $9.10^{-9}$  and 0.000235, respectively.

144  
 145 Reference:

146 Marshall, T.C., Slate, J.B.K.E., Kruuk, L.E.B. & Pemberton, J.M. (1998). Statistical confidence for  
 147 likelihood-based paternity inference in natural populations. *Mol. Ecol.* 7(5), 639-655.

148 **Table S3S5.** Parameters of the model, in which temporary emigration is random, survival is age-  
 149 dependent ~~survival~~ and there is heterogeneity in the detection process, estimated using a Bayesian  
 150 robust-design capture-recapture (CR) approach.

	Mean	SD	Naive SE	Time-series SE
beta[1]	8.52E-01	0.0402	0.8991	0.0014
beta[2]	9.33E-01	0.0333	0.0007	0.0009
beta[3]	9.46E-01	0.0175	0.0004	0.0004
deviance	2.31E+03	17.9349	0.4010	0.5163
gamma	6.62E-02	0.0218	0.0005	0.0006
mean.p	4.29E-01	0.0272	0.0006	0.0019
pstar[1]	8.50E-01	0.0188	0.0004	0.0009
pstar[2]	8.50E-01	0.0188	0.0004	0.0009
pstar[3]	8.50E-01	0.0188	0.0004	0.0009
pstar[4]	8.50E-01	0.0188	0.0004	0.0009
pstar[5]	8.50E-01	0.0188	0.0004	0.0009
pstar[6]	8.50E-01	0.0188	0.0004	0.0009
pstar[7]	8.50E-01	0.0188	0.0004	0.0009
pstar[8]	8.50E-01	0.0188	0.0004	0.0009
pstar[9]	8.50E-01	0.0188	0.0004	0.0009
pstar[10]	8.50E-01	0.0188	0.0004	0.0009
pstar[11]	8.50E-01	0.0188	0.0004	0.0009
pstar[12]	8.50E-01	0.0188	0.0004	0.0009
sdeps	2.10E-01	0.0214	0.0005	0.0007

151 Note: beta[i]: age-specific survival for age i (with 1: cubs, 2: subadults, 3: adults), gamma: probability  
 152 of emigration; mean.p: mean detection probability, sdeps: SD of the random effect, pstar[j]: averaged  
 153 detection over individuals for year j, with j ranging from 2008 to 2019.

154 **Table S4S6.** Comparison of the annual abundance of the Pyrenean brown bear population, estimated  
 155 from a Bayesian Pollock's robust design (PCRD) capture-recapture (CR) approach (with associated  
 156 97.5% Credible Interval), with Minimum Detected Size (MDS, total number of different individuals  
 157 detected in the population during the year) and Minimum Retained Size (MRS, reassessment of the  
 158 MDS in the light of the information newly collected in the following subsequent years) values from  
 159 2008 to 2020.

160

Year	PCRD Estimate	97.5% CI	MDS value	MRS value
2008	13.0	12.8 - 13.3	16	15
2009	17.4	17.0 - 17.8	17	16
2010	16.3	15.9 - 16.7	18	20
2011	19.5	19.1 - 20.0	22	23
2012	23.9	23.4 - 24.4	22	24
2013	21.7	21.3 - 22.2	25	25
2014	26.0	25.5 - 26.7	31	31
2015	29.3	28.7 - 30.0	29	32
2016	41.2	40.4 - 42.2	39	41
2017	41.2	40.4 - 42.2	43	48
2018	39.1	38.3 - 40.0	40	<u>5152</u>
2019	49.9	48.9 - 51.1	52	<u>5859</u>
2020	66.2	64.8 - 67.8	64	<u>NA68</u>

161

162 Note: MRS count for 2020 is provisional and probably slightly underestimated.

163



164 **Table S7.** Evolution of the sex ratio of the Pyrenean brown bear population from 2008 to 2020  
 165 among all individuals and among adult only.

166

	<u># adult males</u>	<u># adult females</u>	<u>Adult sex ratio</u>	<u># males</u>	<u># females</u>	<u>Sex ratio</u>
<u>2008</u>	<u>5</u>	<u>4</u>	<u>1.25</u>	<u>8</u>	<u>6</u>	<u>1.33</u>
<u>2009</u>	<u>6</u>	<u>5</u>	<u>1.20</u>	<u>8</u>	<u>8</u>	<u>1.00</u>
<u>2010</u>	<u>5</u>	<u>6</u>	<u>0.83</u>	<u>8</u>	<u>12</u>	<u>0.67</u>
<u>2011</u>	<u>6</u>	<u>6</u>	<u>1.00</u>	<u>8</u>	<u>14</u>	<u>0.57</u>
<u>2012</u>	<u>6</u>	<u>7</u>	<u>0.86</u>	<u>9</u>	<u>14</u>	<u>0.64</u>
<u>2013</u>	<u>6</u>	<u>9</u>	<u>0.67</u>	<u>9</u>	<u>14</u>	<u>0.64</u>
<u>2014</u>	<u>8</u>	<u>11</u>	<u>0.73</u>	<u>11</u>	<u>17</u>	<u>0.65</u>
<u>2015</u>	<u>7</u>	<u>12</u>	<u>0.58</u>	<u>13</u>	<u>19</u>	<u>0.68</u>
<u>2016</u>	<u>7</u>	<u>13</u>	<u>0.54</u>	<u>19</u>	<u>21</u>	<u>0.90</u>
<u>2017</u>	<u>8</u>	<u>14</u>	<u>0.57</u>	<u>21</u>	<u>26</u>	<u>0.81</u>
<u>2018</u>	<u>8</u>	<u>19</u>	<u>0.42</u>	<u>22</u>	<u>28</u>	<u>0.79</u>
<u>2019</u>	<u>11</u>	<u>21</u>	<u>0.52</u>	<u>23</u>	<u>28</u>	<u>0.82</u>
<u>2020</u>	<u>12</u>	<u>23</u>	<u>0.52</u>	<u>29</u>	<u>29</u>	<u>1.00</u>

167

168 **Table S5.** Combinaison of microsatellite markers used in each PCR mix and type of fluorescent dye  
 169 used for each microsatellite marker from 2017 to 2020.

170

<b>Mix</b>	<b>Locus-name</b>	<b>Dye</b>	<b>Publication</b>
A	UA03	6FAM	De-Barba <i>et al.</i> 2017
A	UA06	6FAM	De-Barba <i>et al.</i> 2017
A	UA25	NED <sup>TM</sup>	De-Barba <i>et al.</i> 2017
A	UA67	NED <sup>TM</sup>	De-Barba <i>et al.</i> 2017
A	UA64	PET <sup>TM</sup>	De-Barba <i>et al.</i> 2017
A	UA63	PET <sup>TM</sup>	De-Barba <i>et al.</i> 2017
A	UA16	VIC <sup>TM</sup>	De-Barba <i>et al.</i> 2017
A	UA14	VIC <sup>TM</sup>	De-Barba <i>et al.</i> 2017
B	UA17	6FAM	De-Barba <i>et al.</i> 2017
B	UA57	6FAM	De-Barba <i>et al.</i> 2017
B	UA51	NED <sup>TM</sup>	De-Barba <i>et al.</i> 2017
B	UA65	PET <sup>TM</sup>	De-Barba <i>et al.</i> 2017
B	UA68	VIC <sup>TM</sup>	De-Barba <i>et al.</i> 2017
B	Our-ZFX	6FAM	Bidon <i>et al.</i> 2013
B	Our-318	6FAM	Bidon <i>et al.</i> 2013
B	Our-SMCY	6FAM	Bidon <i>et al.</i> 2013


171

172 ~~De-Barba, M., Miquel, C., Lobréaux, S., Quenette, P. Y., Swenson, J. E., & Taberlet, P. (2017). High-~~  
 173 ~~throughput microsatellite genotyping in ecology: Improved accuracy, efficiency, standardization and~~  
 174 ~~success with low quantity and degraded DNA. Molecular Ecology Resources, 17(3), 492-507.~~

175 ~~Bidon, T., Frosch, C., Eiken, H. G., Kutschera, V. E., Hagen, S. B., Aarnes, S. G., ... & Hailer, F.~~  
 176 ~~(2013). A sensitive and specific multiplex PCR approach for sex identification of ursine and~~  
 177 ~~tremarettine bears suitable for non-invasive samples. Molecular Ecology Resources, 13(3), 362-368.~~

178

179 **Fig. S1.** Map of the camera traps and transects used in 2020 in France within the framework of the  
180 systematic monitoring of the Pyrenean brown bear population.



181